


PRV

PATENT- OCH REGISTRERINGSVERKET
Patentavdelningen

PCT / SE 2004 / 001960



Intyg Certificate

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

Ansökan ingavs ursprungligen på engelska.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

The application was originally filed in English.

(71) Sökande AstraZeneca AB, Södertälje SE
Applicant (s)

(21) Patentansökningsnummer 0401907-1
Patent application number

(86) Ingivningsdatum 2004-07-21
Date of filing

Stockholm, 2004-11-09

För Patent- och registreringsverket
For the Patent- and Registration Office

Hjordis Segerlund

Avgift
Fee 170:-

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)



BEST AVAILABLE COPY

PATENT- OCH
REGISTRERINGSVERKET
SWEDEN

Postadress/Adress
Box 5055
S-102 42 STOCKHOLM

Telefon/Phone
+46 8 782 25 00
Vx 08-782 25 00

Telex
17978
PATOREG S

Telefax
+46 8 666 02 86
08-666 02 86

CHEMICAL COMPOUNDS

This invention relates to 2-azetidinone derivatives, or pharmaceutically acceptable salts, solvates, solvates of such salts and prodrugs thereof. These 2-azetidinones possess
5 cholesterol absorption inhibitory activity and are accordingly of value in the treatment of disease states associated with hyperlipidaemic conditions. They are therefore useful in methods of treatment of a warm-blooded animal, such as man. The invention also relates to processes for the manufacture of said 2-azetidinone derivatives, to pharmaceutical compositions containing them and to their use in the manufacture of medicaments to inhibit
10 cholesterol absorption in a warm-blooded animal, such as man. A further aspect of this invention relates to the use of the compounds of the invention in the treatment of dyslipidemic conditions.

Atherosclerotic coronary artery disease is a major cause of death and morbidity in the western world as well as a significant drain on healthcare resources. It is well-known that
15 hyperlipidaemic conditions associated with elevated concentrations of total cholesterol and low density lipoprotein (LDL) cholesterol are major risk factors for cardiovascular atherosclerotic disease (for instance "Coronary Heart Disease: Reducing the Risk; a Worldwide View" Assman G., Carmena R. Cullen P. *et al*; Circulation 1999, 100, 1930-1938 and "Diabetes and Cardiovascular Disease: A Statement for Healthcare Professionals from the
20 American Heart Association" Grundy S, Benjamin I., Burke G., *et al*; Circulation, 1999, 100, 1134-46).

The concentration of plasma cholesterol depends on the integrated balance of endogenous and exogenous pathways of cholesterol metabolism. In the endogenous pathway, cholesterol is synthesized by the liver and extra hepatic tissues and enters the circulation as
25 lipoproteins or is secreted into bile. In the exogenous pathway cholesterol from dietary and biliary sources is absorbed in the intestine and enters the circulation as component of chylomicrons. Alteration of either pathway will affect the plasma concentration of cholesterol.

The precise mechanism by which cholesterol is absorbed from the intestine is however not clear. The original hypothesis has been that cholesterol is crossing the intestine by
30 unspecific diffusion. But more recent studies are suggesting that there are specific transporters involved in the intestinal cholesterol absorption. (See for instance New molecular targets for cholesterol-lowering therapy Izzat, N.N., Deshazer, M.E. and Loose-Mitchell D.S. JPET 293:315-320, 2000.)

A clear association between reduction of total cholesterol and (LDL) cholesterol and decreased instance of coronary artery disease has been established, and several classes of pharmaceutical agents are used to control serum cholesterol. There major options to regulate plasma cholesterol include (i) blocking the synthesis of cholesterol by agents such as
5 HMG-CoA reductase inhibitors, for example statins such as simvastatin and fluvastatin, which also by up-regulation of LDL-receptors will promote the cholesterol removal from the plasma; (ii) blocking the bile acid reabsorption by specific agents resulting in increased bile acid excretion and synthesis of bile acids from cholesterol with agents such as bile acid binders, such as resins e.g. cholestyramine and cholestipol; and (iii) by blocking the intestinal
10 uptake of cholesterol by selective cholesterol absorption inhibitors. High density lipoprotein (HDL) elevating agents such as fibrates and nicotinic acid analogues have also been employed.


Even with the current diverse range of therapeutic agents, a significant proportion of the hypercholesterolaemic population is unable to reach target cholesterol levels, or drug
15 interactions or drug safety preclude the long term use needed to reach the target levels. Therefore there is still a need to develop additional agents that are more efficacious and are better tolerated.

Compounds possessing such cholesterol absorption inhibitory activity have been described, see for instance the compounds described in WO 93/02048, WO 94/17038,
20 WO 95/08532, WO 95/26334, WO 95/35277, WO 96/16037, WO 96/19450, WO 97/16455, WO 02/50027, WO 02/50060, WO 02/50068, WO 02/50090, WO 02/66464, WO 04/000803, WO 04/000804, WO04/000805, US 5756470, US 5767115 and US RE37721.

The present invention is based on the discovery that certain 2-azetidinone derivatives surprisingly inhibit cholesterol absorption. Such properties are expected to be of value in the
25 treatment of disease states associated with hyperlipidaemic conditions. The compounds of the present invention are not disclosed in any of the above applications and we have surprisingly found that the compounds of the present invention possess beneficial efficacious, metabolic and toxicological profiles that make them particularly suitable for *in vivo* administration to a warm blooded animal, such as man. In particular certain compounds of the present invention
30 have a low degree of absorption compared to compounds of the prior art whilst retaining their ability to inhibit cholesterol absorption.

Accordingly there is provided a compound of formula (I):

(I)



(I)

5 wherein:

R^1 is hydrogen, C_{1-6} alkyl, C_{3-6} cycloalkyl or aryl; wherein said C_{1-6} alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, carbamoyl, carboxy, C_{1-6} alkoxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, C_1 - C_6 alkylcarbonylamino C_{1-6} alkylS(O) $_a$ wherein a is 0-2, C_{3-6} cycloalkyl or aryl; and wherein any aryl group may be
 10 optionally substituted by one or two substituents selected from halo, hydroxy, C_{1-6} alkyl or C_{1-6} alkoxy;

R^2 is hydrogen, a branched or unbranched C_{1-6} alkyl, C_{3-6} cycloalkyl or aryl; wherein said C_{1-6} alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, carbamoyl, carboxy, C_{1-6} alkoxy, $(C_1-C_4)_3Si$, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino,
 15 C_{1-6} alkylS(O) $_a$ wherein a is 0-2, C_{3-6} cycloalkyl or aryl; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C_{1-6} alkyl or C_{1-6} alkoxy;

R^3 is hydrogen, alkyl, halo or C_{1-6} alkoxy;

∴ R^4 is hydrogen, halo or C_{1-6} alkoxy;

20 R^5 is hydrogen, a branched or unbranched C_{1-6} alkyl, arylalkyl, or aryl C_{1-6} alkyl;

∴ R^6 is hydrogen, C_{1-6} alkyl, or aryl C_{1-6} alkyl;

∴ wherein R^5 and R^2 may form a ring with 2-7 carbon atoms and wherein R^6 and R^2 may form a ring with 3-6 carbon atoms;

∴ or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof;

25 with the proviso that said compound is not 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-2-hydroxyethylsulphanyl]-4-[4-(N -{ N -[(R)-1-(carboxy)-2-(hydroxy)ethyl]carbamoylmethyl} carbamoylmethoxy)phenyl]azetidin-2-one; or 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-

2-hydroxyethylsulphanyl]-4-{4-[N-((R)- α -{N-[(S)-1-(carboxy)-2-(hydroxy)ethyl]carbamoyl}benzyl)carbamoylmethoxy]phenyl}azetidin-2-one.

- In one aspect of the invention R^1 is hydrogen. According to another aspect of the invention,
- 5 R^2 is hydrogen, a branched or unbranched C_{1-6} alkyl, C_{3-6} cycloalkyl or aryl; wherein said C_{1-6} alkyl may be optionally substituted by one or more hydroxy, amino, C_{1-6} alkylS(O)_a wherein a is 0-2, C_{3-6} cycloalkyl or aryl; and wherein any aryl group may be optionally substituted by hydroxy. According to a further aspect of the invention, R^3 is hydrogen, methoxy, or a C_1 - C_6 alkyl, for instance methyl, or a halogen, for instance chlorine or fluorine.
- 10 According to yet another aspect of the invention R^4 is hydrogen or halo, for instance chlorine or fluorine. According to a further aspect of the invention, R^6 is hydrogen, aryl C_{1-6} or R^6 and R^2 form a ring with 3-6 carbon atoms.

- According to one aspect of the invention R^1 is hydrogen, R^2 is a branched or unbranched
- 15 C_{1-4} alkyl, optionally substituted by a C_{3-6} cycloalkyl, R^3 and R^4 are halo, R^5 is hydrogen or C_{1-6} alkyl, and R^6 is hydrogen.

The invention further provides for one or more compounds chosen from:

- 20 N-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-N⁶-acetyl-D-lysine;
- 1-(4-Fluorophenyl)-3-(R)-[2-(4-fluorophenyl)-2-hydroxyethylthio]-4-(R)-{4-[N-{N-[2-(phenyl)-1-(R)-(carboxy)ethyl]carbamoylmethyl}carbamoylmethoxy]phenyl}azetidin-2-one;
- 25 N-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-valine;
- 30 N-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-tyrosine;
- N-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-proline;
- 35 N-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-lysine;
- N-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-hydroxy-2-(4-methoxyphenyl)ethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-valine;
- 40 N-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-2-butylnorleucine;

- 5 *N*-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*S*-methyl-*L*-cysteine;
- N*-{[4-((2*R*,3*R*)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-cyclohexyl-*D*-alanine;
- 10 *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-cyclohexyl-*D*-alanine;
- N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-4-methylleucine;
- 15 *N*-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-*L*-alanyl-*D*-valine;
- N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-hydroxy-2-(4-methylphenyl)ethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*D*-valine;
- 20 *N*-{[4-((2*R*,3*R*)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*D*-valine;
- N*-{[4-((2*R*,3*R*)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-methyl-*D*-valine;
- 25 *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-(2-naphthyl)-*D*-alanine;
- 30 *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-methyl-*D*-valine;
- N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-(3*R*,4*S*,5*R*)-3,4,5,6-tetrahydroxy-*D*-norleucine.
- 35

In this specification the term "alkyl" includes both straight and branched chain alkyl groups but references to individual alkyl groups such as "propyl" are specific for the straight chain version only. For example, "C₁₋₆alkyl" and "C₁₋₄alkyl" include propyl, isopropyl and *t*-butyl. However, references to individual alkyl groups such as 'propyl' are specific for the straight chained version only and references to individual branched chain alkyl groups such as 'isopropyl' are specific for the branched chain version only. A similar convention applies to other radicals, for example "phenylC₁₋₆alkyl" would include benzyl, 1-phenylethyl and 2-phenylethyl. The term "halo" refers to fluoro, chloro, bromo and iodo.

Where optional substituents are chosen from "one or more" groups it is to be understood that this definition includes all substituents being chosen from one of the specified groups or the substituents being chosen from two or more of the specified groups.

The term "aryl" refers to a 4-10 membered aromatic mono or bicyclic ring containing 5 0 to 5 heteroatoms independently selected from nitrogen, oxygen or sulphur. Examples of aryls include phenyl, pyrrolyl, furanyl, imidazolyl, triazolyl, tetrazolyl, pyrazinyl, pyrimidinyl, pyridazinyl, pyridyl, isoxazolyl, oxazolyl, 1,2,4 oxadiazolyl, isothiazolyl, thiazolyl, 1,2,4-triazolyl, thienyl, naphthyl, benzofuranyl, benzimidazolyl, benzthienyl, benzthiazolyl, benzisothiazolyl, benzoxazolyl, benzisoxazolyl, 1,3-benzodioxolyl, indolyl, 10 pyridoimidazolyl, pyrimidoimidazolyl, quinolyl, isoquinolyl, quinoxalyl, quinazolinyl, phthalazinyl, cinnolyl and naphthyridinyl. Particularly "aryl" refers to phenyl, thienyl, pyridyl, imidazolyl or indolyl.

Examples of "C₁₋₆alkoxy" include methoxy, ethoxy and propoxy. Examples of "C₁₋₆alkylS(O)_a wherein a is 0 to 2" include methylthio, ethylthio, methylsulphanyl, 15 ethylsulphanyl, mesyl and ethylsulphonyl. Examples of "N-(C₁₋₆alkyl)amino" include methylamino and ethylamino. Examples of "N,N-(C₁₋₆alkyl)₂amino" include di-N-methylamino, di-(N-ethyl)amino and N-ethyl-N-methylamino. "C₃₋₆cycloalkyl" refers to cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

A suitable pharmaceutically acceptable salt of a compound of the invention, or other 20 compounds disclosed herein, is, for example, an acid-addition salt of a compound of the invention which is sufficiently basic, for example, an acid-addition salt with, for example, an inorganic or organic acid, for example hydrochloric, hydrobromic, sulphuric, phosphoric, trifluoroacetic, citric, acetate or maleic acid. In addition a suitable pharmaceutically acceptable salt of a compound of the invention which is sufficiently acidic is an alkali metal 25 salt, for example a sodium or potassium salt, an alkaline earth metal salt, for example a calcium or magnesium salt, an ammonium salt or a salt with an organic base which affords a physiologically-acceptable cation, for example a salt with methylamine, dimethylamine, trimethylamine, piperidine, morpholine or tris-(2-hydroxyethyl)amine.

The compounds of the formula (I), or other compounds disclosed herein, may be 30 administered in the form of a pro-drug which is broken down in the human or animal body to give a compound of the formula (I). examples of pro-drugs include *in vivo* hydrolysable esters and *in vivo* hydrolysable amides of a compound of the formula (I).

An *in vivo* hydrolysable ester of a compound of the formula (I), or other compounds disclosed herein, containing carboxy or hydroxy group is, for example, a pharmaceutically acceptable ester which is hydrolysed in the human or animal body to produce the parent acid or alcohol. Suitable pharmaceutically acceptable esters for carboxy include C₁₋₆alkoxymethyl esters for example methoxymethyl, C₁₋₆alkanoyloxymethyl esters for example pivaloyloxymethyl, phthalidyl esters, C₃₋₈cycloalkoxycarbonyloxyC₁₋₆alkyl esters for example 1-cyclohexylcarbonyloxyethyl; 1,3-dioxolen-2-onylmethyl esters for example 5-methyl-1,3-dioxolen-2-onylmethyl; and C₁₋₆alkoxycarbonyloxyethyl esters for example 1-methoxycarbonyloxyethyl and may be formed at any carboxy group in the compounds of this invention.

An *in vivo* hydrolysable ester of a compound of the formula (I), or other compounds disclosed herein, containing a hydroxy group includes inorganic esters such as phosphate esters and α -acyloxyalkyl ethers and related compounds which as a result of the *in vivo* hydrolysis of the ester breakdown to give the parent hydroxy group. Examples of α -acyloxyalkyl ethers include acetoxymethoxy and 2,2-dimethylpropionyloxy-methoxy. A selection of *in vivo* hydrolysable ester forming groups for hydroxy include alkanoyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl, alkoxycarbonyl (to give alkyl carbonate esters), dialkylcarbamoyl and *N*-(dialkylaminoethyl)-*N*-alkylcarbamoyl (to give carbamates), dialkylaminoacetyl and carboxyacetyl. Examples of substituents on benzoyl include morpholino and piperazino linked from a ring nitrogen atom via a methylene group to the 3- or 4- position of the benzoyl ring.

A suitable value for an *in vivo* hydrolysable amide of a compound of the formula (I), or other compounds disclosed herein, containing a carboxy group is, for example, a *N*-C₁₋₆alkyl or *N,N*-di-C₁₋₆alkyl amide such as *N*-methyl, *N*-ethyl, *N*-propyl, *N,N*-dimethyl, *N*-ethyl-*N*-methyl or *N,N*-diethyl amide.

Some compounds of the formula (I) may have chiral centres and/or geometric isomeric centres (*E*- and *Z*- isomers), and it is to be understood that the invention encompasses all such optical, diastereoisomers and geometric isomers that possess cholesterol absorption inhibitory activity.

The invention relates to any and all tautomeric forms of the compounds of the formula (I) that possess cholesterol absorption inhibitory activity.


It is also to be understood that certain compounds of the formula (I) can exist in solvated as well as unsolvated forms such as, for example, hydrated forms. It is to be

understood that the invention encompasses all such solvated forms which possess cholesterol absorption inhibitory activity.

Preferred aspects of the invention are those which relate to the compound of formula (I) or a pharmaceutically acceptable salt thereof.


Another aspect of the present invention provides a process for preparing a compound of formula (I) or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof which process (wherein variable groups are, unless otherwise specified, as defined in formula (I)) comprises of:

10 *Process 1*) reacting a compound of formula (II):



(II)


with a compound of formula (III):



(III)

wherein L is a displaceable group;


Process 2) reacting an acid of formula (IV):



(IV)


20 or an activated derivative thereof; with an amine of formula (V):

- 9 -




(V)

Process 3): reacting an acid of formula (VI):




(VI)

or an activated derivative thereof, with an amine of formula (VII):




(VII)

Process 4): reducing a compound of formula (VIII):




(VIII)

Process 5): reacting a compound of formula (IX):



(IX)


with a compound of formula (X):



(X)


wherein L is a displaceable group;

Process 6): reacting a compound of formula (XI):




(XI)

10 wherein L is a displaceable group; with a compound of formula (XII):



(XII)

Process 7): De-esterifying a compound of formula (XIII)



(XIII)

wherein the group C(O)OR is an ester group;

and thereafter if necessary or desirable:

- 5 i) converting a compound of the formula (I) into another compound of the formula (I);
- ii) removing any protecting groups;
- iii) forming a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug; or
- iv) separating two or more enantiomers.

L is a displaceable group, suitable values for L are for example, a halogeno or
 10 sulphonyloxy group, for example a chloro, bromo, methanesulphonyloxy or toluene-4-sulphonyloxy group.


C(O)OR is an ester group, suitable values for C(O)OR are methoxycarbonyl, ethoxycarbonyl, *t*-butoxycarbonyl and benzyloxycarbonyl.

The starting materials used in the present invention can be prepared by modifications
 15 of the routes described in EP 0 792 264 B1. Alternatively they can be prepared by the following reactions.

Process 1): Alcohols of formula (II) may be reacted with compounds of formula (III) in the presence of a base for example an inorganic base such as sodium carbonate, or an organic base such as Hunigs base, in the presence of a suitable solvent such as acetonitrile,

20 dichloromethane or tetrahydrofuran at a temperature in the range of 0°C to reflux, preferably at or near reflux.

Compounds of formula (II) may be prepared according to the following scheme:



Scheme 1

wherein pMeOBz is para methoxy benzyl.


- 5 Compounds of formula (IIb), (IIId), (IIg) and (III) are commercially available compounds, or they are known in the literature, or they are prepared by standard processes known in the art.

- 13 -


A compound of formula (III) may also be reacted with a compound of formula (XIV).

Compounds of formula (XIV) may be prepared according to the following route:

CONFIDENTIAL



Compounds of formula XIVi may be prepared by the following route:



5 *Process 2)* and *Process 3)*: Acids and amines may be coupled together in the presence of a suitable coupling reagent. Standard peptide coupling reagents known in the art can be employed as suitable coupling reagents, for example carbonyldiimidazole and dicyclohexyl-carbodiimide, optionally in the presence of a catalyst such as dimethylaminopyridine or 4-pyrrolidinopyridine, optionally in the presence of a base for
10 example triethylamine, pyridine, or 2,6-di-*alkyl*-pyridines such as 2,6-lutidine or 2,6-di-*tert*-butylpyridine. Suitable solvents include dimethylacetamide, dichloromethane, benzene, tetrahydrofuran and dimethylformamide. The coupling reaction may conveniently be performed at a temperature in the range of -40 to 40°C.

Suitable activated acid derivatives include acid halides, for example acid chlorides,
15 and active esters, for example pentafluorophenyl esters. The reaction of these types of compounds with amines is well known in the art, for example they may be reacted in the presence of a base, such as those described above, and in a suitable solvent, such as those described above. The reaction may conveniently be performed at a temperature in the range of -40 to 40°C.

20 Acids of formula (IV) and (VI) may be prepared from compounds of formula (II) by reacting them with the appropriate, optionally protected, side chain using the conditions of *Process 1)*. Alternatively, acids of formula (IV) and (VI) may be prepared by a modification of *Scheme I*.

Amines of formula (V) and (VII) are commercially available compounds, or they are
25 known in the literature, or they are prepared by standard processes known in the art.

Process 4): Reduction of compounds of formula (VIII) could be performed with a hydride reagent such as sodium borohydride in a solvent such as methanol at temperatures suitable between -20-40°C.

30 Compounds of formula (VIII) can be prepared from compounds of formula (III), by deprotecting the benzyl group and performing *Process 1*. Alternatively compound (IIIk) could be debenzylated, *Process 1* could be performed and the resulting compound deprotected to reveal the ketone.

Process 5) and *Process 6)*: these compounds may be reacted together in the presence of a base for example an inorganic base such as sodium carbonate, or an organic base such as

Hunigs base, in the presence of a suitable solvent such as acetonitrile, dichloromethane or tetrahydrofuran at a temperature in the range of 0°C to reflux, preferably at or near reflux.

Compounds of formula (IX) and (XI) may be prepared by an appropriate modification of *Scheme 1*.

5 Compounds of formula (X) and (XII) are commercially available compounds, or they are known in the literature, or they are prepared by standard processes known in the art.

Process 7): Esters of formula (XIII) may be deprotected under standard conditions such as those described below, for example a methyl or ethyl ester may be deprotected with sodium hydroxide in methanol at room temperature.

10 Compounds of formula (XIII) may be prepared by a modification of any of the processes described herein for the preparation of compounds of formula (I).

It will be appreciated that certain of the various ring substituents in the compounds of the present invention may be introduced by standard aromatic substitution reactions or generated by conventional functional group modifications either prior to or immediately
15 following the processes mentioned above, and as such are included in the process aspect of the invention. Such reactions and modifications include, for example, introduction of a substituent by means of an aromatic substitution reaction, reduction of substituents, alkylation of substituents and oxidation of substituents. The reagents and reaction conditions for such procedures are well known in the chemical art. Particular examples of aromatic substitution
20 reactions include the introduction of a nitro group using concentrated nitric acid, the introduction of an acyl group using, for example, an acyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; the introduction of an alkyl group using an alkyl halide and Lewis acid (such as aluminium trichloride) under Friedel Crafts conditions; and the introduction of a halogeno group. Particular examples of modifications
25 include the reduction of a nitro group to an amino group by for example, catalytic hydrogenation with a nickel catalyst or treatment with iron in the presence of hydrochloric acid with heating; oxidation of alkylthio to alkylsulphinyl or alkylsulphonyl.

It will also be appreciated that in some of the reactions mentioned herein it may be necessary/desirable to protect any sensitive groups in the compounds. The instances where
30 protection is necessary or desirable and suitable methods for protection are known to those skilled in the art. Conventional protecting groups may be used in accordance with standard practice (for illustration see T.W. Green, *Protective Groups in Organic Synthesis*, John Wiley

and Sons, 1999). Thus, if reactants include groups such as amino, carboxy or hydroxy it may be desirable to protect the group in some of the reactions mentioned herein.

A suitable protecting group for an amino or alkylamino group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an alkoxycarbonyl group, for example a methoxycarbonyl, ethoxycarbonyl or *t*-butoxycarbonyl group, an arylmethoxycarbonyl group, for example benzyloxycarbonyl, or an aroyl group, for example benzoyl. The deprotection conditions for the above protecting groups necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or alkoxycarbonyl group or an aroyl group may be removed for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an acyl group such as a *t*-butoxycarbonyl group may be removed, for example, by treatment with a suitable acid as hydrochloric, sulphuric or phosphoric acid or trifluoroacetic acid and an arylmethoxycarbonyl group such as a benzyloxycarbonyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon, or by treatment with a Lewis acid for example boron tris(trifluoroacetate). A suitable alternative protecting group for a primary amino group is, for example, a phthaloyl group which may be removed by treatment with an alkylamine, for example dimethylaminopropylamine, or with hydrazine.

A suitable protecting group for a hydroxy group is, for example, an acyl group, for example an alkanoyl group such as acetyl, an aroyl group, for example benzoyl, or an arylmethyl group, for example benzyl. The deprotection conditions for the above protecting groups will necessarily vary with the choice of protecting group. Thus, for example, an acyl group such as an alkanoyl or an aroyl group may be removed, for example, by hydrolysis with a suitable base such as an alkali metal hydroxide, for example lithium or sodium hydroxide. Alternatively an arylmethyl group such as a benzyl group may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

A suitable protecting group for a carboxy group is, for example, an esterifying group, for example a methyl or an ethyl group which may be removed, for example, by hydrolysis with a base such as sodium hydroxide, or for example a *t*-butyl group which may be removed, for example, by treatment with an acid, for example an organic acid such as trifluoroacetic acid, or for example a benzyl group which may be removed, for example, by hydrogenation over a catalyst such as palladium-on-carbon.

The protecting groups may be removed at any convenient stage in the synthesis using conventional techniques well known in the chemical art.

As stated hereinbefore the compounds defined in the present invention possess cholesterol absorption inhibitory activity. These properties may be assessed, using the following biological tests.

In vivo testing of cholesterol absorption inhibitors (A)

5 C57BL/6 female mice were maintained on regular chow diet and housed in individual cages to collect faeces. Mice were fasted for 3 hours and then gavaged with vehicle or compound. Half an hour later the mice were gavaged with radiolabelled cholesterol. Six hours after the ^{14}C -cholesterol gavage blood samples were taken via the tail and plasma prepared to determine how much cholesterol were absorbed. 24 hours after the gavage of ^{14}C -cholesterol
10 the mice were bled and plasma were prepared for analysis. Faeces were collected for 24 hours to assess absorption efficiency.

In vivo testing of cholesterol absorption inhibitors (B).

C57BL/6 female mice were maintained on regular chow diet and housed in individual cages to collect faeces. Mice were fasted for 3 hours and then gavaged with vehicle or
15 compound. One to ten hours later the mice were gavaged with radiolabelled cholesterol. Six hours after the ^{14}C -cholesterol gavage blood sample was taken via the tail and plasma prepared to determine how much cholesterol was absorbed. 24 hours after the gavage of ^{14}C -cholesterol the mice were bled and plasma analysed for radioactivity. Faeces were also collected for 24 hours to assess absorption efficiency.

20 References

1. E. A. Kirk, G. L. Moe, M. T. Caldwell, J. Å. Lernmark, D. L. Wilson, R. C. LeBoeuf. Hyper- and hypo-responsiveness to dietary fat and cholesterol among inbred mice: searching for level and variability genes. J. Lipid Res. 1995 36:1522-1532.
2. C. P. Carter, P. N. Howles, D. Y. Hui. Genetic variation in cholesterol absorption
25 efficiency among inbred strains of mice. J. Nutr. 1997 127:1344-1348.
3. C. D. Jolley, J. M. Dietschy, S. D. Turley. Genetic differences in cholesterol absorption in 129/Sv and C57BL/6 mice: effect on cholesterol responsiveness. Am. J. Physiol. 1999 276:G1117-G1124.

Administration of 5 $\mu\text{mol/kg}$ of Example 3 gave 87% inhibition of ^{14}C -cholesterol
30 absorption (procedure A). Administration of 5 $\mu\text{mol/kg}$ of Example 1 gave 89% inhibition of ^{14}C -cholesterol absorption (procedure A).

According to a further aspect of the invention there is provided a pharmaceutical

composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore in association with a pharmaceutically-acceptable diluent or carrier.

The composition may be in a form suitable for oral administration, for example as a
5 tablet or capsule, for parenteral injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion) as a sterile solution, suspension or emulsion, for topical administration as an ointment or cream or for rectal administration as a suppository.

In general the above compositions may be prepared in a conventional manner using conventional excipients.

10 The compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, will normally be administered to a warm-blooded animal at a unit dose within the range of approximately 0.02-100 mg/kg, preferably 0.02 -50 mg/kg, and this normally provides a therapeutically-effective dose. A unit dose form such as a tablet
15 or capsule will usually contain, for example 1-250 mg of active ingredient. Preferably a daily dose in the range of 1-50 mg/kg, particularly 0.1-10 mg/kg is employed. In another aspect a daily dose in the range of 0.01-20 mg/kg is employed. In one aspect of the invention the daily dose of a compound of formula (I) is less than or equal to 100mg. However the daily dose will necessarily be varied depending upon the host treated, the particular route of
20 administration, and the severity of the illness being treated. Accordingly the optimum dosage may be determined by the practitioner who is treating any particular patient.

According to a further aspect of the present invention there is provided a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore for use in a method of prophylactic or therapeutic treatment of a warm-blooded animal, such as man.

25 We have found that the compounds defined in the present invention, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, are effective cholesterol absorption inhibitors, and accordingly have value in the treatment of disease states associated with hyperlipidaemic conditions.

Thus according to this aspect of the invention there is provided a compound of the
30 formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore for use as a medicament.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a

prodrug thereof, as defined hereinbefore in the manufacture of a medicament for use in the production of a cholesterol absorption inhibitory effect in a warm-blooded animal, such as man.

According to another feature of the invention there is provided the use of a compound
5 of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore in the production of a cholesterol absorption inhibitory effect in a warm-blooded animal, such as man.

Herein, where the production of a cholesterol absorption inhibitory effect or a cholesterol lowering effect is stated, suitably this relates to the treatment of hyperlipidaemic
10 conditions in a warm-blooded animal, such as man. Additionally is relates to the treatment of dyslipidemic conditions and disorders such as hyperlipidaemia, hypertriglyceridemia, hyperbetalipoproteinemia (high LDL), hyperprebetalipoproteinemia (high VLDL), hyperchylomicronemia, hypolipoproteinemia, hypercholesterolemia, hyperlipoproteinemia and hypoalphalipoproteinemia (low HDL) in a warm-blooded animal, such as man.
15 Furthermore it relates to the treatment of different clinical conditions such as atherosclerosis, arteriosclerosis, arrhythmia, hyper-thrombotic conditions, vascular dysfunction, endothelial dysfunction, heart failure, coronary heart diseases, cardiovascular diseases, myocardial infarction, angina pectoris, peripheral vascular diseases, inflammation of cardiovascular tissues such as heart, valves, vasculature, arteries and veins, aneurisms, stenosis, restenosis,
20 vascular plaques, vascular fatty streaks, leukocytes, monocytes and/or macrophage infiltration, intimal thickening, medial thinning, infectious and surgical trauma and vascular thrombosis, stroke and transient ischaemic attacks in a warm-blooded animal, such as man. It also relates to the treatment of atherosclerosis, coronary heart diseases, myocardial infarction, angina pectoris, peripheral vascular diseases, stroke and transient ischaemic attacks in a
25 warm-blooded animal, such as man.

The production of a cholesterol absorption inhibitory effect or a cholesterol lowering effect also relates to a method of treating and/or preventing atherosclerotic lesions, a method of preventing plaque rupture and a method of promoting lesion regression. Furthermore it relates to a method of inhibiting monocytes-macrophage accumulation in atherosclerotic
30 lesions, a method of inhibiting expression of matrix metalloproteinases in atherosclerotic lesions, a method of inhibiting the destabilization of atherosclerotic lesions, a method for preventing atherosclerotic plaque rupture and a method of treating unstable angina.

The production of a cholesterol absorption inhibitory effect or a cholesterol lowering effect also relates to a method of treating sitosterolemia.

Compounds of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof may also have value in the treatment or prevention of

5 Alzheimer's Disease (see for example WO 02/096415). Therefore in a further aspect of the invention, there is provided a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, for use in the treatment or prevention of Alzheimer's Disease.

Compounds of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of

10 such a salt or a prodrug thereof may also have value in the treatment or prevention of vascular inflammation (see for example WO 03/026644). Therefore in a further aspect of the invention, there is provided a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, for use in the treatment or prevention of vascular inflammation.

15 According to a further feature of this aspect of the invention there is provided a method for producing a cholesterol absorption inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

20 The cholesterol absorption inhibitory activity defined hereinbefore may be applied as a sole therapy or may involve, in addition to a compound of the invention, one or more other substances and/or treatments. Such conjoint treatment may be achieved by way of the simultaneous, sequential or separate administration of the individual components of the treatment. According to this aspect of the invention there is provided a pharmaceutical

25 product comprising a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, as defined hereinbefore and an additional cholesterol absorption inhibitory substance as defined hereinbefore and an additional hypolipidaemic agent for the conjoint treatment of hyperlipidaemia.

In another aspect of the invention, the compound of formula (I), or a pharmaceutically

30 acceptable salt, solvate, solvate of such a salt or a prodrug thereof, may be administered in association with cholesterol biosynthesis inhibitors, or pharmaceutically acceptable salts, solvates, solvates of such salts or prodrugs thereof. Suitable cholesterol biosynthesis inhibitors include HMG Co-A reductase inhibitors, squalene synthesis inhibitors and squalene

epoxidase inhibitors. A suitable squalene synthesis inhibitor is squalestatin 1 and a suitable squalene epoxidase inhibitor is NB-598.

In this aspect of the invention, the compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, may be administered in association with an HMG Co-A reductase inhibitor, or pharmaceutically acceptable salts, solvates, solvates of such salts or prodrugs thereof. Suitable HMG Co-A reductase inhibitors, pharmaceutically acceptable salts, solvates, solvates of such salts or prodrugs thereof are statins well known in the art. Particular statins are fluvastatin, lovastatin, pravastatin, simvastatin, atorvastatin, cerivastatin, bervastatin, dalvastatin, mevastatin and rosuvastatin, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof. A further particular statin is pitavastatin, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof. A particular statin is atorvastatin, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof. A more particular statin is atorvastatin calcium salt. A further particular statin is rosuvastatin, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof. A preferable particular statin is rosuvastatin calcium salt.

Therefore in an additional feature of the invention, there is provided a combination of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

Therefore in an additional feature of the invention, there is provided a method for producing a cholesterol lowering effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in association with a pharmaceutically acceptable diluent or carrier.

According to a further aspect of the present invention there is provided a kit comprising a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

5 According to a further aspect of the present invention there is provided a kit comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a first unit dosage form;
- b) an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate
- 10 of such a salt or a prodrug thereof; in a second unit dosage form; and
- c) container means for containing said first and second dosage forms.

According to a further aspect of the present invention there is provided a kit comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a
- 15 salt or a prodrug thereof, together with a pharmaceutically acceptable diluent or carrier, in a first unit dosage form;
- b) an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a second unit dosage form; and
- c) container means for containing said first and second dosage forms.

20 According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and an HMG Co-A reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the production of a cholesterol lowering effect.

25 According to a further aspect of the present invention there is provided a combination treatment comprising the administration of an effective amount of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with the simultaneous, sequential or separate administration of an effective amount of an HMG Co-A

30 reductase inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier to a warm-blooded animal, such as man in need of such therapeutic treatment.

According to an additional further aspect of the present invention there is provided a combination treatment comprising the administration of an effective amount of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with the simultaneous, sequential or separate administration of a matrix metalloproteinase inhibitor.


In another aspect of the invention, the compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, may be administered in association with an ileal bile acid (IBAT) inhibitor or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof. Suitable compounds possessing IBAT inhibitory activity for use in combination with compounds of the present invention have been described, see for instance the compounds described in WO 93/16055, WO 94/18183, WO 94/18184, WO 94/24087, WO 96/05188, WO 96/08484, WO 96/16051, WO 97/33882, WO 98/07749, WO 98/38182, WO 98/40375, WO 98/56757, WO 99/32478, WO 99/35135, WO 99/64409, WO 99/64410, WO 00/01687, WO 00/20392, WO 00/20393, WO 00/20410, WO 00/20437, WO 00/35889, WO 01/34570, WO 00/38725, WO 00/38726, WO 00/38727, WO 00/38728, WO 00/38729, WO 00/47568, WO 00/61568, WO 01/66533, WO 01/68096, WO 01/68637, WO 02/08211, DE 19825804, JP 10072371, US 5070103, EP 251 315, EP 417 725, EP 489 423, EP 549 967, EP 573 848, EP 624 593, EP 624 594, EP 624 595, EP 864 582, EP 869 121 and EP 1 070 703 and the contents of these patent applications are incorporated herein by reference. Particularly the named examples of these patent applications are incorporated herein by reference. More particularly claim 1 of these patent application are incorporated herein by reference.

Other suitable classes of IBAT inhibitors for use in combination with compounds of the present invention are the 1,2-benzothiazepines, 1,4-benzothiazepines and 1,5-benzothiazepines. A further suitable class of IBAT inhibitors is the 1,2,5-benzothiadiazepines.

One particular suitable compound possessing IBAT inhibitory activity for use in combination with compounds of the present invention is (3R,5R)-3-butyl-3-ethyl-1,1-dioxido-5-phenyl-2,3,4,5-tetrahydro-1,4-benzothiazepin-8-yl β -D-glucopyranosiduronic acid (EP 864 582).

A further suitable compound possessing IBAT inhibitory activity for use in combination with compounds of the present invention is S-8921 (EP 597 107).

A further suitable IBAT inhibitor for use in combination with compounds of the present invention is the compound:



WO 99/32478

- 5 A particular IBAT inhibitor for use in combination with compounds of the present invention is selected from any one of Examples 1-120 of WO 02/50051, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and the compounds of Examples 1-120 are incorporated herein by reference. Claims 1-15 of WO 02/50051 are also incorporated herein by reference. A particular IBAT inhibitor selected from WO 02/50051 for
- 10 use in combination with compounds of the present invention is selected from any one of:
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)-1'-phenyl-1'-[*N'*-(carboxymethyl)carbamoyl]methyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(carboxymethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 15 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)-1'-phenyl-1'-[*N'*-(2-sulphoethyl)carbamoyl]methyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)-1'-phenyl-1'-[*N'*-(2-sulphoethyl)carbamoyl]methyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 20 1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

- 1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-carboxyethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 5 1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(5-carboxypentyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{ α -[*N'*-(2-sulphoethyl)carbamoyl]-2-fluorobenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 10 1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(R)-(2-hydroxy-1-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(R)-(2-hydroxy-1-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 15 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-{*N*-[(R)- α -(*N'*-{(R)-1-[*N''*-(R)-(2-hydroxy-1-carboxyethyl)carbamoyl]-2-hydroxyethyl}carbamoyl)benzyl]carbamoylmethoxy}-2,3,4,5-tetrahydro-1,5-benzothiazepine;
1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{ α -[*N'*-(carboxymethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 20 1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{ α -[*N'*-((ethoxy)(methyl)phosphorylmethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-{*N*-[(R)- α -(*N'*-{2-[(hydroxy)(methyl)phosphoryl]ethyl}carbamoyl)benzyl]carbamoylmethoxy}-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 25 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-methylthio-1-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-{*N*-[(R)- α -(*N'*-{2-[(methyl)(ethyl)phosphoryl]ethyl}carbamoyl)-4-hydroxybenzyl]carbamoylmethoxy}-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 30 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-{*N*-[(R)- α -(*N'*-{2-[(methyl)(hydroxy)phosphoryl]ethyl}carbamoyl)-4-hydroxybenzyl]carbamoylmethoxy}-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[(R)-*N'*-(2-methylsulphinyl)-1-carboxyethyl]carbamoyl}benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
and

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methoxy-8-[*N*-{(R)- α -[*N'*-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy]-2,3,4,5-tetrahydro-1,5-benzothiazepine;
or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

A particular IBAT inhibitor for use in combination with compounds of the present invention is selected from any one of Examples 1-44 of WO 03/020710, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and the compounds of

10 Examples 1-44 are incorporated herein by reference. Claims 1-10 of WO 03/020710 are also incorporated herein by reference. A particular IBAT inhibitor selected from WO 03/020710 for use in combination with compounds of the present invention is selected from any one of:

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-

15 benzothiazepine;

1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-(S)-1-carbamoyl-2-hydroxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(hydroxycarbonylmethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-[*N*-{(R)- α -[*N'*-(2-(*N'*-pyrimidin-2-ylureido)ethyl)carbamoyl}benzyl}carbamoylmethoxy]-2,3,4,5-tetrahydro-1,5-

25 benzothiazepine;

1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-[*N*-{(R)- α -[*N'*-(2-(*N'*-pyridin-2-ylureido)ethyl)carbamoyl}benzyl}carbamoylmethoxy]-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(1-*t*-

30 butoxycarbonylpiperidin-4-ylmethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2,3-dihydroxypropyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-[*N*-{(R)- α -{*N'*-[2-(3,4-dihydroxyphenyl)-2-methoxyethyl]carbamoyl}benzyl}carbamoylmethoxy]-2,3,4,5-tetrahydro-1,5-benzothiazepine

1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-aminoethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(piperidin-4-ylmethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine; or
1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2,*N,N*-dimethylaminosulphamoyl)ethyl]carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

15 A particular IBAT inhibitor for use in combination with compounds of the present invention is selected from any one of Examples 1-7 of WO 03/022825, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and the compounds of Examples 1-7 are incorporated herein by reference. Claims 1-8 of WO 03/022825 are also incorporated herein by reference. A particular IBAT inhibitor selected from WO 03/022825
20 for use in combination with compounds of the present invention is selected from any one of:

1,1-dioxo-3(R)-3-butyl-3-ethyl-5-(R)-5-phenyl-8-[*N*-{(R)- α -carboxybenzyl}carbamoylmethoxy]-2,3,4,5-tetrahydro-1,4-benzothiazepine;

1,1-dioxo-3(S)-3-butyl-3-ethyl-5-(S)-5-phenyl-8-[*N*-{(R)- α -carboxybenzyl}carbamoylmethoxy]-2,3,4,5-tetrahydro-1,4-benzothiazepine;

25 1,1-dioxo-3(R)-3-butyl-3-ethyl-5-(R)-5-phenyl-8-(*N*-{(R)- α -[*N*-(carboxymethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine;

1,1-dioxo-3(S)-3-butyl-3-ethyl-5-(S)-5-phenyl-8-(*N*-{(R)- α -[*N*-(carboxymethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine;

3,5-*trans*-1,1-dioxo-3-ethyl-3-butyl-5-phenyl-7-bromo-8-(*N*-{(R)- α -[*N*-(carboxymethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine;

30 (carboxymethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine;

3,5-*trans*-1,1-dioxo-3-(S)-3-ethyl-3-butyl-4-hydroxy-5-(S)-5-phenyl-7-bromo-8-(N-{(R)- α -[N-(carboxymethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine

3,5-*trans*-1,1-dioxo-3-(R)-3-ethyl-3-butyl-4-hydroxy-5-(R)-5-phenyl-7-bromo-8-(N-{(R)- α -[N-(carboxymethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine;

3,5-*trans*-1,1-dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-(N-{(R)- α -[N-(carboxymethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine;

10 3,5-*trans*-1,1-dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-(N-{(R)- α -[N-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine ammonia salt;

1,1-dioxo-3-(S)-3-ethyl-3-butyl-5-(S)-5-phenyl-7-methylthio-8-(N-{(R)- α -[N-(carboxymethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-

15 benzothiazepine diethylamine salt; and

1,1-dioxo-3-(R)-3-ethyl-3-butyl-5-(R)-5-phenyl-7-methylthio-8-(N-{(R)- α -[N-(carboxymethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine diethylamine salt;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

20 A particular IBAT inhibitor for use in combination with compounds of the present invention is selected from any one of Examples 1-4 of WO 03/022830, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and the compounds of Examples 1-4 are incorporated herein by reference. Claims 1-8 of WO 03/022830 are also incorporated herein by reference. A particular IBAT inhibitor selected from WO 03/022830

25 for use in combination with compounds of the present invention is selected from any one of:

1,1-dioxo-3-butyl-3-ethyl-4-hydroxy-5-phenyl-7-(N-{(R)- α -[N-(carboxymethyl)carbamoyl]benzyl}carbamoylmethylthio)-2,3,4,5-tetrahydrobenzothiepine

1,1-dioxo-3-butyl-3-ethyl-4-hydroxy-5-phenyl-7-(N-{(R)- α -[N-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethylthio)-2,3,4,5-tetrahydrobenzothiepine ammonia salt

30 1,1-dioxo-3-butyl-3-ethyl-4-hydroxy-5-phenyl-7-{N-[α -(carboxy)-2-fluorobenzyl]carbamoylmethylthio}-2,3,4,5-tetrahydrobenzothiepine; and
1,1-dioxo-3-butyl-3-ethyl-4-hydroxy-5-phenyl-7-{N-[1-(carboxy)-1-(thien-2-yl)methyl]carbamoylmethylthio}-2,3,4,5-tetrahydrobenzothiepine

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

A particular IBAT inhibitor for use in combination with compounds of the present invention is selected from any one of Examples 1-39 of WO 03/022286, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and the compounds of

5 Examples 1-39 are incorporated herein by reference. Claims 1-10 of WO 03/022286 are also incorporated herein by reference. A particular IBAT inhibitor selected from WO 03/022286 for use in combination with compounds of the present invention is selected from any one of:

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((R)-1-carboxy-2-methylthio-ethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-

10 benzothiadiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxy-2-(R)-hydroxypropyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxy-2-

15 methylpropyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxybutyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

20 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxypropyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxy-2-(R)-hydroxypropyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-

25 benzothiadiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-

30 carboxyethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(N-{(R)- α -[N-((R)-1-carboxy-2-methylthioethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(N-{(R)- α -[N-((S)-1-[N-((S)-2-hydroxy-1-carboxyethyl)carbamoyl]propyl}carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(N-{(R)- α -[N-((S)-1-carboxy-2-methylpropyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

10 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(N-{(R)- α -[N-((S)-1-carboxypropyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine; and

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-[N-((R)- α -carboxy-4-hydroxybenzyl)carbamoylmethoxy]-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

15 or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

A particular IBAT inhibitor for use in combination with compounds of the present invention is selected from any one of Examples 1-7 of WO 03/091232, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and the compounds of Examples 1-7 are incorporated herein by reference. Claims 1-10 of WO 03/091232 are also
20 incorporated herein by reference. A particular IBAT inhibitor selected from WO 03/091232 for use in combination with compounds of the present invention is selected from any one of:

1,1-Dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(N-{(R)- α -[N-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

25 1,1-Dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(N-{(R)- α -[N-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

1,1-Dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-[N-((R/S)- α -{N-[1-(R)-2-(S)-1-hydroxy-1-(3,4-dihydroxyphenyl)prop-2-yl]carbamoyl}-4-hydroxybenzyl)carbamoylmethoxy]-2,3,4,5-
30 tetrahydro-1,2,5-benzothiadiazepine;

1,1-Dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-{N-[(R)- α -(N-{2-(S)-[N-(carbamoylmethyl)carbamoyl]pyrrolidin-1-ylcarbonylmethyl}carbamoyl)benzyl}carbamoylmethoxy}-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;


1,1-Dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-[N-((R)- α -{N-[2-(3,4,5-trihydroxyphenyl)ethyl]carbamoyl}benzyl)carbamoylmethoxy]-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine; and

1,1-Dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(N-{(R)- α -[N-(2-(R)-3-(S)-4-(S)-5-(R)-3,4,5,6-tetrahydroxytetrahydropyran-2-ylmethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

Further suitable compounds possessing IBAT inhibitory activity for use in combination with compounds of the present invention have the following structure of formula

10 (AI):



(AI)

wherein:

R¹ and R² are independently selected from C₁₋₄alkyl;

15 R³ is hydrogen, hydroxy or halo;

R⁴ is C₁₋₄alkyl optionally substituted by hydroxy, methoxy and methylS(O)_a wherein a is 0-2

R⁵ is hydroxy or HOC(O)CH(R⁶)NH-;

20 R⁶ is selected from hydrogen and C₁₋₃alkyl optionally substituted by hydroxy, methoxy and methylS(O)_a wherein a is 0-2;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof;

with the proviso that when R¹ and R² are both butyl, R⁵ is hydroxy and R⁴ is

methylthiomethyl, methylsulphinylmethyl, methylthiomethyl, hydroxymethyl,

methoxymethyl; R³ is not hydrogen; and with the proviso that when R¹ and R² are both butyl,

25 R⁵ is HOC(O)CH(R⁶)NH-, R⁶ is hydroxymethyl and R⁴ is hydroxymethyl; R³ is not hydrogen.

Suitable IBAT inhibitors having the above structure for use in combination with compounds of the present invention are selected from any one of:

- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxyethyl) carbamoyl]benzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 5 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxypropyl) carbamoyl]benzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxybutyl) carbamoyl]benzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 10 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-2-methylpropyl) carbamoyl]benzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-2-methylbutyl) carbamoyl]benzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-3-methylbutyl) carbamoyl]benzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 15 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-2-hydroxypropyl) carbamoyl]benzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-2-mesyloethyl) carbamoyl]benzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 20 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-3-methylsulphonylpropyl) carbamoyl]benzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-3-mesylpropyl) carbamoyl]benzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 25 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxyethyl) carbamoyl]-4-hydroxybenzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxypropyl) carbamoyl]-4-hydroxybenzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxybutyl) carbamoyl]-4-hydroxybenzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 30 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-2-methylpropyl) carbamoyl]-4-hydroxybenzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-

benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-2-methylbutyl)carbamoyl]-4-hydroxybenzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

5 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-3-methylbutyl)carbamoyl]-4-hydroxybenzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-2-hydroxyethyl)carbamoyl]-4-hydroxybenzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-

10 benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-2-hydroxypropyl)carbamoyl]-4-hydroxybenzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-2-

15 methylthioethyl)carbamoyl]-4-hydroxybenzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-2-methylsulphinyethyl)carbamoyl]-4-hydroxybenzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

20 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-2-mesylethyl)carbamoyl]-4-hydroxybenzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-2-methoxyethyl)carbamoyl]-4-hydroxybenzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-

25 benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-3-methylthiopropyl)carbamoyl]-4-hydroxybenzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-3-

30 methylsulphonylpropyl)carbamoyl]-4-hydroxybenzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxy-3-mesylpropyl)carbamoyl]-4-hydroxybenzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-


benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxypropyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine; or

- 5 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(*R*)- α -[*N'*-((*S*)-1-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine.

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

Further suitable IBAT inhibitors for use in combination with compounds of the present invention are those having the structure (BI):



(BI)

wherein

M^1 is $-CH_2-$ or $-NR^{21}-$;


M^2 is $-CR^{22}R^{23}-$ or $-NR^{24}-$; provided that if M^1 is $-NR^{21}-$, M^2 is $-CR^{22}R^{23}-$;

- 15 One of R^1 and R^2 are selected from hydrogen, C_{1-6} alkyl or C_{2-6} alkenyl and the other is selected from C_{1-6} alkyl or C_{2-6} alkenyl;

R^3 is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkanoylamino, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl) $_2$ carbamoyl, C_{1-6} alkylS(O) $_a$ wherein a is 0 to 2, C_{1-6} alkoxycarbonyl, N -(C_{1-6} alkyl)sulphamoyl and N,N -(C_{1-6} alkyl) $_2$ sulphamoyl;

v is 0-5;

one of R^5 and R^6 is a group of formula (BIA):



(BIA)

R⁴ and R⁷ and the other of R⁵ and R⁶ are independently selected from hydrogen, halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl, mercapto, sulphamoyl, C₁₋₄alkyl,

5 C₂₋₄alkenyl, C₂₋₄alkynyl, C₁₋₄alkoxy, C₁₋₄alkanoyl, C₁₋₄alkanoyloxy, *N*-(C₁₋₄alkyl)amino, *N,N*-(C₁₋₄alkyl)₂amino, C₁₋₄alkanoylamino, *N*-(C₁₋₄alkyl)carbamoyl,

N,N-(C₁₋₄alkyl)₂carbamoyl, C₁₋₄alkylS(O)_a wherein a is 0 to 2, C₁₋₄alkoxycarbonyl,

N-(C₁₋₄alkyl)sulphamoyl and *N,N*-(C₁₋₄alkyl)₂sulphamoyl; wherein R⁴ and R⁷ and the other of R⁵ and R⁶ may be optionally substituted on carbon by one or more R²⁵;

10 Z is -O-, -N(R^a)-, -S(O)_b- or -CH(R^a)-; wherein R^a is hydrogen or C₁₋₆alkyl and b is 0-2;

R⁸ is hydrogen, C₁₋₄alkyl, carbocyclyl or heterocyclyl; wherein R⁸ may be optionally substituted on carbon by one or more substituents selected from R²⁶; and wherein if said heterocyclyl contains an -NH- group, that nitrogen may be optionally substituted by a group
15 selected from R²⁷;


R⁹ is hydrogen or C₁₋₄alkyl;

R¹⁰ and R¹¹ are independently selected from hydrogen, C₁₋₄alkyl, carbocyclyl or heterocyclyl; or R¹⁰ and R¹¹ together form C₂₋₆alkylene; wherein R¹⁰ and R¹¹ or R¹⁰ and R¹¹ together may be independently optionally substituted on carbon by one or more substituents
20 selected from R²⁸; and wherein if said heterocyclyl contains an -NH- moiety, that nitrogen may be optionally substituted by one or more R²⁹;

R¹² is hydrogen, C₁₋₄alkyl, carbocyclyl or heterocyclyl; wherein R¹² may be optionally substituted on carbon by one or more substituents selected from R³⁰; and wherein if said heterocyclyl contains an -NH- moiety, that nitrogen may be optionally substituted by one or
25 more R³¹;

R¹³ is hydrogen, halo, nitro, cyano, hydroxy, amino, carbamoyl, mercapto, sulphamoyl, hydroxyaminocarbonyl, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl, C₁₋₁₀alkoxy, C₁₋₁₀alkoxycarbonyl, C₁₋₁₀alkanoyl, C₁₋₁₀alkanoyloxy, *N*-(C₁₋₁₀alkyl)amino, *N,N*-(C₁₋₁₀alkyl)₂amino, *N,N,N*-(C₁₋₁₀alkyl)₃ammonio, C₁₋₁₀alkanoylamino,
30 *N*-(C₁₋₁₀alkyl)carbamoyl, *N,N*-(C₁₋₁₀alkyl)₂carbamoyl, C₁₋₁₀alkylS(O)_a wherein a is 0 to 2, *N*-(C₁₋₁₀alkyl)sulphamoyl, *N,N*-(C₁₋₁₀alkyl)₂sulphamoyl, *N*-(C₁₋₁₀alkyl)sulphamoylamino,

N,N-(C₁₋₁₀alkyl)₂sulphamoylamino, C₁₋₁₀alkoxycarbonylamino, carbocyclyl, carbocyclylC₁₋₁₀alkyl, heterocyclic group, heterocyclylC₁₋₁₀alkyl, carbocyclyl-(C₁₋₁₀alkylene)_e-R³²-(C₁₋₁₀alkylene)_f or heterocyclyl-(C₁₋₁₀alkylene)_g-R³³-(C₁₋₁₀alkylene)_h; wherein R¹³ may be optionally substituted on carbon by one or more substituents selected from R³⁶; and wherein if said heterocyclyl contains an -NH- group, that nitrogen may be optionally substituted by a group selected from R³⁷; or R¹³ is a group of formula (BIB):



(BIB)

10 wherein:

X is $-N(R^{38})-$, $-N(R^{38})C(O)-$, $-O-$, and $-S(O)_a-$; wherein a is 0-2 and R^{38} is hydrogen or C_{1-4} alkyl;

R¹⁴ is hydrogen or C₁₋₄alkyl;

R¹⁵ and **R¹⁶** are independently selected from hydrogen, halo, nitro, cyano, hydroxy,

15 amino, carbamoyl, mercapto, sulphamoyl, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆alkoxy, C₁₋₆alkanoyl, C₁₋₆alkanoyloxy, *N*-(C₁₋₆alkyl)amino, *N,N*-(C₁₋₆alkyl)₂amino,

C₁₋₆alkanoylamino, *N*-(C₁₋₆alkyl)carbamoyl, *N,N*-(C₁₋₆alkyl)₂carbamoyl, C₁₋₆alkylS(O)_a wherein a is 0 to 2, C₁₋₆alkoxycarbonyl, *N*-(C₁₋₆alkyl)sulphamoyl,

N,N-(C₁₋₆alkyl)₂sulphamoyl, carbocyclyl or heterocyclic group; wherein R¹⁵ and R¹⁶ may be

20 independently optionally substituted on carbon by one or more substituents selected from R⁴¹;
and wherein if said heterocyclyl contains an -NH- group, that nitrogen may be optionally
substituted by a group selected from R⁴²;

R¹⁷ is selected from hydrogen, halo, nitro, cyano, hydroxy, amino, carbamoyl, mercapto, sulphamoyl, hydroxyaminocarbonyl, C₁₋₁₀alkyl, C₂₋₁₀alkenyl, C₂₋₁₀alkynyl,

25 C₁₋₁₀alkoxy, C₁₋₁₀alkanoyl, C₁₋₁₀alkanoyloxy, *N*-(C₁₋₁₀alkyl)amino, *N,N*-(C₁₋₁₀alkyl)₂amino, C₁₋₁₀alkanoylamino, *N*-(C₁₋₁₀alkyl)carbamoyl, C₁₋₁₀alkoxycarbonyl,

N,N-(C₁₋₁₀alkyl)₂carbamoyl, C₁₋₁₀alkylS(O)_a wherein a is 0 to 2, *N*-(C₁₋₁₀alkyl)sulphamoyl,


N,N-(C₁₋₁₀alkyl)₂sulphamoyl, *N*-(C₁₋₁₀alkyl)sulphamoylamino,

N,N-(C₁₋₁₀alkyl)₂sulphamoylamino, carbocyclyl, carbocyclylC₁₋₁₀alkyl, heterocyclic group,

30 heterocyclylC₁₋₁₀alkyl, carbocyclyl-(C₁₋₁₀alkylene)_e-R⁴³-(C₁₋₁₀alkylene)_f or

heterocyclyl-(C₁₋₁₀alkylene)_g-R⁴⁴-(C₁₋₁₀alkylene)_h-; wherein R¹⁷ may be optionally substituted

on carbon by one or more substituents selected from R^{47} ; and wherein if said heterocyclyl contains an -NH- group, that nitrogen may be optionally substituted by a group selected from R^{48} ; or R^{17} is a group of formula (BIC):



(BIC)

wherein:

R^{18} is selected from hydrogen or C_{1-4} alkyl;

R^{19} is selected from hydrogen, halo, nitro, cyano, hydroxy, amino, carbamoyl, mercapto, sulphamoyl, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkoxy, C_{1-6} alkanoyl, C_{1-6} alkanoyloxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl)₂amino, C_{1-6} alkanoylamino, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl)₂carbamoyl; C_{1-6} alkylS(O)_a wherein a is 0 to 2, C_{1-6} alkoxycarbonyl, N -(C_{1-6} alkyl)sulphamoyl, N,N -(C_{1-6} alkyl)₂sulphamoyl, carbocyclyl or heterocyclic group; where R^{19} may be independently optionally substituted on carbon by one or more substituents selected from R^{51} ; and wherein if said heterocyclyl contains an -NH- group, that nitrogen may be optionally substituted by a group selected from R^{52} ;

R^{20} is selected from halo, nitro, cyano, hydroxy, amino, carbamoyl, mercapto, sulphamoyl, hydroxyaminocarbonyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy, C_{1-10} alkoxycarbonyl, C_{1-10} alkanoyl, C_{1-10} alkanoyloxy, N -(C_{1-10} alkyl)amino, N,N -(C_{1-10} alkyl)₂amino, N,N,N -(C_{1-10} alkyl)₃ammonio, C_{1-10} alkanoylamino, N -(C_{1-10} alkyl)carbamoyl, N,N -(C_{1-10} alkyl)₂carbamoyl, C_{1-10} alkylS(O)_a wherein a is 0 to 2, N -(C_{1-10} alkyl)sulphamoyl, N,N -(C_{1-10} alkyl)₂sulphamoyl, N -(C_{1-10} alkyl)sulphamoylamino, N,N -(C_{1-10} alkyl)₂sulphamoylamino, C_{1-10} alkoxycarbonylamino, carbocyclyl, carbocyclyl C_{1-10} alkyl, heterocyclic group, heterocyclyl C_{1-10} alkyl, carbocyclyl-(C_{1-10} alkylene)_e- R^{53} -(C_{1-10} alkylene)_f or heterocyclyl-(C_{1-10} alkylene)_g- R^{54} -(C_{1-10} alkylene)_h; wherein R^{20} may be independently optionally substituted on carbon by one or more R^{57} ; and wherein if said heterocyclyl contains an -NH- group, that nitrogen may be optionally substituted by a group selected from R^{58} ;

p is 1-3; wherein the values of R^{15} may be the same or different;

q is 0-1;

r is 0-3; wherein the values of R^{16} may be the same or different;

m is 0-2; wherein the values of R^{12} may be the same or different;

n is 1-2; wherein the values of R^8 may be the same or different;

z is 0-3; wherein the values of R^{19} may be the same or different;

R^{21} is selected from hydrogen or C_{1-6} alkyl;

R^{22} and R^{23} are independently selected from hydrogen, hydroxy, amino, mercapto,

- 5 C_{1-6} alkyl, C_{1-6} alkoxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl)₂amino, C_{1-6} alkylS(O)_a wherein a is 0 to 2;

R^{24} is selected from hydrogen, hydroxy, C_{1-6} alkyl, C_{1-4} alkoxy and C_{1-6} alkanoyloxy;

R^{25} is selected from halo, nitro, cyano, hydroxy, amino, carboxy, carbamoyl,

mercapto, sulphamoyl, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{1-4} alkoxy, C_{1-4} alkanoyl,

- 10 C_{1-4} alkanoyloxy, N -(C_{1-4} alkyl)amino, N,N -(C_{1-4} alkyl)₂amino, C_{1-4} alkanoylamino, N -(C_{1-4} alkyl)carbamoyl, N,N -(C_{1-4} alkyl)₂carbamoyl, C_{1-4} alkylS(O)_a wherein a is 0 to 2, C_{1-4} alkoxycarbonyl, N -(C_{1-4} alkyl)sulphamoyl and N,N -(C_{1-4} alkyl)₂sulphamoyl; wherein R^{25} , may be independently optionally substituted on carbon by one or more R^{67} ;

R^{26} , R^{28} , R^{30} , R^{36} , R^{41} , R^{47} , R^{51} and R^{57} are independently selected from halo, nitro,

- 15 cyano, hydroxy, amino, carbamoyl, mercapto, sulphamoyl, hydroxyaminocarbonyl, C_{1-10} alkyl, C_{2-10} alkenyl, C_{2-10} alkynyl, C_{1-10} alkoxy, C_{1-10} alkanoyl, C_{1-10} alkanoyloxy, C_{1-10} alkoxycarbonyl, N -(C_{1-10} alkyl)amino, N,N -(C_{1-10} alkyl)₂amino, N,N,N -(C_{1-10} alkyl)₃ammonio, C_{1-10} alkanoylamino, N -(C_{1-10} alkyl)carbamoyl, N,N -(C_{1-10} alkyl)₂carbamoyl, C_{1-10} alkylS(O)_a wherein a is 0 to 2, N -(C_{1-10} alkyl)sulphamoyl,
- 20 N,N -(C_{1-10} alkyl)₂sulphamoyl, N -(C_{1-10} alkyl)sulphamoylamino, N,N -(C_{1-10} alkyl)₂sulphamoylamino, C_{1-10} alkoxycarbonylamino, carbocyclyl, carbocyclyl C_{1-10} alkyl, heterocyclic group, heterocyclyl C_{1-10} alkyl, carbocyclyl-(C_{1-10} alkylene)_e- R^{59} -(C_{1-10} alkylene)_f or heterocyclyl-(C_{1-10} alkylene)_g- R^{60} -(C_{1-10} alkylene)_h; wherein R^{26} , R^{28} , R^{30} , R^{36} , R^{41} , R^{47} , R^{51}

- 25 and R^{57} may be independently optionally substituted on carbon by one or more R^{63} ; and wherein if said heterocyclyl contains an -NH- group, that nitrogen may be optionally substituted by a group selected from R^{64} ;

R^{27} , R^{29} , R^{31} , R^{37} , R^{42} , R^{48} , R^{52} , R^{58} and R^{64} are independently selected from

C_{1-6} alkyl, C_{1-6} alkanoyl, C_{1-6} alkylsulphonyl, sulphamoyl, N -(C_{1-6} alkyl)sulphamoyl,

- 30 N,N -(C_{1-6} alkyl)₂sulphamoyl, C_{1-6} alkoxycarbonyl, carbamoyl, N -(C_{1-6} alkyl)carbamoyl, N,N -(C_{1-6} alkyl)₂carbamoyl, benzyl, phenethyl, benzoyl, phenylsulphonyl and phenyl;

$R^{32}, R^{33}, R^{43}, R^{44}, R^{53}, R^{54}, R^{59}$ and R^{60} are independently selected from -O-, -NR⁶⁵-, -S(O)_x-, -NR⁶⁵C(O)NR⁶⁶-, -NR⁶⁵C(S)NR⁶⁶-, -OC(O)N=C-, -NR⁶⁵C(O)- or -C(O)NR⁶⁵-; wherein R^{65} and R^{66} are independently selected from hydrogen or C₁₋₆alkyl, and x is 0-2;

R^{63} and R^{67} are independently selected from halo, hydroxy, cyano, carbamoyl, ureido, amino, nitro, carbamoyl, mercapto, sulphamoyl, trifluoromethyl, trifluoromethoxy, methyl, ethyl, methoxy, ethoxy, vinyl, allyl, ethynyl, methoxycarbonyl, formyl, acetyl, formamido, acetylamino, acetoxymethyl, methylamino, dimethylamino, *N*-methylcarbamoyl, *N,N*-dimethylcarbamoyl, methylthio, methylsulphinyl, mesyl, *N*-methylsulphamoyl and *N,N*-dimethylsulphamoyl; and

10 e, f, g and h are independently selected from 0-2;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

Additional suitable IBAT inhibitors having the above structure for use in combination with compounds of the present invention are selected from any one of:

- (+/-)-trans-1,1-dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-(S)-3-(R)-4-
15 (R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine;
- (+/-)-trans-1,1-dioxo-3-ethyl-3-butyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-(S)-3-(R)-4-
(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,4-benzothiazepine;
- 20 1,1-dioxo-3-ethyl-3-butyl-4-hydroxy-5-phenyl-7-(*N*-{ α -[*N'*-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]-2-fluorobenzyl}carbamoylmethylthio)-2,3,4,5-tetrahydrobenzothiazepine; or
- 1,1-dioxo-3-butyl-3-ethyl-4-hydroxy-5-phenyl-7-(*N*-{1-[*N'*-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]-1-(cyclohexyl)methyl}carbamoylmethylthio)-
25 2,3,4,5-tetrahydrobenzothiazepine.

Compounds of formula (AI) and (BI) or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof may be prepared by processes known in the art.

In a particular aspect of the invention an IBAT inhibitor or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof is an IBAT inhibitor or a
30 pharmaceutically acceptable salt thereof.

Therefore in an additional feature of the invention, there is provided a combination of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a

salt or a prodrug thereof and an IBAT inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

Therefore in an additional feature of the invention, there is provided a method for producing a cholesterol lowering effect in a warm-blooded animal, such as man, in need of
5 such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of an IBAT inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

10 According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and an IBAT inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in association with a pharmaceutically acceptable diluent or carrier.

15 According to a further aspect of the present invention there is provided a kit comprising a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and an IBAT inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

According to a further aspect of the present invention there is provided a kit
20 comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a first unit dosage form;
- b) an IBAT inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof; in a second unit dosage form; and
- 25 c) container means for containing said first and second dosage forms.

According to a further aspect of the present invention there is provided a kit comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, together with a pharmaceutically acceptable diluent or carrier, in a
30 first unit dosage form;
- b) an IBAT inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a second unit dosage form; and
- c) container means for containing said first and second dosage forms.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and an IBAT inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the
5 production of a cholesterol lowering effect in a warm-blooded animal, such as man.

According to a further aspect of the present invention there is provided a combination treatment comprising the administration of an effective amount of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with the
10 simultaneous, sequential or separate administration of an effective amount of an IBAT inhibitor, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier to a warm-blooded animal, such as man in need of such therapeutic treatment.

In another aspect of the invention, the compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, may be administered in
15 association with a PPAR alpha and/or gamma agonist, or pharmaceutically acceptable salts, solvates, solvates of such salts or prodrugs thereof. Suitable PPAR alpha and/or gamma agonists, pharmaceutically acceptable salts, solvates, solvates of such salts or prodrugs thereof are well known in the art. These include the compounds described in WO 01/12187, WO
20 01/12612, WO 99/62870, WO 99/62872, WO 99/62871, WO 98/57941, WO 01/40170, WO03/051821, WO03/051822, WO03/051826, PCT/GB03/02584, PCT/GB03/02591, PCT/GB03/02598, J Med Chem, 1996, 39, 665, Expert Opinion on Therapeutic Patents, 10 (5), 623-634 (in particular the compounds described in the patent applications listed on page 634) and J Med Chem, 2000, 43, 527 which are all incorporated herein by reference.

25 Particularly a PPAR alpha and/or gamma agonist refers to WY-14643, clofibrate, fenofibrate, bezafibrate, GW 9578, troglitazone, pioglitazone, rosiglitazone, eglitazone, proglitazone, NN622/Ragaglitazar, BMS 298585, BRL-49634, KRP-297, JTT-501, SB 213068, GW 1929, GW 7845, GW 0207, L-796449, L-165041 and GW 2433. Particularly a PPAR alpha and/or gamma agonist refers to (S)-2-ethoxy-3-[4-(2-{4-methanesulphonyloxyphenyl}ethoxy)
30 phenyl]propanoic acid and pharmaceutically acceptable salts thereof.

Therefore in an additional feature of the invention, there is provided a combination of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a

salt or a prodrug thereof and a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

Therefore in an additional feature of the invention, there is provided a method for producing a cholesterol lowering effect in a warm-blooded animal, such as man, in need of
5 such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

10 According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in association with a pharmaceutically acceptable diluent or carrier.

15 According to a further aspect of the present invention there is provided a kit comprising a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

According to a further aspect of the present invention there is provided a kit
20 comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a first unit dosage form;
- b) a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof; in a second unit dosage form; and
- 25 c) container means for containing said first and second dosage forms.

According to a further aspect of the present invention there is provided a kit comprising:

- a) a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, together with a pharmaceutically acceptable diluent or carrier, in a
30 first unit dosage form;
- b) a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a second unit dosage form; and
- c) container means for containing said first and second dosage forms.

According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament
5 for use in producing a cholesterol lowering effect in a warm-blooded animal, such as man.

According to a further aspect of the present invention there is provided a combination treatment comprising the administration of an effective amount of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with the
10 simultaneous, sequential or separate administration of an effective amount of a PPAR alpha and/or gamma agonist, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier to a warm-blooded animal, such as man in need of such therapeutic treatment.

In another aspect of the invention, the compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, may be administered in association with a nicotinic acid derivative or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof. As used herein "nicotinic acid derivative" means a compounds comprising a pyridine-3-carboxylate structure or a pyrazine-2-carboxylate structure. Examples of nicotinic acid derivatives include nicotinic acid, niceritrol,
15 nicofuranose, NIASPAN® and acipimox.
20

Therefore, in an additional feature of the invention, there is provided a combination of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof and a nicotinic acid derivative or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

25 Therefore in an additional feature of the invention, there is provided a method for producing a cholesterol lowering effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective
30 amount of a nicotinic acid derivative, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable

salt, solvate, solvate of such a salt or a prodrug thereof, and a nicotinic acid derivative, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in association with a pharmaceutically acceptable diluent or carrier.

According to another feature of the invention there is provided the use of a compound
5 of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a nicotinic acid derivative, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the production of a cholesterol lowering effect in a warm-blooded animal, such as man.

In another aspect of the invention, the compound of formula (I), or a pharmaceutically
10 acceptable salt, solvate, solvate of such a salt or a prodrug thereof, may be administered in association with a bile acid sequestrant or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof. Suitable bile acid sequestrants include cholestyramine, cholestipol and cosevelam hydrochloride.

Therefore, in an additional feature of the invention, there is provided a combination of
15 a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof and a bile acid sequestrant or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

Therefore in an additional feature of the invention, there is provided a method for
producing a cholesterol lowering effect in a warm-blooded animal, such as man, in need of
20 such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of a bile acid sequestrant, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

25 According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a bile acid sequestrant, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in association with a pharmaceutically acceptable diluent or carrier.

30 According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a bile acid sequestrant, or a pharmaceutically acceptable salt, solvate,

solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the production of a cholesterol lowering effect in a warm-blooded animal, such as man.

According to an additional further aspect of the present invention there is provided a combination treatment comprising the administration of an effective amount of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with the simultaneous, sequential or separate administration one or more of the following agents selected from Group X:

- an antihypertensive compound (for example althiazide, benzthiazide, captopril, carvedilol, chlorothiazide sodium, clonidine hydrochloride, cyclothiazide, delapril hydrochloride, dilevalol hydrochloride, doxazosin mesylate, fosinopril sodium, guanfacine hydrochloride, methyldopa, metoprolol succinate, moexipril hydrochloride, monatepil maleate, pelanserine hydrochloride, phenoxybenzamine hydrochloride, prazosin hydrochloride, primidolol, quinapril hydrochloride, quinaprilat, ramipril, terazosin hydrochloride, candesartan, candesartan cilexetil, telmisartan, amlodipine besylate, amlodipine maleate and bevantolol hydrochloride);
- an angiotensin converting enzyme inhibitor (for example alacepril, alatriopril, altiopril calcium, ancovenin, benazepril, benazepril hydrochloride, benazeprilat, benzoylcaptopril, captopril, captopril-cysteine, captopril-glutathione, ceranapril, ceranopril, ceronapril, cilazapril, cilazaprilat, delapril, delapril-diacid, enalapril, enalaprilat, enapril, epicaptopril, foroxymithine, fosfenopril, fosenopril, fosenopril sodium, fosinopril, fosinopril sodium, fosinoprilat, fosinoprilic acid, glycopril, hemorphan-4, idrapril, imidapril, indolapril, indolaprilat, libenzapril, lisinopril, lyciumin A, lyciumin B, mixanapril, moexipril, moexiprilat, moveltipril, muracein A, muracein B, muracein C, pentopril, perindopril, perindoprilat, pivalopril, pivopril, quinapril, quinapril hydrochloride, quinaprilat, ramipril, ramiprilat, spirapril, spirapril hydrochloride, spiraprilat, spiropril, spiropril hydrochloride, temocapril, temocapril hydrochloride, teprotide, trandolapril, trandolaprilat, utibapril, zabicipril, zabiciprilat, zofenopril and zofenoprilat);
- an angiotensin II receptor antagonist (for example candesartan, candesartan cilexetil, losartan, valsartan, irbesartan, tasosartan, telmisartan and eprosartan);
- an adrenergic blocker (for example bretylium tosylate, dihydroergotamine so mesylate, phentolamine mesylate, solypertine tartrate, zolertine hydrochloride,

carvedilol or labetalol hydrochloride); an alpha andrenergic blocker (for example fenspiride hydrochloride, labetalol hydrochloride, proroxan and alfuzosin hydrochloride); a beta andrenergic blocker (for example acebutolol, acebutolol hydrochloride, alprenolol hydrochloride, atenolol, bunolol hydrochloride, carteolol hydrochloride, celiprolol hydrochloride, cetamolol hydrochloride, cicloprolol hydrochloride, dexpropranolol hydrochloride, diacetolol hydrochloride, dilevalol hydrochloride, esmolol hydrochloride, exaprolol hydrochloride, fleistolol sulfate, labetalol hydrochloride, levobetaxolol hydrochloride, levobunolol hydrochloride, metalol hydrochloride, metoprolol, metoprolol tartrate, nadolol, pamatolol sulfate, penbutolol sulfate, practolol, propranolol hydrochloride, sotalol hydrochloride, timolol, timolol maleate, tiprenolol hydrochloride, tolamolol, bisoprolol, bisoprolol fumarate and nebivolol); or a mixed alpha/beta andrenergic blocker;

➤ an andrenergic stimulant (for example combination product of chlorothiazide and methyl dopa, the combination product of methyldopa hydrochlorothiazide and methyl dopa, clonidine hydrochloride, clonidine, the combination product of chlorthalidone and clonidine hydrochloride and guanfacine hydrochloride);

➤ channel blocker, for example a calcium channel blocker (for example clentiazem maleate, amlodipine besylate, isradipine, nimodipine, felodipine, nilvadipine, nifedipine, teludipine hydrochloride, diltiazem hydrochloride, belfosdil, verapamil hydrochloride or fostedil);

➤ a diuretic (for example the combination product of hydrochlorothiazide and spironolactone and the combination product of hydrochlorothiazide and triamterene);

➤ anti-anginal agents (for example amlodipine besylate, amlodipine maleate, betaxolol hydrochloride, bevantolol hydrochloride, butoprozine hydrochloride, carvedilol, cinepazet maleate, metoprolol succinate, molsidomine, monatepil maleate, primidolol, ranolazine hydrochloride, tosifen or verapamil hydrochloride);

➤ vasodilators for example coronary vasodilators (for example fostedil, azaclozine hydrochloride, chromonar hydrochloride, clonitrate, diltiazem hydrochloride, dipyrindamole, droprenilamine, erythrityl tetranitrate, isosorbide dinitrate, isosorbide mononitrate, lidoflazine, mioflazine hydrochloride, mixidine, molsidomine, nicorandil, nifedipine, nisoldipine, nitroglycerine, oxprenolol hydrochloride, pentrinitrol, perhexiline maleate, prenylamine, propatyl nitrate, terodiline hydrochloride, tolamolol and verapamil);

- anti-coagulants (selected from argatroban, bivalirudin, dalteparin sodium, desirudin, dicumarol, Iyapolate sodium, nafamostat mesylate, phenprocoumon, tinzaparin sodium and warfarin sodium);
- antithrombotic agents (for example anagrelide hydrochloride, bivalirudin, cilostazol, dalteparin sodium, danaparoid sodium, dazoxiben hydrochloride, efegatran sulfate, enoxaparin sodium, fluretofen, ifetroban, ifetroban sodium, lamifiban, lotrafiban hydrochloride, napsagatran, orbofiban acetate, roxifiban acetate, sibrafiban, tinzaparin sodium, trifenagrel, abciximab and zolimomab aritox);
- fibrinogen receptor antagonists (for example roxifiban acetate, fradafiban, orbofiban, lotrafiban hydrochloride, tirofiban, xemilofiban, monoclonal antibody 7E3 and sibrafiban)
- platelet inhibitors (for example cilostezol, clopidogrel bisulfate, epoprostenol, epoprostenol sodium, ticlopidine hydrochloride, aspirin, ibuprofen, naproxen, sulindae, indomethacin, mefenamate, droxicam, diclofenac, sulfinpyrazone and piroxicam, dipyridamole);
- platelet aggregation inhibitors (for example acadesine, beraprost, beraprost sodium, ciprostone calcium, itezigrel, lifarizine, lotrafiban hydrochloride, orbofiban acetate, oxagrelate, fradafiban, orbofiban, tirofiban and xemilofiban)
- hemorrhheologic agents (for example pentoxifylline);
- lipoprotein associated coagulation inhibitors;
- Factor VIIa inhibitors;
- Factor Xa inhibitors;
- low molecular weight heparins (for example enoxaparin, nardroparin, dalteparin, certroparin, parnaparin, reviparin and tinzaparin);
- squalene synthase inhibitors;
- squalene epoxidase inhibitors;
- liver X receptor (LXR) agonists for example GW-3965 and those described in WO00224632, WO00103705, WO02090375 and WO00054759 (claim 1 and the named examples of these four application are incorporated herein by reference);
- microsomal triglyceride transfer protein inhibitors for example implitapide and those described in WO03004020, WO03002533, WO02083658 and WO 00242291 (claim 1 and the named examples of these four application are incorporated herein by reference);

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier to a warm-blooded animal, such as man in need of such therapeutic treatment.

Therefore, in an additional feature of the invention, there is provided a combination of
5 a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof and a compound from Group X or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

Therefore in an additional feature of the invention, there is provided a method for producing a cholesterol lowering effect in a warm-blooded animal, such as man, in need of
10 such treatment which comprises administering to said animal an effective amount of a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of a compound from Group X, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

15 According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a compound from Group X, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in association with a pharmaceutically acceptable diluent or carrier.

20 According to another feature of the invention there is provided the use of a compound of the formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and a compound from Group X, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the production of a cholesterol lowering effect in a warm-blooded animal, such as man.

25 In addition to their use in therapeutic medicine, the compounds of formula (I), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, are also useful as pharmacological tools in the development and standardisation of *in vitro* and *in vivo* test systems for the evaluation of the effects of inhibitors of cholesterol absorption in laboratory animals such as cats, dogs, rabbits, monkeys, rats and mice, as part of the search
30 for new therapeutic agents.

Many of the intermediates described herein are novel and are thus provided as a further feature of the invention. For example compounds of formula (VI) show cholesterol

absorption inhibitory activity when tested in the above referenced *in vitro* test assay and are thus claimed as a further feature of the invention.

Thus in a further feature of the invention, there is provided a compound of formula (VI), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, with the proviso that said compound is not 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-2-hydroxyethylsulphanyl]-4-{4-[N-(carboxymethyl)carbamoylmethoxy] phenyl}azetidin-2-one.

Therefore according to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula (VI), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in association with a pharmaceutically-acceptable diluent or carrier, with the proviso that said compound is not 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-2-hydroxyethylsulphanyl]-4-{4-[N-(carboxymethyl)carbamoylmethoxy]phenyl}azetidin-2-one.

According to an additional aspect of the present invention there is provided a compound of the formula (VI), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, for use in a method of prophylactic or therapeutic treatment of a warm-blooded animal, such as man, with the proviso that said compound is not 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-2-hydroxyethylsulphanyl]-4-{4-[N-(carboxymethyl)carbamoylmethoxy]phenyl}azetidin-2-one.

Thus according to this aspect of the invention there is provided a compound of the formula (VI), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, for use as a medicament, with the proviso that said compound is not 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-2-hydroxyethylsulphanyl]-4-{4-[N-(carboxymethyl)carbamoylmethoxy]phenyl}azetidin-2-one.

According to another feature of the invention there is provided the use of a compound of the formula (VI), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the production of a cholesterol absorption inhibitory effect in a warm-blooded animal, such as man, with the proviso that said compound is not 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-2-hydroxyethylsulphanyl]-4-{4-[N-(carboxymethyl)carbamoylmethoxy] phenyl}azetidin-2-one.

According to another feature of the invention there is provided the use of a compound of the formula (VI), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the treatment of hyperlipidaemic conditions in a warm-blooded animal, such as man, with the proviso that said

compound is not 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-2-hydroxyethylsulphanyl]-4-{4-[N-(carboxymethyl)carbamoylmethoxy] phenyl}azetidin-2-one.

According to a further feature of this aspect of the invention there is provided a method for producing a cholesterol absorption inhibitory effect in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (VI), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, with the proviso that said compound is not 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-2-hydroxyethylsulphanyl]-4-{4-[N-(carboxymethyl)carbamoylmethoxy] phenyl}azetidin-2-one.

According to a further feature of this aspect of the invention there is provided a method of treating hyperlipidemic conditions in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula (VI), or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, with the proviso that said compound is not 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-2-hydroxyethylsulphanyl]-4-{4-[N-(carboxymethyl)carbamoylmethoxy]phenyl}azetidin-2-one.

In the above other pharmaceutical composition, process, method, use and medicament manufacture features, the alternative and preferred embodiments of the compounds of the invention described herein also apply.

20 Examples

The invention will now be illustrated in the following non limiting Examples, in which standard techniques known to the skilled chemist and techniques analogous to those described in these Examples may be used where appropriate, and in which, unless otherwise stated:

- (i) evaporations were carried out by rotary evaporation *in vacuo* and work up procedures were carried out after removal of residual solids such as drying agents by filtration;
- (ii) all reactions were carried out under an inert atmosphere at ambient temperature, typically in the range 18-25°C, with solvents of HPLC grade under anhydrous conditions, unless otherwise stated;
- (iii) column chromatography (by the flash procedure) was performed on Silica gel 40-63 µm (Merck);
- (iv) yields are given for illustration only and are not necessarily the maximum attainable;
- (v) the structures of the end products of the formula (I) were generally confirmed by nuclear (generally proton) magnetic resonance (NMR) and mass spectral techniques; magnetic

resonance chemical shift values were measured in deuterated CDCl_3 (unless otherwise stated) on the delta scale (ppm downfield from tetramethylsilane); proton data is quoted unless otherwise stated; spectra were recorded on a Varian Mercury-300 MHz, Varian Unity plus-400 MHz, Varian Unity plus-600 MHz or on Varian Inova-500 MHz spectrometer unless otherwise stated data was recorded at 400MHz; and peak multiplicities are shown as follows: s, singlet; d, doublet; dd, double doublet; t, triplet; tt, triple triplet; q, quartet; tq, triple quartet; m, multiplet; br, broad; ABq, AB quartet; ABd, AB doublet, ABdd, AB doublet of doublets; dABq, doublet of AB quartets; LCMS were recorded on a Waters ZMD, LC column xTerra MS C_8 (Waters), detection with a HP 1100 MS-detector diode array equipped; mass spectra (MS) (loop) were recorded on VG Platform II (Fisons Instruments) with a HP-1100 MS-detector diode array equipped; unless otherwise stated the mass ion quoted is (MH^+); unless further details are specified in the text, analytical high performance liquid chromatography (HPLC) was performed on Prep LC 2000 (Waters), Cromasil C_8 , 7 μm , (Akzo Nobel); MeCN and de-ionised water 10 mM ammonium acetate as mobile phases, with suitable composition;

(vii) intermediates were not generally fully characterised and purity was assessed by thin layer chromatography (TLC), HPLC, infra-red (IR), MS or NMR analysis;

(viii) where solutions were dried sodium sulphate was the drying agent; and

(ix) the following abbreviations may be used hereinbefore or hereinafter:-

20	DCM	dichloromethane;
	DMF	<i>N,N</i> -dimethylformamide;
	TBTU	<i>o</i> -Benzotriazol-1-yl- <i>N,N,N',N'</i> -tetramethyluronium tetrafluoroborate;
	EtOAc	ethyl acetate;
	MeCN	acetonitrile;
25	TFA	trifluoroacetic acid;
	DMAP	4-(dimethylamino)pyridine;
	BSA	<i>N,O</i> -Bis(trimethylsilyl)acetamide; and
	TBAF	tetrabutylammonium fluoride;
	NMM	<i>N</i> -methyl morpholine;
30	TEA	triethylamine;
	DBN	1,5-diazabicyclo-[4,3,0]-non-5-ene.

5 Example 1

(2R)-[[(N-[[4-((2R,3R)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl]glycyl)amino](phenyl)acetic acid

- 10 *N*-[[4-((2R,3R)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl]glycine (44 mg, 0.081 mmol), *N*-methylmorpholine (18 ul, 0.16 mmol), *tert*-butyl (2R)-amino(phenyl)acetate (23 mg, 0.11 mmol) and TBTU (36 mg, 0.11 mmol) were added to methylene chloride (3 ml) and the reaction mixture was stirred at ambient
- 15 temperature for 1.5 h. The mixture was purified by column chromatography on silica gel using methylene chloride/ethylacetate (1/1) as eluent. The resulting intermediate ester was dissolved in formic acid (2 ml) and the mixture was stirred overnight at 45 °C. The solvent was evaporated under reduced pressure and was co-evaporated with toluene. The residue was dissolved in methanol (2 ml) and NaBH₄ 10 mg, 0.26 mmol) was added. The reaction mixture
- 20 was stirred at ambient temperature for 15 min. A small amount of ammoniumacetate buffer was added and the methanol was evaporated off. The residue was purified by preparativ HPLC using acetonitril/ammonium acetat buffer (45:55) as eluent. The collected fractions were lyophilized to obtain 39 mg (71%, 3 steps) of the title compound. (¹H-NMR, 400 MHz, CD₃OD): 1.8-2.0 (m, 2H), 3.75 (d, 2H), 4.3 (d, 1H), 4.55 (s, 2H), 4.65-4.80 (m, 2H), 5.05 (d,
- 25 1H), 7.0-7.4 (m, 17 H), 7.95 (dd, 1H) 8.25 (t, 1H)

Example 2

- 30 *N*-[[4-((2R,3R)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl]glycyl-*N*⁶-acetyl-D-lysine

- N*-[[4-((2R,3R)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl]glycine (20 mg, 0.037 mmol), *N*-methylmorpholine (18 ul, 0.16 mmol) and TBTU (13 mg, 0.041 mol) were added to CH₂Cl₂ (2 ml) and the reaction mixture was stirred at 30 °C for 1.5 h. To the mixture were *N*⁶-acetyl-D-lysine (14 mg, 0.074 mmol) and DMF (5 drops) added and the mixture was stirred at ambient temperature. overnight. The solvent was evaporated under reduced pressure and methanol (2 ml) and NaBH₄ (30 mg, 0.79 mmol) were added to the residue. The reaction mixture was stirred at ambient temperature for
- 40 1 h.. A small amount of ammoniumacetate buffer was added and the methanol was evaporated off. The residue was purified by preparativ HPLC using acetonitril/ammonium acetat buffer (40:60) as eluent. The collected fractions were lyophilized, solved in *tert*-butanol and lyophilized again to obtain 4 mg (15 % 2 steps) of the title compound. (¹H-NMR, 400 MHz, DMSO-*d*₆): 1.2-1.8 (m, 11H), 2.85-2.95 (m, 2H), 3.7-3.8 (m, 2H), 4.0-4.1 (bs, 1H), 4.25-4.3
- 45 (m, 1H), 4.5 (s, 2H), 4.7-4.8 (m, 1H), 5.05 (dd, 1H), 6.95-7.4 (m, 12H), 7.7-7.8 (m, 1H), 7.95-8.05 (m, 1H), 8.25-8.3 (m, 1H)

Example 3

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}-*D*-alanine-*D*-phenylalanine

5

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}-*D*-alanine (12 mg, 0.022 mmol), *tert*-butyl *D*-phenylalaninate hydrochloride (7 mg, 0.028 mmol) and 4-methylmorpholine (7mg, 0.065 mmol) were dissolved in 0.5 ml DCM. The mixture was stirred at room temperature for 5 minutes. TBTU (9 mg, 0.028mmol) was added and the reaction mixture was stirred overnight and then purified on silica (2g), first with DCM and then with DCM:MeOH, 10:1 as eluent. The fractions containing the product were concentrated and formic acid (1 ml) was added and the solution was stirred overnight. The formic acid was evaporated whereafter residual formic acid was removed by addition of toluene and evaporation. The residue was stirred in methanol: triethylamine, 40:1, (2 ml) for 2.5 days (hydrolysis of the formed formic acid ester), the mixture was concentrated and the residue was purified by preparative HPLC using a gradient from 20 % to 100% CH₃CN in 0.1% ammonium acetate buffer as eluent. Freeze-drying gave 7.3 mg (48 %) of the title product. *M/z*: 702.3 (*M* - 1), NMR (DMSO, 400 MHz): 1.18 (d,3H), 2.85-2.96 (m,3H), 3.00-3.07 (m,1H), 4.05-4.12 (m,1H), 4.19-4.32 (m,2H), 4.48 (dd,2H), 4.68-4.76 (m,1H), 5.0-5.07 (m,1H), 6.89-6.95 (m,2H), 7.05-7.17 (m,10H), 7.19-7.25 (m,2H), 7.30-7.36 (m,4H), 7.65-7.75 (m,1H), 8.12-8.18 (m,1H).

25

Example 4

*N*²-[[(2*R*)-2-([4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl)amino]-2-phenylacetyl]-*D*-lysine

30

(2*R*)-([4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl)amino)(phenyl)acetic acid (10 mg, 0.016mmol), *tert*-butyl *N*⁶-(*tert*-butoxycarbonyl)-*D*-lysinate hydrochloride (7mg, 0.021mmol) and *N*-methylmorpholine (5mg, 0.048mmol) were dissolved in DCM (0.5ml). The mixture was stirred and after 5 min TBTU (7mg, 0.021) was added and the reaction mixture was stirred for two hours. The product was purified on a silica plug with DCM:MeOH, 100:5 as the eluent. Formic acid (0.5ml) was added to the purified product and the mixture was stirred two hours at 40 °C and then at ambient temperature overnight. The formic acid was removed under reduced pressure and toluene was added whereafter the solvent was removed. The residue was dissolved in MeOH and a small amount of triethylamine and stirred overnight at ambient temperature. The product was purified by preparative HPLC (CH₃CN/ 0.1% ammonium acetate 20:80-100:0). The fractions containing the product were lyophilized and 4.0mg (33%) of the title compound was obtained. *M/z*: 745.4 (*M* - 1). NMR (DMSO, 400 MHz): 1.19-2.30 (m,2H), 1.30-1.40 (m,2H), 1.40-1.52 (m,2H), 1.52-1.68 (m,2H), 2.63-2.75 (m,2H), 2.83-2.95 (m,2H), 3.7-3.8 (m,1H), 4.28 (dd,1H), 4.60 (dd, 2H), 4.68-4.76 (m,1H), 5.04 (dd,1H), 5.48 (d,1H), 6.95 (d, 2H), 7.05-7.17 (m,4H), 7.19-7.28 (m,5H), 7.30-7.40 (m,6H), 7.88 (d,1H), 8.55 (d,1H).

50

Example 5

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}-*L*-tryptophyl-*D*-phenylalanine

5

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}-*L*-tryptophan (0.015 g, 0.022 mmol) was dissolved in CH₂Cl₂ (2 ml). *H*-*D*-Phe-OTBU hydrochloride (0.007 g, 0.027 mmol) and *N*-Methylmorpholine (0.007 g, 0.067 mmol) were added. After 10 minutes TBTU (0.009 g, 0.029 mmol) was added. The reaction mixture was stirred at room temperature for 2h. The mixture was then passed through a short pad of silica gel and eluted with EtOAc/CH₂Cl₂ 25/75. The pure fractions were collected and concentrated. The resulting tert-butyl ester was hydrolysed by the addition of formic acid (0.5 ml) followed by stirring at room temperature overnight. The resulting product was concentrated and dissolved in 2 ml of MeOH. Et₃N (0.1ml) was added in order to hydrolyze the formylated product. The mixture was allowed to stir overnight. The solvent was evaporated and the residue was purified by preparative HPLC using a gradient of 20-60% CH₃CN in 0.1 M NH₄OAc buffer as mobile phase. Freeze-drying afforded the title product (0.004 g, 22%) as a colourless solid. *M/z*: 817.3 (*M* - 1). ¹H NMR (CD₃CN, 400 MHz) δ 2.82-3.20 (m, 6H), 4.07-4.15 (m, 1H), 4.29-4.39 (m, 2H), 4.54-4.89 (m, 4H), 6.74-7.53 (m, 22H), 9.14-9.20 (m, 1H).

Example 6

25

N-((2*R*)-2-[[[4-((2*R*,3*R*)-3-[(2-hydroxy-2-phenylethyl)thio]-4-oxo-1-phenylazetidin-2-yl)phenoxy]acetyl]amino}-2-phenylacetyl)glycine

30

[4-((2*R*,3*R*)-3-[[[5,5-Dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl]thio]-4-oxo-1-phenylazetidin-2-yl)phenoxy]acetic acid (0.050 g, 0.094 mmol) was dissolved in CH₂Cl₂ (5 ml). *tert*-Butyl *N*-[(2*R*)-2-amino-2-phenylacetyl]glycinate (0.030 g, 0.112 mmol) and *N*-methylmorpholine (0.028 g, 0.281 mmol) were added. After 5 minutes TBTU (0.039 g, 0.122 mmol) was added. After 1h, HPLC showed a full conversion to the corresponding tert-butylester. The reaction mixture was purified by chromatography on silica gel and eluted with EtOAc/CH₂Cl₂ (25/75). Pure fractions were collected and concentrated. The residue was dissolved in 3 ml CH₂Cl₂ and 1 ml trifluoroacetic acid was added. The reaction mixture was allowed to stir at room temperature for 2h after which the resulting acid was concentrated in vacuo. The remaining TFA was removed through co-evaporation with toluene (2 ml). The acid was dissolved in 3 ml of MeOH and sodiumborohydride (10 mg, 0.264 mmol) was added. After 5 minutes, HPLC showed a full conversion to the corresponding alcohol. The reaction was quenched by the addition of 1 ml of 0.1M NH₄OAc buffer followed by concentration of the mixture. Purification by preparative HPLC of the residue using a gradient of 20-60% CH₃CN in 0.1 M NH₄OAc buffer as mobile phase and lyophilisation afforded the title product (0.032 g, 54%) as a colourless solid. *M/z*: 638.5 (*M* - 1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 2.90-2.94 (m, 2H), 3.45-3.65 (m, 2H), 4.25-4.27 (m, 1H), 4.59 (d, 1H), 4.64 (d, 1H), 4.68-4.75 (m, 1H), 5.02-5.05 (m, 1H), 5.57 (d, 1H), 6.94-7.41 (m, 19H), 8.28 (s, br, 1H), 8.55 (d, 1H).

50

Example 7

N-[(4-{(2*R*,3*R*)-3-[(2-hydroxy-2-phenylethyl)thio]-4-oxo-1-phenylazetidin-2-yl}phenoxy)acetyl]glycyl-D-phenylalanine

N-[(4-{(2*R*,3*R*)-4-Oxo-3-[(2-oxo-2-phenylethyl)thio]-1-phenylazetidin-2-yl}phenoxy)acetyl]glycine was dissolved in CH₂Cl₂ (3 ml). (*R*)-Phenylalanine tert-butyl ester hydrochloride (0.012 g, 0.045 mmol) and *N*-methylmorpholine (0.011 g, 0.113 mmol) were added. After 5 minutes TBTU (0.016 g, 0.049 mmol) was added. After 1 h, full conversion to the corresponding tert-butyl ester was obtained. The reaction mixture purified by chromatography on silica gel and eluted with EtOAc/ CH₂Cl₂ 25/75. Pure fractions were collected and concentrated. The residue was dissolved in 3 ml CH₂Cl₂ and 0.5 ml trifluoroacetic acid and the reaction was allowed to stir at room temperature overnight. The resulting acid was concentrated and dissolved in 3 ml of MeOH. Sodiumborohydride (0.010 g, 0.264 mmol) was added. Full conversion to the corresponding alcohol was obtained after 5 minutes. The reaction was quenched by the addition of 1 ml of 0.1M NH₄OAc buffer. Concentration followed by preparative HPLC using a gradient of 20-60% CH₃CN in 0.1M NH₄OAc buffer as mobile phase afforded the title product (0.021 g, 84%) as a colourless solid after freeze-drying. *M/z*: 652.3 (*M* - 1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 2.85-3.05 (m, 4H), 3.62-3.80 (m, 2H), 4.18-4.35 (m, 2H), 4.48 (d, 1H), 4.51 (d, 1H), 4.68-4.78 (m, 1H), 4.99-5.03 (m, 1H), 6.93-7.35 (m, 19H), 7.65-7.80 (m, 1H), 8.23-8.29 (m, 1H).

Example 8

N-[(2*R*)-2-([4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy)acetyl]amino)-2-phenylacetyl]-L-alanine

(2*R*)-([4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl]amino)(phenyl)acetic acid (0.040 g, 0.065 mmol) was dissolved in CH₂Cl₂ (5 ml). *tert*-Butyl L-alaninate hydrochloride (0.014 g, 0.078 mmol) and *N*-methyl-morpholine (0.020 g, 0.195 mmol) were added. After 5 minutes TBTU (0.027 g, 0.084 mmol) was added. The reaction was allowed to stir for 1 h after which the resulting tert-butyl ester was purified by chromatography on silica gel and eluted with EtOAc/ CH₂Cl₂ 25/75. Pure fractions were concentrated and dissolved in CH₂Cl₂ (4 ml) and trifluoroacetic acid (0.5 ml). After 1.5 h at room temperature, full conversion to the corresponding acid was obtained. The reaction mixture was concentrated and the residue of TFA was removed by co-evaporation with toluene (3 ml). The crude acid was dissolved in MeOH (3 ml) and NaBH₄ (0.010 g, 0.260 mmol) was added. Full conversion to the corresponding alcohol was obtained after 5 minutes. The reaction was quenched by the addition of 0.1M NH₄OAc buffer (1 ml). The mixture was concentrated and the residue was purified by preparative HPLC using a gradient of 20-50% CH₃CN in 0.1M NH₄OAc buffer as mobile phase. Freeze-drying afforded the title compound (0.020 g, 45%) as a colourless solid. *M/z*: 688.4 (*M* - 1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 2.30-2.35 (m, 3H), 2.86-2.94 (m, 2H), 3.15-3.31 (m, 1H), 4.26-4.28

(m, 1H), 4.59 (d, 1H), 4.64 (d, 1H), 4.70-4.76 (m, 1H), 5.04-5.07 (m, 1H), 5.46 (d, 1H), 6.94-7.36 (m, 17H), 8.43-8.55 (m, 2H).

5 Example 9

***N*-[*(2R)*-2-([4-((*2R,3R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl]amino)-2-phenylacetyl]-*D*-alanine**

10

(*2R*)-([4-((*2R,3R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl]amino)(phenyl)acetic acid (0.040 g, 0.065 mmol) was dissolved in CH₂Cl₂ (5 ml). *tert*-Butyl *D*-alaninate hydrochloride (0.014 g, 0.078 mmol) and 15 *N*-methyl-morpholine (0.020 g, 0.195 mmol) were added. After 5 minutes TBTU (0.027 g, 0.084 mmol) was added. The reaction was allowed to stir overnight and the resulting *tert*-butyl ester was purified on silica gel and eluted with EtOAc/ CH₂Cl₂ (25/75). The pure fractions were concentrated. The residue was dissolved in CH₂Cl₂ (4 ml) and trifluoroacetic acid (0.5 ml). After 1.5h at room temperature, full conversion to the corresponding acid was 20 obtained. The reaction mixture was concentrated and the remaining TFA was removed through co-evaporation with toluene (3 ml). The crude acid was dissolved in MeOH (3 ml) and NaBH₄ (0.010 g, 0.260 mmol) was added. Full conversion to the corresponding alcohol was obtained after 5 minutes. The reaction was quenched by the addition of 0.1M NH₄OAc buffer (1 ml). The mixture was concentrated and the residue was purified by preparative 25 HPLC using a gradient of 20-50% CH₃CN in 0.1M NH₄OAc buffer as mobile phase. Freeze-drying gave the title compound (0.024 g, 54%) as a colourless solid. *M/z*: 688.6 (*M*-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 1.10-1.24 (m, 3H), 2.89-2.94 (m, 2H), 3.99-4.08 (m, 1H), 4.26-4.30 (m, 1H), 4.60 (d, 1H), 4.64 (d, 1H), 4.70-4.78 (m, 1H), 5.02-5.08 (m, 1H), 5.53-5.56 (d, 1H), 6.94-7.40 (m, 17H), 8.39-8.59 (m, 2H).

30

Example 10

***(2R)*-{[*(2R)*-2-([4-((*2R,3R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl]amino)-2-phenylacetyl]amino}(phenyl)acetic acid**

35

(*2R*)-([4-((*2R,3R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl]amino)(phenyl)acetic acid (0.040 g, 0.065 mmol) was dissolved in CH₂Cl₂ (5 ml) and *tert*-butyl (*2R*)-amino(phenyl)acetate (0.016 g, 0.078 mmol) and *N*-methyl-morpholine (0.020 g, 0.195 mmol) were added. After 5 minutes TBTU (0.027 g, 0.084 mmol) was added. The reaction was allowed to stir at room temperature for 3 h after 45 which the resulting *tert*-butyl ester was purified on silica gel and eluted with EtOAc/ CH₂Cl₂ 25/75. Pure fractions were concentrated. The residue was dissolved in CH₂Cl₂ (3 ml) and TFA (0.5 ml). The reaction mixture was allowed to stir overnight at room temperature. The resulting acid was concentrated and the remaining trace of TFA was removed by co-evaporation with toluene (3 ml). The acid was dissolved in MeOH (3 ml) and NaBH₄ (0.010 g, 0.260 mmol) was added. After 5 minutes, full conversion to the corresponding alcohol was 50

obtained. The reaction was quenched by the addition of 1 ml of 0.1M NH₄OAc buffer. Concentration followed by purification on preparative HPLC using a gradient of 20-50% CH₃CN in 0.1M NH₄OAc buffer as mobile phase and freeze-drying of the pure fractions afforded the title compound (0.034 g, 70%) as a colourless solid. M/z: 750.4 (M-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 2.88-2.94 (m, 2H), 4.23-4.29 (m, 1H), 4.56-4.65 (m, 2H), 4.70-4.78 (m, 1H), 4.91-5.06 (m, 2H), 5.65-5.75 (m, 1H), 6.93-7.42 (m, 22H), 8.54-8.69 (m, 2H).

Example 11

(2S)-{[(2R)-2-({[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}amino)-2-phenylacetyl]amino}(phenyl)acetic acid

(2R)-({[4-((2R,3R)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}amino)(phenyl)acetic acid (0.040 g, 0.065 mmol) was dissolved in CH₂Cl₂ (5 ml). *tert*-Butyl (2S)-amino(phenyl)acetate (0.016 g, 0.078 mmol) and N-methyl-morpholine (0.020 g, 0.195 mmol) were added. After 5 minutes, TBTU (0.027 g, 0.084 mmol) was added. The reaction mixture was stirred at room temperature for 3 h. The resulting *tert*-butyl ester was purified on silica gel and eluted with EtOAc/ CH₂Cl₂ (25/75). The pure fractions were concentrated. The residue was dissolved in CH₂Cl₂ (3 ml) and TFA (0.5 ml). The reaction mixture was stirred overnight at room temperature. The resulting acid was concentrated and the remaining trace of TFA was removed by co-evaporation with toluene (3 ml). The acid was dissolved in MeOH (3 ml) and NaBH₄ (0.010 g, 0.260 mmol) was added. After 5 minutes, full conversion to the corresponding alcohol was obtained. The reaction was quenched by the addition of 1 ml of 0.1M NH₄OAc buffer. Concentration followed by purification on preparative HPLC using a gradient of 20-50% CH₃CN in 0.1M NH₄OAc buffer as mobile phase and freeze-drying of the pure fractions afforded the title compound (0.037 g, 76%) as a colourless solid. M/z: 750.6 (M-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 2.87-2.94 (m, 2H), 4.25-4.28 (m, 1H), 4.58-4.78 (m, 3H), 5.01-5.07 (m, 2H), 5.65-5.74 (m, 1H), 6.94-7.39 (m, 22H), 8.53-8.72 (m, 2H).

Example 12

N-[(2R)-2-({[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}amino)-2-phenylacetyl]-D-serine

(2R)-({[4-((2R,3R)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}amino)(phenyl)acetic acid (0.040 g, 0.065 mmol), *tert*-butyl O-(*tert*-butyl)-D-serinate hydrochloride (0.020 g, 0.078 mmol) and N-methyl-morpholine (0.020 g, 0.195 mmol) were dissolved in CH₂Cl₂ (5 ml) at room temperature. After 5 minutes, TBTU (0.027 g, 0.084 mmol) was added. After 1h, full conversion to the corresponding *tert*-butyl ester was obtained. The reaction mixture was purified on silica gel and eluted with EtOAc/ CH₂Cl₂ (25/75). Pure fractions were collected and concentrated. The residue was dissolved in 3 ml CH₂Cl₂ and 1 ml TFA. The reaction mixture was stirred for 1.5h and concentrated. The remaining trace of TFA was azeotropically removed by co-evaporation with toluene (3 ml). The crude acid was dissolved in MeOH (3 ml) and NaBH₄ (0.010 g,

0.260 mmol) was added. Full conversion to the corresponding alcohol was obtained after 5 minutes. The reaction was quenched by the addition of 0.1M NH₄OAc buffer (2 ml).

- 5 Purification on preparative HPLC using a gradient of 20-50% CH₃CN in 0.1M NH₄OAc buffer as mobile phase followed by freeze-drying of the pure fractions afforded the title compound (0.032 g, 71%) as a colourless solid. M/z: 704.7 (M-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 2.88-2.94 (m, 2H), 3.42-3.46 (m, 1H), 3.58-3.61 (m, 1H), 3.85-3.90 (m, 1H), 4.26-4.29 (m, 1H), 4.59 (d, 1H), 4.64 (d, 1H), 4.70-4.77 (m, 1H), 5.04-5.07 (m, 1H), 5.57-5.65 (m, 1H), 6.95-7.41 (m, 17H), 8.08-8.20 (m, 1H), 8.53-8.56 (m, 1H).

10

Example 13

N-[*(2R)*-2-([4-((*2R,3R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl]amino)-2-phenylacetyl]-*L*-threonine

15

- (*2R*)-([4-((*2R,3R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl]amino)(phenyl)acetic acid (0.030 g, 0.049 mmol), *tert*-butyl *O*-(*tert*-butyl)-*L*-threoninate (0.014 g, 0.058 mmol) and *N*-methyl-morpholine (0.015 g, 0.146 mmol) were dissolved in CH₂Cl₂ (4 ml). After 5 minutes, TBTU (0.020 g, 0.063 mmol) was added. Full conversion to the corresponding *tert*-butyl ester was obtained after 1 h. The reaction mixture was purified on silica gel and eluted with EtOAc/ CH₂Cl₂ (25/75). Pure fractions were collected and concentrated. The residue was dissolved in 3 ml CH₂Cl₂ and 1 ml
- 25 TFA and the mixture was stirred for 1.5 h. The resulting acid was concentrated. The remaining trace of TFA was azeotropically removed by co-evaporation with toluene (3 ml). The crude acid was dissolved in MeOH (3 ml) and NaBH₄ (0.007 g, 0.195 mmol) was added. Full conversion to the corresponding alcohol was obtained after 5 minutes. The reaction was quenched by the addition of 0.1M NH₄OAc buffer (2 ml). Purification by preparative HPLC
- 30 using a gradient of 20-50% CH₃CN in 0.1M NH₄OAc buffer as eluent followed by freeze-drying of the pure fractions afforded the title compound (0.025 g, 71%) as a colourless solid. M/z: 721.1. ¹H NMR [(CD₃)₂SO], 400 MHz] δ 0.69 (d, 3H), 2.88-2.93 (m, 2H), 3.79-3.85 (m, 1H), 3.96-4.01 (m, 1H), 4.27-4.30 (m, 1H), 4.60 (d, 1H), 4.64 (d, 1H), 4.71-4.79 (m, 1H), 5.02-5.08 (m, 1H), 5.65-5.69 (m, 1H), 6.95-7.42 (m, 17H), 8.01-8.09 (m, 1H), 8.51-8.59 (m,
- 35 1H).

Example 14

- 40 *N*²-[*(2R)*-2-([4-((*2R,3R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl]amino)-2-phenylacetyl]-*L*-asparagine

- 45 (*2R*)-([4-((*2R,3R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl]amino)(phenyl)acetic acid (0.030 g, 0.049 mmol), *tert*-butyl *L*-asparaginate hydrochloride (0.013 g, 0.058 mmol) and *N*-methyl-morpholine (0.015 g, 0.146 mmol) were dissolved in CH₂Cl₂ (4 ml). TBTU (0.020 g, 0.063 mmol) was added after 5 minutes. Full conversion to the corresponding *tert*-butyl ester was obtained after 1 h.
- 50 The reaction mixture was purified on silica gel and eluted with EtOAc. The pure fractions

were collected and concentrated. The residue was dissolved in 3 ml CH_2Cl_2 and 1 ml TFA and the mixture was stirred for 0.5h. The resulting acid was concentrated. The remaining trace of TFA was azeotropically removed by co-evaporation with toluene (3 ml). The crude acid was dissolved MeOH (3 ml) and NaBH_4 (0.007 g, 0.195 mmol) was added. Full conversion to the corresponding alcohol was obtained after 5 minutes. The reaction was quenched by the addition of 0.1M NH_4OAc buffer (2 ml). Purification by preparative HPLC using a gradient of 20-40% CH_3CN in 0.1M NH_4OAc buffer as eluent followed by freeze-drying of the pure fractions afforded the title compound (0.024 g, 67%) as a colourless solid. M/z : 731.6 (M-1). ^1H NMR [$(\text{CD}_3)_2\text{SO}$], 400 MHz] δ 2.16-2.23 (m, 1H), 2.32-2.42 (m, 1H), 2.87-2.95 (m, 2H), 4.20-4.30 (m, 2H), 4.59-4.78 (m, 3H), 5.03-5.07 (m, 1H), 5.54-5.59 (m, 1H), 6.68-6.73 (m, 1H), 6.94-7.38 (m, 17H), 7.78-7.83 (m, 1H), 8.22-8.37 (m, 1H), 8.50-8.58 (m, 1H).

Example 15

15 *N*-[(2*R*)-2-({[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}amino)-2-phenylacetyl]-L-aspartic acid

20 (2*R*)-({[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}amino)(phenyl)acetic acid (0.020 g, 0.032 mmol), di-*tert*-butyl L-aspartate hydrochloride (0.011 g, 0.039 mmol) and N-methyl-morpholine (0.010 g, 0.097 mmol) were dissolved in CH_2Cl_2 (3 ml). After 5 minutes, TBTU (0.014 g, 0.042 mmol) was added. Full conversion to the corresponding tert-butyl ester was obtained after 1 h. The reaction mixture was purified on silica gel and eluted with EtOAc/ CH_2Cl_2 (25/75). The pure fractions were collected and concentrated. The residue was dissolved in 3 ml of CH_2Cl_2 and 1 ml of TFA. The reaction mixture was stirred for 2h and the resulting acid was concentrated. The remaining trace of TFA was azeotropically removed by co-evaporation with toluene (3 ml). The crude acid was dissolved in MeOH (3 ml) and NaBH_4 (0.005 g, 0.130 mmol) was added. Full conversion to the corresponding alcohol was obtained after 5 minutes. The reaction was quenched by the addition of 0.1M NH_4OAc buffer (2 ml). Purification by preparative HPLC using a gradient of 10-40% CH_3CN in 0.1M NH_4OAc buffer as eluent followed by freeze-drying of the pure fractions afforded the title compound (0.018 g, 76%) as a colourless solid. M/z : 715.7 (M-18). ^1H NMR [$(\text{CD}_3)_2\text{SO}$], 400 MHz] δ 2.00-2.08 (m, 1H), 2.31-2.50 (m, 1H), 2.88-2.94 (m, 2H), 4.18-4.22 (m, 1H), 4.27-4.30 (m, 1H), 4.60-4.78 (m, 3H), 5.03-5.08 (m, 1H), 5.55-5.65 (m, 1H), 6.94-7.40 (m, 17H), 8.22-8.39 (m, 1H), 8.50-8.59 (m, 1H).

40 Example 16

N-[(2*R*)-2-({[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}amino)-2-phenylacetyl]-D-valine

45

(2*R*)-({[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}amino)(phenyl)acetic acid (0.010 g, 0.016 mmol), *tert*-butyl D-valinate hydrochloride (0.004 g, 0.020 mmol) and N-methyl-morpholine (0.005 g, 0.049 mmol) were dissolved in CH_2Cl_2 (3 ml). After 5 minutes, TBTU (0.007 g, 0.021 mmol)

was added. Full conversion to the corresponding tert-butyl ester was obtained after 1h. The reaction mixture was purified on silica gel and eluted with EtOAc/ CH₂Cl₂ (25/75). The pure fractions were collected and concentrated. The residue was dissolved in 3 ml CH₂Cl₂ and 0.5 ml TFA. The reaction mixture was stirred for 1.5 h and concentrated. The remaining trace of TFA was azeotropically removed by co-evaporation with toluene (3 ml). The crude acid was dissolved in MeOH (2 ml) and NaBH₄ (0.002 g, 0.065 mmol) was added. Full conversion to the corresponding alcohol was obtained after 5 minutes. The reaction was quenched by the addition of 0.1M NH₄OAc buffer (2 ml). Purification by preparative HPLC using a gradient of 20-50% CH₃CN in 0.1M NH₄OAc buffer as mobile phase followed by freeze-drying of the pure fractions afforded the title compound (0.004 g, 34%) as a colourless solid. M/z: 716.6 (M-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 0.81-0.85 (m, 6H), 1.98-2.09 (m, 1H), 2.88-2.93 (m, 2H), 3.92-3.98 (m, 1H), 4.22-4.30 (m, 1H), 4.60 (d, 1H), 4.64 (d, 1H), 4.70-4.78 (m, 1H), 5.02-5.07 (m, 1H), 5.62 (d, 1H), 6.94-7.40 (m, 17H), 8.18-8.22 (m, 1H), 8.52 (d, 1H).

Example 17

N-[(2*R*)-2-({[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}amino)-2-phenylacetyl]-L-valine

(2*R*)-({[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}amino)(phenyl)acetic acid (0.020 g, 0.032 mmol), *tert*-butyl L-valinate hydrochloride (0.008 g, 0.039 mmol) and N-methyl-morpholine (0.010 g, 0.097 mmol) were dissolved in CH₂Cl₂ (4 ml). After 5 minutes, TBTU (0.014 g, 0.042 mmol) was added. Full conversion to the corresponding tert-butyl ester was obtained after 1 h. The reaction mixture was purified on silica gel and eluted with EtOAc/ CH₂Cl₂ (25/75). The pure fractions were collected and concentrated. The residue was dissolved in 3 ml CH₂Cl₂ and 1 ml TFA. The reaction mixture was stirred for 1.5h and concentrated. The remaining trace of TFA was azeotropically removed by co-evaporation with toluene (3 ml). The crude acid was dissolved in MeOH (3 ml) and NaBH₄ (0.005 g, 0.130 mmol) was added. Full conversion to the corresponding alcohol was obtained after 5 minutes. The reaction was quenched by the addition of 0.1M NH₄OAc buffer (2 ml). Purification by preparative HPLC using a gradient of 20-50% CH₃CN in 0.1M NH₄OAc buffer as eluent followed by freeze-drying of the pure fractions afforded the title compound (0.011 g, 47%) as a colourless solid. M/z: 716.5 (M-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 0.60 (d, 3H), 0.68 (d, 3H), 1.95-2.05 (m, 1H), 2.88-2.94 (m, 2H), 4.00-4.08 (m, 1H), 4.27-4.30 (m, 1H), 4.61 (d, 1H), 4.64 (d, 1H), 4.70-4.78 (m, 1H), 5.02-5.09 (m, 1H), 5.64-5.75 (m, 1H), 6.94-7.44 (m, 17H), 8.39-8.45 (m, 1H), 8.50-8.58 (m, 1H).

Example 18

N-((2*R*)-2-({[4-((2*R*,3*R*)-3-[(2-hydroxy-2-phenylethyl)thio]-4-oxo-1-phenylazetidin-2-yl)phenoxy]acetyl}amino)-2-phenylacetyl)-L-serine

[4-((2*R*,3*R*)-3-{[(5,5-Dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl]thio}-4-oxo-1-phenylazetidin-2-yl)phenoxy]acetic acid (0.030 g, 0.056 mmol), *tert*-butyl *N*-[(2*R*)-2-amino-2-phenylacetyl]-*O*-(*tert*-butyl)-*L*-serinate (0.024 g, 0.068 mmol) and *N*-methyl-morpholine (0.017 g, 0.169 mmol) were dissolved in CH₂Cl₂ (5 ml). After 5 minutes, TBTU (0.023 g, 0.073 mmol) was added. Full conversion to the corresponding *tert*-butyl ester was obtained after 1 h. The reaction mixture was purified on silica gel and eluted with EtOAc/ CH₂Cl₂ (25/75). The pure fractions were collected and concentrated. The residue was dissolved in 3 ml CH₂Cl₂ and 1 ml TFA. The reaction mixture was stirred overnight and concentrated. The remaining trace of TFA was azeotropically removed by co-evaporation with toluene (3 ml). The crude acid was dissolved in MeOH (4 ml) and NaBH₄ (0.009 g, 0.225 mmol) was added. Full conversion to the corresponding alcohol was obtained after 5 minutes. The reaction was quenched by the addition of 0.1M NH₄OAc buffer (2 ml). Purification by preparative HPLC using a gradient of 20-50% CH₃CN in 0.1M NH₄OAc buffer as mobile phase followed by freeze-drying of the pure fractions afforded the title compound (0.022 g, 58%) as a colourless solid. *M/z*: 668.5 (*M*-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 2.89-2.94 (m, 2H), 3.30-3.35 (m, 1H), 3.46-3.50 (m, 1H), 3.94-4.00 (m, 1H), 4.25-4.27 (m, 1H), 4.60 (d, 1H), 4.65 (d, 1H), 4.69-4.77 (m, 1H), 5.00-5.05 (m, 1H), 5.59-5.66 (m, 1H), 6.94-7.39 (m, 19H), 8.20-8.25 (m, 1H), 8.50-8.58 (m, 1H).

Example 19

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-*D*-alanyl-*D*-valine

25

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-*D*-alanine (0.040g, 0.072 mmol) was dissolved in CH₂Cl₂ (5 ml). *tert*-Butyl *D*-valinate hydrochloride (0.018 g, 0.087 mmol), *N*-methylmorpholine (0.022 g, 0.216 mmol) and TBTU (0.030 g, 0.094 mmol) were added. The *tert*-butyl ester was obtained after 1.5 h. The reaction mixture was purified on silica gel and eluted with CH₂Cl₂/EtOAc (75/25). The pure fractions were collected and concentrated. The residue was dissolved in CH₂Cl₂ (4 ml) and TFA (1 ml). The reaction mixture was stirred for 3 h and concentrated. The crude acid was dissolved in MeOH (4 ml) and NaBH₄ (0.011 g, 0.289 mmol) was added. Full conversion to the corresponding alcohol was obtained after 5 minutes. The reaction was quenched by the addition of 0.1M NH₄OAc buffer (1 ml). The reaction mixture was concentrated and the residue was purified by preparative HPLC using an eluent of 20-50% CH₃CN in 0.1M NH₄OAc buffer. This gave the title compound (0.039 g, 82%) as a colourless solid. *M/z*: 654.5 (*M*-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 0.79 (d, 6H), 1.21 (d, 3H), 1.96-2.05 (m, 1H), 2.85-2.95 (m, 2H), 3.91-3.97 (m, 1H), 4.22-4.29 (m, 1H), 4.35-4.42 (m, 1H), 4.48 (d, 1H), 4.53 (d, 1H), 4.70-4.78 (m, 1H), 5.02-5.04 (m, 1H), 6.94-7.36 (m, 12H), 7.70-7.77 (m, 1H), 8.15 (d, 1H).

Example 20

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-*D*-alanyl-*D*-alanine

50

- 5 *N*-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-alanine (0.040g, 0.072 mmol) was dissolved in CH₂Cl₂ (5 ml). *tert*-Butyl D-alaninate hydrochloride (0.016 g, 0.087 mmol), *N*-methylmorpholine (0.022 g, 0.216 mmol) and TBTU (0.030 g, 0.094 mmol) were added. Full conversion to the *tert*-butyl-ester was obtained after 1.5 h. The reaction mixture was purified on silica gel and eluted with
- 10 EtOAc/ CH₂Cl₂ (25/75). The pure fractions were collected and concentrated. The residue was dissolved in CH₂Cl₂ (3 ml) and TFA (1 ml). Full conversion to the acid was obtained after 2 h at room temperature. The reaction mixture was concentrated and the remaining trace of TFA was azeotropically removed by co-evaporation with toluene (3 ml). The crude acid was dissolved in MeOH (3 ml) and NaBH₄ (0.011 g, 0.289 mmol) was added. Full conversion to
- 15 the corresponding alcohol was obtained after 5 minutes. The reaction was quenched by the addition of 0.1M NH₄OAc buffer (1 ml). The reaction mixture was concentrated and the residue purified by preparative HPLC using a gradient of 20-50% CH₃CN in 0.1M NH₄OAc buffer as mobile phase. Freeze-drying of the pure fractions gave the title compound (0.039 g, 86%) as a colourless solid. M/z: 626.4 (M-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 1.17-1.22
- 20 (m, 6H), 2.82-2.96 (m, 2H), 3.90-3.98 (m, 1H), 4.25-4.35 (m, 2H), 4.48-4.54 (m, 2H), 4.70-4.78 (m, 1H), 5.02-5.06 (m, 1H), 6.94-7.37 (m, 12H), 7.85-7.92 (m, 1H), 8.16 (d, 1H).

Example 21

25

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-alanyl-D-lysine

30

- N*-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-alanine (0.040g, 0.072 mmol) was dissolved in CH₂Cl₂ (5 ml). *tert*-Butyl *N*⁶-(*tert*-butoxycarbonyl)-D-lysinate hydrochloride (0.029 g, 0.087 mmol), *N*-methylmorpholine (0.022 g, 0.216 mmol) and TBTU (0.030 g, 0.094 mmol) were added.
- 35 After 1.5 h full conversion to the ester was obtained. The reaction mixture was purified on silica gel and eluted with EtOAc/ CH₂Cl₂ (25/75). The pure fractions were collected and concentrated. The residue was dissolved in CH₂Cl₂ (3 ml) and TFA (1 ml) and stirred overnight and the corresponding acid was formed. The reaction mixture was concentrated and the remaining trace of TFA was azeotropically removed by co-evaporation with toluene (3
- 40 ml). The crude acid was dissolved in MeOH (3 ml) and NaBH₄ (0.011 g, 0.288 mmol) was added. The reduction was completed after 5 minutes. The reaction was quenched by the addition of 0.1M NH₄OAc buffer (1 ml). The mixture was concentrated and the residue purified by preparative HPLC using a gradient of 10-50% CH₃CN in 0.1M NH₄OAc buffer as eluent. Freeze-drying of the pure fractions afforded the title compound (0.039 g, 79%) as a
- 45 colourless solid. M/z: 685.1. ¹H NMR [(CD₃)₂SO], 400 MHz] δ 1.21 (d, 3H), 1.22-1.69 (m, 6H), 2.62-2.70 (m, 2H), 2.82-2.93 (m, 2H), 3.78-3.84 (m, 1H), 4.22-4.33 (m, 2H), 4.48 (d, 1H), 4.52 (d, 1H), 4.70-4.78 (m, 1H), 5.01-5.06 (m, 1H), 6.95-7.37 (m, 12H), 7.56-7.63 (m, 1H), 8.27 (d, 1H).

50

Example 22

***N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-valyl-D-serine**

5

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-valine (11 mg, 18.9 μ mol) was dissolved in DCM (2 ml). *tert*-Butyl *O*-(*tert*-butyl)-D-serinate hydrochloride (6.3 mg, 24.8 μ mol) and *N*-methylmorpholine (10 μ l, 91 μ mol) were added. After 5 minutes, TBTU (8.2 mg, 25.5 μ mol) was added and the reaction mixture, a white suspension, was stirred overnight. The formation of the ester was confirmed. *M/z*: 780.5 (M-H). The solvent was removed under reduced pressure. The residue was dissolved in formic acid (1 ml) and stirred at 50°C for 5 h and at ambient temperature overnight. LC-MS analysis showed the formation of the formate adduct of the product. *M/z*: 698.2 (M+H) and 696.2 (M-H). The formic acid was removed under reduced pressure, toluene (3x1 ml) was used to assist this removal. The yellowish oily residue was dissolved in methanol (1 ml) and triethylamine (150 μ l, 0.12 mmol) was added. The reaction mixture was stirred for 1 h. The formate was hydrolyzed; *M/z*: 670.1 (M+1) and 668.0 (M-1). Sodium borohydride (8.4 mg, 0.22 mmol) was added to the methanol solution. The mixture was stirred for 5 minutes. Ammonium acetate (7 mg) was added and the solvent was removed under reduced pressure. The residue was purified with preparative HPLC on a C8 column. A gradient from 20 to 50 % MeCN in 0.1M ammonium acetate buffer was used as eluent. After freeze-drying, the title compound was obtained as a white solid (3.8 mg, 30 %). H-NMR (400 MHz, DMS-*d*₆): 0.80 (m, 6H), 1.95-2.05 (m, 1H), 2.87-2.93 (m, 2H), 3.46-3.60 (m, 2H), 3.86 (brs, 1H), 4.19-4.32 (m, 2H), 4.59 (br, 2H), 4.69-4.77 (m, 1H), 5.05 (m, 1H), 6.96 (d, 2H), 7.06-7.18 (m, 4H), 7.20-7.26 (m, 2H), 7.30-7.39 (m, 4H), 7.70-7.77 (brs, 1H), 7.96-8.01 (d, 1H). *M/z*: 670.1(M-H).

30

Example 23

***N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-valine**

35

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-valine (0.022 g, 0.034 mmol) was dissolved in methanol (2 ml). NaBH₄ (0.0025 g, 0.066 mmol) was added and when the reaction was complete according to LC-MS a few drops of acetic acid was added. The solvent was removed under reduced pressure and the residue was purified by preparative HPLC on a Kromasil C8- column using 35% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, 0.015 g (68 %) of the title product was obtained. NMR (400 MHz, CD₃COOD) 0.90 (d, 3H), 0.93 (d, 3H), 2.10-2.20 (m, 1H), 2.90-3.06 (m, 2H), 3.99 (s, 2H), 4.03 (d, 0.5H), 4.05 (d, 0.5H), 4.27-4.34 (m, 1H), 4.60 (s, 2H), 4.79-4.84 (m, 1H), 4.89 (d, 0.5H), 4.91 (d, 0.5H), 6.95-7.03 (m, 4H), 7.06 (d, 2H), 7.25-7.30 (m, 2H); 7.30-7.37 (m, 4H).

50

Example 24

(2*R*)-cyclohexyl[(*N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl)amino]acetic acid

5

(2*R*)-cyclohexyl[(*N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl)amino]acetic acid (0.0085 g, 0.013 mmol) was dissolved in methanol (2 ml). NaBH₄ (0.006 g, 0.159 mmol) was added and when the reaction was complete according to LC-MS a few drops of acetic acid was added. The solvent was removed under reduced pressure and the residue was purified by preparative HPLC on a Kromasil C8- column using a stepwise gradient of 35%, 40% then 50% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, 0.007 g (82 %) of the title product was obtained. NMR (500 MHz, CD₃COOD) 1.05-1.33 (m, 5H), 1.60-1.87 (m, 6H), 2.91-3.07 (m, 2H), 3.94-4.06 (m, 3H), 4.32 (d, 1H), 4.60 (s, 2H), 4.79-4.86 (m, 1H), 4.90 (d, 0.5H), 4.92 (d, 0.5H), 6.96-7.03 (m, 4H), 7.03-7.08 (brd, 2H), 7.25-7.31 (m, 2H), 7.31-7.38 (m, 4H)

20 **Example 25**

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-(trimethylsilyl)alanine

25

Ethyl 3-(trimethylsilyl)alaninate (20mg, 0.106mmol) was dissolved in 1.5ml Et₃N, 0.2ml MeOH and 0.2ml H₂O and stirred for 5 days. The solvent was evaporated under reduced pressure at 40°C. Et₃N, 1ml, was added and evaporated.

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (20mg, 0.037mmol) was dissolved in 1.5ml dry DMF. *N*-Methylmorpholine (0.010ml, 0.109mmol) and TBTU (15mg, 0.047mmol) were added and the mixture was stirred for 1.5h. The Et₃N-salt of the above hydrolyzed amino acid was added and the mixture was stirred for 3h. A few drops of water were added and the mixture was stirred for 15 min. MeOH (2ml) and NaBH₄ (ca 15mg) were added. After 15 min ca 20mg NH₄Ac was added. The mixture was left overnight and was purified using preparative HPLC on a C8 column. A gradient from 20-50% MeCN in 0.1M ammonium acetate was used as mobile phase. A pure product fraction was collected and lyophilized. Mass: 14.5mg. The solid was placed in the vacuum oven at 40°C for 5h. M/z: 684 (M-1). NMR (400 MHz, DMSO-*d*₆): 8.23 (t, 1H), 7.88-7.98 (m, 1H), 7.30-7.38 (m, 4H), 7.20-7.25 (m, 2H), 7.05-7.18 (m, 4H), 6.98 (d, 2H), 5.02-5.07 (m, 1H), 4.67-4.76 (m, 1H), 4.51 (s, 2H), 4.24-4.33 (m, 1H), 4.05-4.15 (m, 1H), 3.65-3.80 (m, 2H), 2.82-2.98 (m, 2H), 0.85-1.03 (m, 2H), 0.04 (s, 9H).

45 **Example 26**

N-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-tyrosine

- 5 A mixture of 3-(R)-4-(R)-1-(4-Fluorophenyl)-3-[(4-fluorobenzoyl)methylthio]-4-{4-[N-(carboxymethyl) carbamoylmethoxy]phenyl}azetidin-2-one (0.0227g, 0.042 mmol), (R)-tyrosine *tert*-butylester (0.0144g, 0.061 mmol), N-methylmorpholin (0.012 ml, 0.111 mmol) in DCM (2ml) was stirred at room temperature. TBTU (0.018 g, 0.056 mmol) was added and the mixture was stirred overnight. Trifluoroacetic acid (0.65 ml) was added and after a couple of
10 hours the hydrolysis was complete. The solvent was removed under reduced pressure and the residue was purified by preparative HPLC on a Kromasil C8- column using a gradient of 5-100% MeCN in 0.15% trifluoroacetic acid buffer as eluent. After removing the solvent under reduced pressure the compound (M/z 704.1) was dissolved in methanol (2 ml). NaBH₄ (0.004 g, 0.105 mmol) was added and when the reaction was complete a few drops of acetic acid was
15 added. The solvent was removed under reduced pressure and the residue was purified by preparative HPLC on a Kromasil C8- column using 35% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, 0.021 g (71 %) of the title product was obtained. NMR (400 MHz, CD₃COOD) 2.85-3.10 (m, 4H), 3.90 (ABq, 2H), 4.03 (d, 0.5H), 4.05 (d, 0.5H), 4.51 (dd, 1H), 4.55 (ABq, 2H), 4.79-4.84 (m, 1H), 4.89 (d, 0.5H), 4.90 (d, 0.5H), 6.62-6.67 (m,
20 2H), 6.95-7.05 (m, 8H), 7.25-7.37 (m, 6H)

Example 27

- 25 N-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-proline

- A mixture of 3-(R)-4-(R)-1-(4-Fluorophenyl)-3-[(4-fluorobenzoyl)methylthio]-4-{4-[N-(carboxymethyl) carbamoylmethoxy]phenyl}azetidin-2-one (0.0197g, 0.036 mmol), (R)-proline *tert*-butylester (0.0118 g, 0.069 mmol), N-methylmorpholin (0.012 ml, 0.111 mmol) in DCM (2ml) was stirred at room temperature. TBTU (0.018 g, 0.056 mmol) was added and the mixture was stirred overnight. Trifluoroacetic acid (0.65 ml) was added and after a couple of
30 hours the hydrolysis was complete. The solvent was removed under reduced pressure and the residue was purified by preparative HPLC on a Kromasil C8- column using a gradient of 5-100% MeCN in 0.15% trifluoroacetic acid buffer as eluent. After removing the solvent under reduced pressure the compound (M/z 638.08) was dissolved in methanol (2 ml). NaBH₄ (0.004 g, 0.106 mmol) was added and when the reaction was complete a few drops of acetic acid was added. The solvent was removed under reduced pressure and the residue was
35 purified by preparative HPLC on a Kromasil C8- column using 35% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, 0.020 g (85 %) of the title product was obtained. NMR (400 MHz, CD₃COOD) 1.80-1.94 (m, 1H), 1.94-2.10 (m, 1.5H), 2.12-2.35 (m, 1.5H), 2.90-3.06 (m, 2H), 3.49-3.67 (m, 2H), 3.85 (d, 0.5H), 4.00-4.07 (m, 1.5H), 4.14-4.24 (m, 1H), 4.35 (dd, 0.5H), 4.41 (brd, 0.5H), 4.57 (s, 1H), 4.59 (s, 1H), 4.78-4.84
40 (m, 1H), 4.89 (d, 0.5H), 4.91 (d, 0.5H), 6.95-7.08 (m, 6H), 7.25-7.38 (m, 6H).

Example 28

- 50 N-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-L-threonine

5

Example 29

15

20

Example 30

40

45

2.90-3.07 (m, 2H), 3.89-4.08(m, 3H), 4.55 (dd, 1H), 4.60 (ABq, 2H), 4.78-4.85 (m, 1H), 4.90 (d, 0.5H), 4.92 (d, 0.5H), 6.96-7.09 (m, 6H), 7.24-7.38 (m, 6H).

5 Example 31

N-([4-((2R,3R)-1-(4-fluorophenyl)-3-([2-(4-fluorophenyl)-2-hydroxyethyl]thio)-4-oxoazetidin-2-yl)phenoxy]acetyl)glycyl-L-methionine

- 10 N-([4-((2R,3R)-1-(4-fluorophenyl)-3-([2-(4-fluorophenyl)-2-oxoethyl]thio)-4-oxoazetidin-2-yl)phenoxy]acetyl)glycyl-L-methionine (0.015 g, 0.022 mmol) was dissolved in methanol (2 ml). NaBH₄ (0.004 g, 0.106 mmol) was added and when the reaction was complete a few drops of acetic acid were added. The solvent was removed under reduced pressure and the residue was purified by preparative HPLC on a Kromasil C8- column using 35% MeCN in 15 0.1M ammonium acetate buffer as eluent. After freeze-drying, 0.015 g (99%) of the title product was obtained. NMR (400 MHz, CD₃COOD) 1.86-2.00 (m, 1H), 2.04 (s, 3H); 2.07-2.18 (m, 1H), 2.45-2.51 (m, 2H), 2.90-3.08 (m, 2H), 3.97 (s, 2H), 4.04 (d, 0.5H), 4.06 (d, 0.5H), 4.36-4.43 (m, 1H), 4.60 (s, 2H), 4.77-4.85 (m, 1H), 4.90 (d, 0.5H), 4.92 (d, 0.5H), 6.95-7.08 (m, 6H), 7.25-7.38 (m, 6H).

20 Example 32

N-([4-((2R,3R)-1-(4-fluorophenyl)-3-([2-hydroxy-2-(4-methoxyphenyl)ethyl]thio)-4-oxoazetidin-2-yl)phenoxy]acetyl)glycyl-D-valine

- 25 N-([4-((2R,3R)-1-(4-fluorophenyl)-3-([2-(4-methoxyphenyl)-2-oxoethyl]thio)-4-oxoazetidin-2-yl)phenoxy]acetyl)glycyl-D-valine (0.015g, 0.023 mmol) was dissolved in methanol (3 ml). NaBH₄ (0.006 g, 0.158 mmol) was added and when the reaction was complete a few drops of 30 acetic acid were added. The solvent was removed under reduced pressure and the residue was purified by preparative HPLC on a Kromasil C8- column using 25% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, 0.014 g (93%) of the title product was obtained. NMR (500 MHz, CD₃COOD) 0.93 (d, 3H), 0.95 (d, 3H), 2.12-2.22 (m, 1H), 2.91-3.07 (m, 2H), 3.75 (s, 1.5H), 3.76 (s, 1.5H), 3.94-4.06 (m, 3H), 4.33 (d, 1H), 4.60 (s, 35 2H), 4.72-4.78 (m, 1H), 4.81-4.88 (m, 1H), 6.79-6.83 (m, 2H), 6.97-7.08 (m, 4H), 7.20-7.36 (m, 6H).

40 Example 33

N-([4-((2R,3R)-1-(4-fluorophenyl)-3-([2-(4-fluorophenyl)-2-hydroxyethyl]thio)-4-oxoazetidin-2-yl)phenoxy]acetyl)glycyl-D-leucine

- 45 A mixture of 3-(R)-4-(R)-1-(4-Fluorophenyl)-3-[(4-fluorobenzoyl)methylthio]-4-[4-[N-(carboxymethyl) carbamoylmethoxy]phenyl]azetidin-2-one (0.015g, 0.028 mmol), *tert*-butyl D-leucinate hydrochloride (0.010g, 0.045 mmol), N-methylmorpholin (0.0092 ml, 0.083 mmol) in DCM (2ml) was stirred at room temperature. TBTU (0.012 g, 0.037 mmol) was added and the mixture was stirred overnight. Trifluoroacetic acid (1.0 ml) was added and after 50 2 h the solvent was removed under reduced pressure and the residue was purified by

preparative HPLC on a Kromasil C8- column using a gradient of 5-100% MeCN in 0.15% trifluoroacetic acid buffer as eluent. After removing the solvent under reduced pressure the compound (M/z 654.25) was dissolved in methanol (2 ml). NaBH₄ (0.005 g, 0.132 mmol) was added and the mixture was stirred for 5 minutes. A few drops of acetic acid was added and the solvent was removed under reduced pressure and the residue was purified by preparative HPLC on a Kromasil C8- column using 35% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, 0.014 g (76 %) of the title product was obtained. NMR (400 MHz, CD₃COOD) 0.90-0.94 (m, 6H), 1.52-1.74 (m, 3H), 2.90-3.07 (m, 2H), 3.97 (s, 2H), 4.02 (d, 0.5H), 4.04 (d, 0.5H), 4.40 (d, 0.5H), 4.42 (d, 0.5H), 4.59 (s, 2H), 4.78-4.84 (m, 1H), 4.89 (d, 0.5H), 4.91 (d, 0.5H), 6.95-7.08 (m, 6H), 7.24-7.37 (m, 6H).

Example 34

15 N-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-hydroxy-2-(4-methoxyphenyl)ethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-lysine

A mixture of N-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-methoxyphenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.0177g, 0.032 mmol) *tert*-butyl N⁶-(*tert*-butoxycarbonyl)-D-lysinate hydrochloride (0.0141 g, 0.042 mmol), N-methylmorpholin (0.0106 ml, 0.096 mmol) in DCM (2ml) was stirred at room temperature. TBTU (0.013 g, 0.042 mmol) was added and the mixture was stirred for 2 h. Additional *tert*-butyl N⁶-(*tert*-butoxycarbonyl)-D-lysinate hydrochloride (0.004 g, 0.012 mmol) was added and after 0.5 h. TBTU (0.008g, 0.025 mmol) was added and the mixture was stirred for additional 10 minutes. Trifluoroacetic acid (0.65 ml) was added and after 3 h the solvent was removed under reduced pressure. The residue was purified by preparative HPLC on a Kromasil C8- column using a gradient of 5-100% MeCN in 0.15% 1M ammonium acetate buffer as eluent. After removing the solvents under reduced pressure, the intermediate (M/z 681.4) was dissolved in methanol (3 ml). NaBH₄ (0.008 g, 0.211mmol) was added and when the reaction was complete a few drops of acetic acid was added. The solvent was removed under reduced pressure and the residue was purified by preparative HPLC on a Kromasil C8- column using 35% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, 0.020 g (91 %) of the title product was obtained. NMR (500 MHz, CD₃COOD) 1.41-1.52 (m, 2H), 1.58-1.78 (m, 3H), 1.90-2.02 (m, 1H), 2.86-3.08 (m, 4H), 3.76 (s, 1.5H), 3.77 (s, 1.5H), 3.90-4.06 (m, 3H), 4.48 (dd, 1H), 4.60-4.62 (m, 2H), 4.73-4.78 (m, 1H), 4.83-4.89 (m, 1H), 6.80-6.84 (m, 2H), 6.98-7.08 (m, 4H), 7.21-7.31 (m, 4H), 7.31-7.36 (m, 2H), 8.20 (d, NH), 8.54 (t, NH).

Example 35

45 N²-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-L-glutaminy-D-phenylalanine

N²-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-L-glutamine (15mg, 0.0244 mmol), *tert*-butyl D-phenylalaninate hydrochloride (8 mg, 0.0310 mmol) and N-methylmorpholine (10 mg, 0.099

mmol) were dissolved in methylene chloride (0.5 ml). TBTU (10 mg, 0.0313 mmol) was added and the mixture was stirred for 1 h at room temperature. The solvent was evaporated and the residue was dissolved in formic acid (0.5 ml). The mixture was heated to 45-50 °C and stirred at this temperature for 6 h. The reaction mixture was evaporated under reduced pressure. Methanol (5 ml) was added and evaporated. The residue was dissolved in methanol (1 ml). Two drops of TEA was added and the mixture was stirred overnight at room temperature. The solvent was evaporated and the residue was purified by preparative HPLC using acetonitrile/ammonium acetate buffer (40:60) as eluent. After freeze-drying 12 mg (64%) of the title compound was obtained. ¹H-NMR, 300 MHz, DMSO): 1.55-1.97 (m, 4H), 2.77-3.10 (m, 4H), 4.20-4.38 (m, 3H), 4.49 (s, 2H), 4.66-4.78 (m, 1H), 4.99-5.07 (m, 1H), 6.70 (s, 1H), 6.88-7.40 (m, 19H), 8.00-8.20 (m, 2H).

Example 36

*N*²-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-L-glutaminyl-D-tyrosine

*N*²-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-L-glutamine (22 mg, 0.0359 mmol), *tert*-butyl D-tyrosinate hydrochloride (10 mg, 0.0421 mmol) and *N*-methylmorpholine (14 mg, 0.138 mmol) were dissolved in methylene chloride (0.5 ml). TBTU (14 mg, 0.0436 mmol) was added and the mixture was stirred overnight at room temperature. The solvent was evaporated and the residue was dissolved in formic acid (1 ml). The mixture was stirred at 45-50 °C for 4 h and was then evaporated under reduced pressure. Methanol (10 ml) was added and evaporated. The residue was dissolved in methanol (2 ml). Three drops of TEA were added and the mixture was stirred overnight at room temperature. The solvent was evaporated and the residue was purified by preparative HPLC using acetonitrile/ammonium acetate buffer (40:60) as eluent. After freeze-drying 15 mg (54%) of the title compound was obtained. ¹H-NMR, 300 MHz, DMSO): 1.60-2.00 (m, 4H), 2.69-3.0 (m, 4H), 4.09-4.18 (m, 1H), 4.22-4.35 (m, 3H), 4.49 (s, 2H), 4.65-4.78 (m, 1H), 4.97-5.08 (m, 1H), 6.56 (d, 2H), 6.70 (s, 1H), 6.86-7.40 (m, 18H), 7.77-7.91 (m, 1H), 8.12 (d, 1H), 9.10 (bs, 1H).

Example 37

N-{[4-((2*R*,3*R*)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-cyclohexyl-D-alanine

N-{[4-((2*R*,3*R*)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (20mg, 0.0349 mmol) and *N*-methylmorpholine (20 mg, 0.198 mmol) were dissolved in methylene chloride (0.5 ml). TBTU (17 mg, 0.0530 mmol) was added and the mixture was stirred for 15 min at room temperature. 3-Cyclohexyl-D-alanine (9 mg, 0.0526 mmol) was added and the stirring was continued for 1 h at room temperature. The solvent was evaporated and the residue was dissolved in methanol (0.5 ml). NaBH₄ (10 mg, 0.264 mmol) was added and the mixture was stirred for 15 min at room temperature. Three drops of acetic acid was added to the reaction mixture. The product was isolated by

preparative HPLC using acetonitrile/ammonium acetate buffer (45:55) as eluent. After freeze-drying 3 mg (12%) of the title compound was obtained. The product was analyzed by LC/Micromass Q TOF micro MS technique. M/z: 728.1973 (calc. Mass 728.1964).

5

Example 38

10 *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-cyclohexyl-D-alanine

15 *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (30 mg, 0.0555 mmol) and *N*-methylmorpholine (30 mg, 0.296 mmol) were dissolved in DMF (0.5 ml). TBTU (23 mg, 0.0717 mmol) was added and the mixture was stirred for 15 min at room temperature. 3-Cyclohexyl-D-alanine (15 mg, 0.0876 mmol) was added and the stirring was continued overnight at room temperature. The solvent was evaporated under reduced pressure and the residue was dissolved in methanol (1 ml).
20 NaBH₄ (10 mg, 0.264 mmol) was added and the mixture was stirred for 10 min at room temperature. Three drops of acetic acid was added to the reaction mixture. The product was isolated by preparative HPLC using acetonitrile/ammonium acetate buffer (40:60) as eluent. After freeze-drying 19 mg (49%) of the title compound was obtained. ¹H-NMR, 300 MHz, DMSO): 0.68-0.93 (m, 2H), 1.0-1.75 (m, 11H), 2.78-3.00 (m, 2H), 3.73 (s, 2H), 4.00-4.14 (m, 25 1H), 4.23-4.35 (m, 1H), 4.51 (s, 2H), 4.65-4.78 (m, 1H), 4.99-5.09 (m, 1H), 6.90-7.44 (m, 12H), 7.72-7.86 (m, 1H), 8.26 (t, 1H).

Example 39

30 *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-L-phenylalanine

35 *N*-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-L-phenylalanine (7mg, 0.010mmol) was dissolved in 1.5ml MeOH and NaBH₄ (2mg, 0.053mmol) was added. The mixture was stirred for 30 min and quenched with an excess of NH₄Ac. LC/MS showed ca 15% starting material. The mixture was diluted
40 with H₂O and purified using preparative chromatography on a C8 column (25x300mm). A gradient from 20% to 40% MeCN in 0.1M ammonium acetate was used as mobile phase. The product fraction was collected and partly concentrated. The mixture was lyophilized to yield 2mg (29%). M/z: 688 (M-1). NMR (400 MHz, MeOD): 6.95-7.36 (m, 17H), 4.90 (dd, 1H), 4.53-4.59 (m, 4H), 4.03 (dd, 1H), 3.90 (q, 2H), 3.17 (dd, 1H), 2.90-3.05 (m, 3H).
45

Example 40

50 *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-4-methylleucine

- tert*-Butyl *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-4-methylleucinate (ca 26mg, 0.036mmol) was dissolved in 1.5 ml formic acid and heated at 40°C for 1.5 h. The formic acid was evaporated under reduced pressure. The intermediate acid was confirmed by LC/MS analysis. *M/z*: 668. The crude mixture was dissolved in 2 ml MeOH. NaBH₄ (15mg, 0.40mmol) was added and the mixture was stirred for 10 min. NH₄Ac (30mg) was added. The crude mixture was diluted with 1ml water and purified using preparative HPLC on a C8 column (50x300mm). A gradient from 20% to 40% MeCN in 0.1M ammonium acetate buffer was used as mobile phase. Lyophilization yielded 36mg white solid. NMR showed the presence of water and HOAc. The product was dried in the vacuum oven for 1.5h at 40°C. Mass 13mg (52%). *M/z*: 668 (M-1). NMR (400 MHz, DMSO): 8.22 (t, 1H), 7.53-7.65 (m, 1H), 7.30-7.38 (m, 4H), 7.20-7.25 (m, 2H), 7.05-7.17 (m, 4H), 6.98 (d, 2H), 5.02-5.06 (m, 1H), 4.68-4.76 (m, 1H), 4.51 (s, 2H), 4.24-4.32 (m, 1H), 4.01-4.09 (m, 1H), 3.60-3.77 (m, 2H), 2.82-2.98 (m, 2H), 1.64 (dd, 1H), 1.33 (dd, 1H), 0.83 (s, 9H).

Example 41

- N*-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-*D*-seryl-*D*-phenylalanine
- A solution of *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-*D*-serine (diastereomeric mixture) (0.008 g, 0.014 mmol), *D*-phenylalanine *tert*-butyl ester hydrochloride (0.005 g, 0.018 mmol) and *N*-Methylmorpholine (0.006 ml, 0.055 mmol) in DCM (3 ml) was stirred at RT for 5 min. TBTU (0.008 g, 0.025 mmol) was added. After 3 days the conversion to the ester (*M/z*: 776.1) was complete and the mixture was concentrated under reduced pressure. The residue was dissolved in formic acid (3 ml) and the solution was stirred for 25 h. The mixture was concentrated under reduced pressure and the residue was dissolved in MeOH (4ml) and TEA (1 ml). The solution was stirred at 40°C for 6 h. The solvent was removed under reduced pressure. The residue was purified by preparative HPLC using a gradient of 20-50% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, 0.005 g (53 % yield) of the title product was obtained as a white solid. *M/z*: 720.1. ¹H NMR (DMSO, 400 MHz): δ 2.78-2.96 (m, 3H), 3.03-3.11 (m, 1H), 3.44-3.60 (m, 2H), 4.20-4.36 (m, 3H), 4.51 (s, 2H), 4.67-4.76 (m, 1H), 5.01-5.06 (m, 1H), 5.70 (bs, 1H), 6.90-6.98 (m, 2H), 7.05-7.25 (m, 1H), 7.29-7.37 (m, 4H), 7.84-8.00 (m, 2H).

Example 42

- N*-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-*D*-seryl-*D*-serine

- A solution of *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-*D*-serine (0.008 g, 0.014 mmol), O-

tert-butyl-D-serine tert-butyl ester hydrochloride (0.005 g, 0.019 mmol) and N-Methylmorpholine (0.006 ml, 0.055 mmol) in DCM (3 ml) was stirred for 5 min. TBTU (0.008 g, 0.025 mmol) was added. After 3 days the conversion to the ester (M/z: 772.5) was complete and the mixture was concentrated under reduced pressure. The residue was dissolved in formic acid (3 ml) and the solution was stirred at RT for 26h. The mixture was concentrated under reduced pressure and the residue was dissolved in MeOH (4ml) and TEA (1 ml). The solution was stirred at 40°C for 6 h. The solvent was removed under reduced pressure. The residue was purified by preparative HPLC using a gradient of 20-50% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, 0.005 g (52 % yield) of the title compound was obtained as a white solid. M/z: 660.1. ¹H NMR (DMSO, 400 MHz): δ 2.83-2.95 (m, 2H), 3.42-3.66 (m, 4), 3.91-4.01 (m, 1H), 4.25-4.31 (m, 1H), 4.32-4.39 (m, 1H), 4.54 (ABq, 2H), 4.68-4.76 (m, 1H), 5.02-5.06 (m, 1H), 5.68 (bs, 1H), 6.94-7.00 (m, 2H), 7.05-7.18 (m, 4H), 7.19-7.26 (m, 2H), 7.29-7.39 (m, 4H), 7.74-7.81 (m, 1H), 7.99 (d, 1H).

15 Example 43

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-2-butylnorleucine

20

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-[[2-(4-fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-2-butylnorleucine (0.009 g, 0.012 mmol) was dissolved in MeOH (3 ml). NaBH₄ (0.007 g, 0.185 mmol) was added and the mixture was stirred for 10 min. Ammonium acetate buffer (0.1M, 3 ml) was added and most of the methanol was removed under reduced pressure. The remaining solution was purified by preparative HPLC, using a gradient of 20-50% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, 0.008 g (94 % yield) of the title compound was obtained as a white solid. M/z: 712.1. ¹H NMR (DMSO, 400 MHz): δ 0.73-0.83 (m, 6H), 0.89-1.22 (m, 8H), 1.59-1.71 (m, 2H), 1.94-2.06 (m, 2H), 2.83-2.96 (m, 2H), 3.73 (d, 2H), 4.23-4.28 (m, 1H), 4.51 (s, 2H), 4.68-4.76 (m, 1H), 5.02-5.07 (m, 1H), 5.68 (bs, 1H), 6.94-7.00 (m, 2H), 7.05-7.18 (m, 4H), 7.19-7.26 (m, 2H), 7.29-7.39 (m, 4H), 7.53 (s, 1H), 8.30-8.36 (m, 1H).

35 Example 44

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*S*-methyl-L-cysteine

40

A solution of *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.015 g, 0.028 mmol), *tert*-butyl *S*-methyl-L-cysteinate (0.014 g, 0.073 mmol) and N-methylmorpholine (0.012 ml, 0.109 mmol) in DCM (5 ml) was stirred for 5 min. TBTU (0.013 g, 0.042 mmol) was added. After 20 h, the conversion to the ester (M/z: 714.1) was complete and the mixture was concentrated under reduced pressure. The residue was dissolved in formic acid (3 ml) and the solution was stirred at 40°C for 22 h. The mixture was diluted with toluene (2 ml) and the solvent was again removed under reduced pressure. The residue was dissolved in MeOH (4 ml) and NaBH₄ was added in small portions to the solution (a total of 0.035 g, 0.925 mmol) until the reduction was complete. Ammonium acetate buffer (0.1M, 3 ml) was added and most of the methanol was

removed under reduced pressure. The remaining solution was purified by preparative HPLC, using a gradient of 20-50% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, 0.013 g (72 % yield) of the title compound was obtained as a white solid.

M/z: 660.3. ¹H NMR (DMSO, 400 MHz): δ 2.66-2.74 (m, 1H), 2.82-3.02 (m, 3H), 3.67-3.82 (m, 2H), 4.02-4.12 (m, 1H), 4.24-4.31 (m, 1H), 4.52 (s, 2H), 4.68-4.77 (m, 1H), 5.01-5.07 (m, 1H), 6.95-7.02 (m, 2H), 7.05-7.18 (m, 4H), 7.19-7.26 (m, 2H), 7.29-7.40 (m, 4H), 7.73-7.82 (m, 1H), 8.30-8.37 (m, 1H).

Example 45

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-L-isoleucine

15 A solution of *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.015 g, 0.028 mmol), L-isoleucine *t*-butyl ester hydrochloride (0.008 g, 0.036 mmol) and *N*-methylmorpholine (0.012 ml, 0.109 mmol) in DCM (5 ml) was stirred for 5 min. TBTU (0.012 g, 0.036 mmol) was added. After 22 h, the conversion to the ester (M/z: 710.2) was complete and the mixture was concentrated under
20 reduced pressure. The residue was dissolved in formic acid (3 ml) and the solution was stirred at 40°C for 20h. The mixture was diluted with toluene (2 ml) and the solvent was removed under reduced pressure. The residue was dissolved in MeOH (4 ml) and NaBH₄ was added in small portions to the solution (a total of 0.060 g, 1.59 mmol) until the reduction was complete.
25 under reduced pressure. The remaining solution was purified by preparative HPLC, using a gradient of 20-50% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, 0.010 g (57 % yield) of the title compound was obtained as a white solid. M/z: 656.2. ¹H
NMR (DMSO, 400 MHz): δ 0.76-0.84 (m, 6H), 1.05-1.17 (m, 1H), 1.32-1.44 (m, 1H), 1.67-1.78 (m, 1H), 2.82-2.95 (m, 2H), 3.78 (d, 2H), 4.03-4.11 (m, 1H), 4.23-4.29 (m, 1H), 4.51 (s,
30 2H), 4.68-4.77 (m, 1H), 5.01-5.07 (m, 1H), 6.94-7.01 (m, 2H), 7.05-7.26 (m, 6H), 7.29-7.39 (m, 4H), 7.79-7.89 (m, 1H), 8.22-8.28 (m, 1H).

Example 46

35 *N*-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-L-alanyl-D-valine

40 A solution of *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-L-alanine (0.015 g, 0.027 mmol), D-valine *tert*-butyl ester hydrochloride (0.008 g, 0.038 mmol) and *N*-methylmorpholine (0.030 ml, 0.272 mmol) in DCM (4 ml) was stirred for 5 min. TBTU (0.013 g, 0.041 mmol) was added. After 3 h, the conversion to the ester (M/z: 710.2) was completed and TFA (3 ml) was added. After 4h, the mixture was diluted with toluene (2 ml) and the solvent was removed under reduced pressure.
45 The residue was dissolved in MeOH (4 ml) and NaBH₄ was added in small portions to the solution (a total of 0.065 g, 1.72 mmol) until the reduction was complete. Ammonium acetate buffer (0.1M, 3 ml) was added and most of the methanol was removed under reduced pressure. The remaining solution was purified by preparative HPLC, using a gradient of 20-50% MeCN in a 0.1M ammonium acetate buffer as eluent. After freeze-drying, 0.014 g (77 %
50 yield) of the title compound was obtained as a white solid. M/z: 656.1. ¹H NMR (DMSO, 400

MHz): δ 0.75-0.83 (m, 6H), 1.22 (d, 3H), 1.95-2.07 (m, 1H), 3.83-3.96 (m, 2H), 3.98-4.06 (m, 1H), 4.24-4.31 (m, 1H), 4.40-4.54 (m, 3H), 4.67-4.76 (m, 1H), 5.01-5.07 (m, 1H), 6.91-7.98 (m, 2H), 7.05-7.17 (m, 4H), 7.19-7.25 (m, 2H), 7.29-7.39 (m, 4H), 7.84-7.95 (m, 1H), 8.07 (d, 1H),

5

Example 47

1-[(*N*-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl)amino]cyclopentanecarboxylic acid

10

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.015 g, 0.028 mmol), NMM (0.012 ml, 0.109 mmol) and TBTU (0.011 g, 0.034 mmol) were dissolved in DMF (2 ml) at 30°C. After 30 min 1-amino-1-cyclopentanecarboxylic acid (0.004 g, 0.030 mmol, 97 %) was added and the mixture was stirred at 30°C for 1h. The reaction was quenched with water (0.2 ml) and the mixture was diluted with MeOH (2 ml). NaBH₄ (0.015 g, 0.397 mmol) was added and the mixture was stirred for 10 min. Ammonium acetate buffer (0.1M, 3 ml) was added and most of the methanol was removed under reduced pressure. The remaining solution was purified by preparative HPLC, using a gradient of 20-50% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, the title compound (0.009 g, 49 % yield) was obtained as a white solid. M/z: 654.0. ¹H NMR (DMSO, 400 MHz): δ 1.56-1.66 (m, 4H), 1.78-1.87 (m, 2H), 1.97-2.07 (m, 2H), 2.84-2.94 (m, 2H), 3.74 (d, 2H), 4.24-4.29 (m, 1H), 4.51 (s, 2H), 4.68-4.76 (m, 1H), 5.02-5.07 (m, 1H), 6.92-7.00 (m, 2H), 7.05-7.26 (m, 6H), 7.29-7.39 (m, 4H), 8.09-8.15 (m, 2H).

30 **Example 48**

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*N*-benzylglycine

35

TBTU (0.011 g, 0.034 mmol) was added to a solution of *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.015 g, 0.028 mmol) and NMM (0.020 ml, 0.182 mmol) in DMF (2 ml) at 30°C. After 30 min, *N*-benzylglycine (0.005 g, 0.030 mmol, 98 %) was added and the mixture was stirred at 30°C for 1h. The reaction was quenched with water (0.2 ml) and the mixture was diluted with MeOH (2 ml). NaBH₄ (0.015 g, 0.397 mmol) was added and the mixture was stirred for 10 min. Ammonium acetate buffer (0.1M, 3 ml) was added and most of the methanol was removed under reduced pressure. The remaining solution was purified by preparative HPLC, using a gradient of 20-50% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, the title compound (0.010 g, 53 % yield) was obtained as a white solid. M/z: 690.0. ¹H NMR (DMSO, 400 MHz): δ 2.83-2.94 (m, 2H), 3.91 (s, 1H), 3.98-4.09 (m, 3H), 4.25-4.29 (m, 1H), 4.50 (s, 2H), 4.54 (s, 1H), 4.62 (s, 1H), 4.68-4.76 (m, 1H), 5.02-5.07 (m, 1H), 6.92-7.02 (m, 2H), 7.05-7.40 (m, 15H), 8.14-8.21 (m, 1H).

45

Example 49

5 [(*N*-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl)amino](diphenyl)acetic acid

TBTU (0.016 g, 0.050 mmol) was added to a solution of *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.020 g, 0.037 mmol) and NMM (0.012 ml, 0.109 mmol) in DMF (2 ml) at 30°C. After 30min, 2,2-diphenylglycine (0.009 g, 0.037 mmol, 98 %) was added and the mixture was stirred at 30°C for 2.5 h. The reaction was quenched with water (0.2 ml) and the mixture was diluted with MeOH (2 ml). NaBH₄ (0.020 g, 0.529 mmol) was added and the mixture was stirred for 10 min. Ammonium acetate buffer (0.1M, 3 ml) was added and most of the methanol was removed under reduced pressure. The remaining solution was purified by preparative HPLC, using a gradient of 20-50% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, the title compound (0.012 g, 44 % yield) was obtained as a white solid. *M/z*: 752.0. ¹H NMR (DMSO, 400 MHz): δ 2.83-2.95 (m, 2H), 3.75 (d, 2H), 4.24-4.28 (m, 1H), 4.54 (s, 1H), 4.68-4.75 (m, 1H), 5.02-5.06 (m, 1H), 6.95-7.03 (m, 2H), 7.04-7.38 (m, 20H), 8.43-8.51 (m, 1H), 8.90 (s, 1H).

Example 50

25 *N*-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycylglycine

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycylglycine (0.011 g, 0.018 mmol) was dissolved in MeOH (3 ml). NaBH₄ (0.013 g, 0.344 mmol) was added and the mixture was stirred for 10 min. Ammonium acetate buffer (0.1M, 3 ml) was added and most of the methanol was removed under reduced pressure. The remaining solution was purified by preparative HPLC, using a gradient of 20-50% MeCN in a 0.1M ammonium acetate buffer as eluent. After freeze-drying, the title compound (0.011 g, 97 % yield) was obtained as a white solid. *M/z*: 600.0. ¹H NMR (DMSO, 400 MHz): 2.97-2.84 (m, 2H), 3.44-3.50 (m, 2H), 3.74 (d, 2H), 4.26-4.32 (m, 1H), 4.46-4.54 (m, 2H), 4.67-4.76 (m, 1H), 5.02-5.07 (m, 1H), 6.95-7.01 (m, 2H), 7.05-7.26 (m, 6H), 7.30-7.40 (m, 4H), 7.60-7.80 (m, 1H), 8.32-8.38 (m, 1H).

40 **Example 51**

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-valyl-L-serine

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-valine (6.9 mg, 11.8 μmol) was dissolved in DCM (3 ml). *tert*-Butyl *O*-(*tert*-butyl)-L-serinate hydrochloride (3.7 mg, 14.2 μmol) and *N*-methylmorpholine (5.5 μl, 50 μmol) were added. After 5 minutes, TBTU (4.6 mg, 14.3 μmol) was added and the reaction mixture was stirred overnight. The formation of the ester was confirmed. *M/z*:

780.57 (M-1). The mixture was extracted between DCM (5 ml) and aqueous KHSO₄ (5 ml, pH of 2). The organic phase was washed with aqueous NaHCO₃ (5 ml, pH of 9). The aqueous phase was extracted with DCM (2x5 ml). The combined organic phases were dried over Na₂SO₄, filtered and concentrated. Formic acid (1.5 ml) was added and the reaction mixture was stirred overnight. The formate of the intermediate acid was obtained. The solvent was removed under reduced pressure, toluene (3x1 ml) was added and evaporated. The residue was dissolved in MeOH (1.5 ml) and triethylamine (90 µl, 0.65 mmol) was added and the reaction mixture was stirred for 1 hour. Sodium borohydride (4.0 mg, 0.11 mmol) was added and the reaction mixture was stirred for 1 hour. Ammonium acetate (15 mg) was added. The solvent was removed under reduced pressure and the residue was purified on preparative HPLC using a C8 column. A gradient from 20% to 50 % MeCN in 0.1M ammonium acetate buffer was used as eluent. After lyophilisation, the title compound was obtained as a white solid (5.3 mg, 67 %). HRMS calcd for C₃₃H₃₅F₂N₃O₈S 671.2113, found 672.2192 [M+H]⁺.

15

Example 52

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}-*D*-valylglycine

20

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-[[2-(4-fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}-*D*-valine (11.8 mg, 0.02 mmol), *tert*-butyl glycinate hydrochloride (4.6 mg, 0.03 mmol) and *N*-methylmorpholine (10 µl, 0.09 mmol) were dissolved in DCM (1.5 ml). After 5 minutes, TBTU was added and the reaction mixture was stirred for 3.5 h. The formation of the ester was confirmed. M/z: 694.0 (M-H). The mixture was extracted between DCM (3 ml) and aqueous KHSO₄ (5 ml, pH of 3). The aqueous phase was extracted with DCM (2x5 ml). The combined organic phases were washed with water (2x5 ml), dried over Na₂SO₄, filtered and concentrated. Formic acid (3 ml) was added and the solution was heated at 40°C overnight. The solvent was removed under reduced pressure, toluene was added and evaporated. The residue was dissolved in methanol (3 ml) and sodium borohydride (8.5 mg, 0.23 mmol) was added. The reaction mixture was stirred for 15 minutes. The solvent was removed under reduced pressure. The residue was purified on preparative HPLC on a C8 column. A gradient from 20% to 50 % MeCN in 0.1M ammonium acetate buffer was used as eluent. After lyophilisation, the title compound was obtained as a white solid (6.2 mg, 48 %). HRMS calcd for C₃₂H₃₃F₂N₃O₇S 641.2007, found 642.2086 [M+H]⁺.

40 **Example 53**

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}-*D*-valyl-L-valine

45

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-[[2-(4-fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}-*D*-valine (11.5, 0.02 mmol), *tert*-butyl L-valinate hydrochloride (5.3 mg, 0.025 mmol) and *N*-methylmorpholine (10 µl, 0.091 mmol) were dissolved in DCM (1.5 ml). After 5 minutes, TBTU (7.8 mg, 0.024 mmol) was added and the reaction mixture was stirred overnight. Additional *tert*-butyl L-valinate hydrochloride (1.5 mg, 7.2 µmol), *N*-

methylmorpholine (6.5 μ l, 58 μ mol) and TBTU (2.0 mg, 6.2 μ mol, 0.31 eq) were added and the mixture was stirred for 2.5 h. The formation of the ester was confirmed. *M/z*: 736.1 (M-H). The reaction mixture was extracted between aqueous KHSO₄ (5 ml, pH of 3) and DCM (5 ml). The organic phase was dried over Na₂SO₄, filtered and concentrated. The residue was dissolved in formic acid (2 ml) and heated at 40 °C overnight. The solvent was removed under reduced pressure. Toluene was added and removed under reduced pressure. The residue was dissolved in MeOH (2 ml) and sodium borohydride (8.1 mg, 0.21 mmol) was added. The mixture was stirred for 30 minutes. Ammonium acetate (16 mg) was added and the solvent was removed under reduced pressure. The residue was purified with preparative HPLC on a C8 column. A gradient from 20% to 50 % MeCN in 0.1 M NH₄OAc buffer was used as eluent. After lyophilisation, the title compound was obtained as a white solid (7.3 mg, 54 %).

H-NMR (400 MHz, DMSO-*d*₆): 0.70 (d, 3H), 0.79 (d, 3H), 0.83 (d, 6H), 2.00 (m, 2H), 2.86-2.92 (m, 2H), 4.07 (brs, 1H), 4.25-4.29 (m, 1H), 4.40 (m, 1H), 4.59 (brs, 2H), 4.72 (m, 1H), 5.03 (d, 0.5H), 5.05 (d, 0.5H), 5.68 (brs, 1H), 6.95 (d, 2H), 7.06-7.16 (m, 4H), 7.19-7.24 (m, 2H), 7.29-7.37 (m, 4H), 7.80 (d, 1H), 8.05-8.15 (brs, 1H). *M/z*: 682.1 (M-H).

Example 54

20 *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-valyl-D-valine

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-Fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-valine (11.9 mg, 0.02 mmol), *tert*-butyl D-valinate hydrochloride (5.6 mg, 0.027 mmol) and *N*-methylmorpholine (10 μ l, 0.092 mmol) were dissolved in DCM (1.5 ml). After 5 minutes, TBTU (8.2 mg, 0.0255 mmol) was added and the reaction mixture was stirred overnight. Additional *tert*-butyl D-valinate hydrochloride (1.7 mg, 8.1 μ mol), *N*-methylmorpholine (6.5 μ l, 58 μ mol), TBTU (2 mg, 6.2 μ mol) were added and the mixture was stirred for 3 h. The formation of the ester was confirmed. *M/z*: 736.2 (M-H). The solvent was removed under reduced pressure and the residue was purified on silica gel (1 g) using DCM: MeOH (8:2) as eluent. The fractions were collected and concentrated. The residue was dissolved in formic acid (1 ml) and the resulting solution was stirred at 25-30 °C overnight. The solvent was evaporated and toluene was added and removed under reduced pressure. The residue was dissolved in methanol (1 ml) and sodium borohydride (8.8 mg, 0.23 mmol) was added. The mixture was stirred for 30 minutes. Ammonium acetate (18 mg) was added and the solvent was removed under reduced pressure. The residue was purified with preparative HPLC on a C8 column. A gradient from 20% to 50 % MeCN in 0.1 M ammonium acetate buffer was used as eluent. After lyophilisation, the title compound was obtained as a white solid (2.5 mg, 18%).

H-NMR (400 MHz, DMSO-*d*₆): 0.75 (d, 3H), 0.79-0.87 (m, 9H), 1.92-2.06 (m, 2H), 2.86-2.92 (m, 2H), 4.01 (brs, 1H), 4.25 (d, 0.5H), 4.28 (d, 0.5H), 4.32 (t, 1H), 4.57 (d, 2H), 4.67-4.76 (m, 1H), 5.03 (d, 0.5H), 5.05 (d, 0.5H), 5.69 (brs, 1H), 6.93 (d, 2H), 7.04-7.17 (m, 4H), 7.18-7.25 (m, 2H), 7.29-7.38 (m, 4H), 7.85 (d, 1H), 7.96 (brs, 1H). *M/z*: 682.1 (M-H).

Example 55

50 *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-valyl-L-methionine

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-valine (13.3 mg, 0.023 mmol), *tert*-butyl L-methioninate hydrochloride (7.4 mg, 0.031 mmol) and *N*-methylmorpholine (10 μ l, 0.091 mmol) were dissolved in 1 ml. After 5 minutes, TBTU (8.9 mg, 0.028 mmol) was added and the resulting suspension was stirred overnight. Additional *tert*-butyl L-methioninate hydrochloride (2.1 mg, 0.0087 mmol), *N*-methylmorpholine (5 μ l, 45 μ mol) and TBTU (2.1 mg, 6.54 μ mol) were added and the mixture was stirred for 2 h. The formation of the ester was confirmed. *M/z* 768.1 (M-H) and 770.0 (M+H). The yellow suspension was purified on silica gel (1g) and eluted with EtOAc:DCM (15:85). The pure fractions were concentrated and formic acid (1.5 ml) was added. The solution was stirred at 50°C overnight. The solvent was removed under reduced pressure. Toluene was added and removed under reduced pressure. The residue was dissolved in methanol (1 ml) and sodium borohydride (9.9 mg, 0.26 mmol) was added. The resulting reaction mixture was stirred for 10 minutes. Ammonium acetate (18.9 mg) was added and the solvent was removed under reduced pressure. The residue was purified with preparative HPLC on a C8 column. A gradient from 20% to 40 % MeCN in 0.1 M ammonium acetate buffer was used as eluent. After lyophilisation, the title compound was obtained as a white solid (4.6 mg, 28%). ¹H-NMR (400 MHz, DMSO-*d*₆): 0.75 (d, 3H), 0.79 (d, 3H), 1.79-1.97 (m, 3H), 1.99 (s, 3H), 2.36-2.44 (m, 2H), 2.86-2.92 (m, 2H), 2.24-4.35 (m, 3H), 4.58 (d, 2H), 4.67-4.76 (m, 1H), 5.03 (d, 0.5H), 5.5 (d, 0.5H), 5.63 (t, 1H), 6.95 (d, 2H), 7.05-7.16 (m, 4H), 7.18-7.24 (m, 2H), 7.30-7.38 (m, 2H), 7.82 (d, 1H), 7.37 (d, 1H). *M/z*: 714.0 (M-1) and 716.1 (M+H).

Example 56

N-[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[(2-hydroxy-2-phenylethyl)thio]-4-oxoazetidin-2-yl)phenoxy]acetyl]glycyl-D-valine

(4-((2*R*,3*R*)-1-(4-Fluorophenyl)-4-oxo-3-[(2-oxo-2-phenylethyl)thio]azetidin-2-yl)phenoxy)acetic acid (15 mg, 0.043 mmol), *tert*-butyl glycyl-D-valinate hydrochloride, (14.3 mg, 0.054 mmol) and *N*-methylmorpholine (14 μ l, 0.13 mmol) were dissolved in DCM (2 ml). After 5 minutes, TBTU (16.9 mg, 0.053 mmol) was added and the reaction mixture was stirred for 2.5 h. The formation of the ester was confirmed. *M/z*: 678.35 (M+H). The solvent was removed under reduced pressure. The residue was dissolved in EtOAc:DCM (1:3) and purified on silica gel (1 g) using EtOAc:DCM (1:3) as eluent. The fractions were collected and concentrated. The residue (0.029 g) was dissolved in DCM (3 ml) and TFA (0.5 ml) was added. The reaction mixture was stirred overnight. The solvent was removed under reduced pressure and toluene was added and removed under reduced pressure. The yellowish residue was dissolved in MeOH (2 ml) and sodium borohydride (16.2 mg, 0.43 mmol) was added. The reaction mixture was stirred for 10 minutes. Ammonium acetate (31.4 mg) was added and the solvent was removed under reduced pressure. The residue was purified with preparative HPLC on a C8 column. A gradient from 20% to 50 % MeCN in 0.1 M ammonium acetate buffer was used as eluent. After lyophilisation, the title compound was obtained as a white solid (7.6 mg, 28 %). H-NMR (400 MHz, DMSO-*d*₆): 0.79 (d, 3H), 0.81 (d, 3H) 1.95-2.05 (m, 1H), 2.84-2.96 (m, 2H), 3.79 (d, 2H), 3.98-4.04 (m, 1H), 4.27 (d, 0.5H), 4.30 (d,

0.5H), 4.51 (s, 2H), 4.66-4.75 (m, 1H), 5.02 (d, 0.5H), 5.04 (d, 0.5H), 6.98 (d, 2H), 7.10-7.17 (m, 2H), 7.19-7.32 (m, 7H), 7.36 (d, 2H), 7.77 (t, 1H), 8.26 (t, 1H). M/z: 622.1 (M-H) and 624.2 (M+H).

5

Example 57

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-hydroxy-2-(4-methylphenyl)ethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-valine

10

[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-methylphenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetic acid (15.0 mg, 0.043 mmol), *tert*-butyl glycyl-D-valinate hydrochloride (10.8 mg, 0.04 mmol), *N*-methylmorpholine (10 μ l, 0.09 mmol) were dissolved in DCM (2 ml). After 5 minutes, TBTU (12.1 mg, 0.04 mmol) was added and the reaction mixture was stirred overnight. The formation of the ester was confirmed. M/z: 690.13 (M-H) and 692.15 (M+H). The reaction mixture was purified on silica gel (1 g) and eluted with EtOAc:DCM (1:4). The collected fractions were concentrated. The oily residue was dissolved in DCM (1.5 ml) and TFA (1 ml) was added. The reaction mixture was stirred for 2.5 h. The solvent was evaporated. Toluene was added and evaporated to assist the removal of TFA. The residue was dissolved in methanol (1.5 ml) and sodium borohydride (12.2 mg, 0.32 mmol) was added. Additional sodium borohydride (4.2 mg, 0.11 mmol) was added and the mixture was stirred for 15 minutes. The solvent was removed under reduced pressure and the residue was purified on preparative HPLC on a C8 column. A gradient from 20% to 50 % MeCN in 0.1M ammonium acetate buffer was used as eluent. After lyophilisation, the title compound was obtained as a white solid (10.4 mg, 52 %). H-NMR (400 MHz, DMSO-*d*₆): 0.80 (d, 6H), 1.95-2.05 (m, 1H), 2.24 (brs, 3H), 2.80-2.94 (m, 2H), 3.78 (d, 2H), 4.00 (brs, 1H), 4.23 (d, 0.5H), 4.27 (brs, 0.5H) 4.51 (s, 2H), 4.61-4.70 (m, 1H), 5.00 (m, 1H), 6.97 (d, 2H) 7.06 (d, 2H), 7.10-7.18 (m, 4H), 7.19-7.25 (m, 2H), 7.34 (d, 2H), 7.76 (brs, 1H), 8.26 (t, 1H). M/z: 636.1 (M-H) and 638.1 (M+H).

Example 58

35

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-valyl-D-tyrosine

40 *N*-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-valine (14.7 mg, 0.025 mmol), added *tert*-butyl D-tyrosinate hydrochloride (10.5 mg, 0.038 mmol) and *N*-methylmorpholine (10 μ l, 91 μ mol) were dissolved in DCM (1.5 ml). After 5 minutes, TBTU (9.9 mg, 0.031 mmol) was added and the reaction mixture was stirred overnight. Additional *tert*-butyl D-tyrosinate hydrochloride (3.6 mg, 0.013 mmol), *N*-methylmorpholine (10 μ l, 91 μ mol) and TBTU (3.1 mg, 9.7 μ mol) were added. The mixture was stirred for 3 h. The formation of the ester was confirmed. M/z: 800.07 (M-H) and 802.08 (M+H). Aqueous KHSO₄ (3 ml) was added and the mixture (pH of 2) was extracted with DCM (3x5 ml). The combined organic phases were washed with water (2x10 ml), dried over Na₂SO₄, filtered and concentrated. The oily residue (22.4 mg) was dissolved in DCM (1.5 ml) and TFA (1.0 ml) was added. The mixture was stirred overnight. The

solvent was evaporated. Toluene was added and removed under reduced pressure. The residue was dissolved in methanol (2 ml) and sodium borohydride (14 mg) was added. The solution was stirred for 10 minutes. Ammonium acetate (15 mg) was added and the solvent was removed under reduced pressure. The residue was purified with preparative HPLC on a C8 column. A gradient from 20% to 50 % MeCN in 0.1M ammonium acetate buffer was used as eluent. After lyophilisation, the title compound was obtained as a white solid (8.4 mg, 45 %).
1H-NMR (400 MHz, DMSO-*d*₆): 0.70 (d, 3H), 0.76 (d, 3H), 1.90-1.99 (m, 1H), 2.73-2.80 (m, 1H), 2.86-2.95 (m, 3H), 3.96-4.04 (m, 1H), 4.06-4.12 (m, 1H), 4.27 (d, 0.5H), 4.29 (d, 0.5H), 4.50-4.61 (m, 2H), 4.67-4.76 (m, 1H), 5.02 (d, 0.5H), 5.04 (d, 0.5H), 6.53 (d, 2H), 6.91 (q, 4H), 7.04-7.15 (m, 4H), 7.18-7.25 (m, 2H), 7.30-7.38 (m, 4H), 7.58-7.65 (brs, 1H), 7.91 (d, 1H). M/z: 746.0 (M-H).

Example 59

15 *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-valyl-D-lysine

20 *N*-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-valine (14.7, 0.025 mmol), *tert*-butyl *N*⁶-(*tert*-butoxycarbonyl)-D-lysinate hydrochloride (10.3, 0.03 mmol) and *N*-methylmorpholine (10 μ l, 91 μ mol) were dissolved in DCM (1.5 ml). After 5 min, TBTU (9.8 mg, 0.03 mmol) was added and the
25 reaction mixture was stirred overnight. Additional *N*⁶-(*tert*-butoxycarbonyl)-D-lysinate hydrochloride (3.4 mg, 0.01 mmol), *N*-methylmorpholine (5 μ l, 45 μ mol) and TBTU (3.3 mg, 0.01 mmol) were added and the mixture was stirred for 2 h. Aqueous KHSO₄ (3 ml) was added and the mixture (pH of 3) was extracted with DCM (3x5 ml). The combined organic phases were washed with water (2x5 ml) and dried over Na₂SO₄. The solvent was removed
30 under reduced pressure. The crude residue (17.1 mg) was dissolved in DCM (1.5 ml) and TFA (1ml). The solution was stirred for 1.5 h. The solvent was removed under reduced pressure. Toluene was added and evaporated to assist the removal of TFA. The residue was dissolved in methanol (2 ml) and sodium borohydride (11.5 mg, 0.30 mmol) was added. The mixture was for ca 15 minutes. After removal of the solvent under reduced pressure, the
35 residue was purified with preparative HPLC on a C8 column. A gradient from 20% to 50 % MeCN in 0.1M ammonium acetate buffer was used as eluent. After lyophilisation, the title compound was obtained as a white solid. (7.8 mg, 43 %). 1H-NMR (400 MHz, DMSO-*d*₆): 0.75 (d, 3H), 0.79 (d, 3H), 1.17-1.63 (m, 4H), 1.96-2.06 (m, 1H), 2.61-2.69 (m, 2H), 2.85-2.93 (m, 2H), 3.72-3.80 (m, 1H), 4.12 (t, 1H), 4.27 (s, 0.5H), 4.30 (s, 0.5H), 4.53-5.64 (m, 2H),
40 4.67-4.76 (m, 1H), 5.01-5.05 (m, 1H), 6.94 (d, 2H), 7.03-7.16 (m, 4H), 7.29-7.39 (m, 4H), 7.50-7.57 (brs, 1H), 8.05 (d, 1H). M/z: 713.1.

Example 60

45 *N*-{[4-((2*R*,3*R*)-3-[(2-hydroxy-2-phenylethyl)thio]-4-oxo-1-phenylazetidin-2-yl)phenoxy]acetyl}glycyl-D-valine

50

[4-((2*R*,3*R*)-3-[[5,5-Dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl]thio}-4-oxo-1-phenylazetidin-2-yl)phenoxy]acetic acid (12.6 mg, 0.024 mmol) and *N*-methylmorpholine (15 μ l, 0.14 mmol) were dissolved in DCM (2 ml). Additional DCM (2 ml), *N*-methylmorpholine (20 μ l, 0.18 mmol) and *tert*-butyl glycyl-D-valinate hydrochloride, (9.0 mg, 0.034 mmol) were added after 0.5 h and the mixture was stirred for 10 minutes. TBTU (10.5 mg, 0.033 mmol) was added and the mixture was stirred overnight. The formation of the ester was confirmed. *M/z*: 746.1. The solvent was removed under reduced pressure and the residue was purified with preparative HPLC on a C8 column. A gradient from 20% to 50% MeCN in 0.1 M ammonium acetate buffer was used as eluent. After lyophilisation, the obtained compound was dissolved in DCM (2 ml) and TFA (1 ml) was added. The reaction mixture was stirred for 2.5 h. The hydrolysis of the ester was confirmed. *M/z*: 604.2. The solvent was removed under reduced pressure. Co-evaporation with toluene was performed to assist the removal of TFA. The residue was dissolved in methanol (2 ml) and sodium borohydride (9.2 mg, 0.24 mmol) was added. After 15 minutes, the solvent was evaporated off and the residue was purified with preparative HPLC on a C8 column. A gradient from 20% to 50 % MeCN in 0.1 M ammonium acetate buffer was used as eluent. After lyophilisation, the title compound was obtained as a white solid was (4.0 mg, 28%). H-NMR (400 MHz, DMSO-*d*₆): 0.78 (d, 6H), 1.95-2.04 (m, 1H), 2.83-2.97 (m, 2H), 3.76 (d, 2H), 3.89-3.95 (m, 1H), 4.26 (d, 0.5H), 4.30 (d, 0.5H), 4.51 (s, 2H), 4.67-4.75 (m, 1H), 5.01 (d, 0.5H), 5.03 (d, 0.5H), 6.98 (d, 2H), 7.03 (t, 1H), 7.17-7.22 (m, 3H), 7.23-7.32 (m, 6H), 7.36 (d, 2H), 7.55-7.65 (m, 1H), 8.29 (t, 1H). *M/z*: 603.96 (M-H).

Example 61

N-[[4-((2*R*,3*R*)-1-(4-chlorophenyl)-3-[[2-(4-chlorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl]glycyl-D-valine

[4-((2*R*,3*R*)-1-(4-Chlorophenyl)-3-[[2-(4-chlorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetic acid (15.3 mg, 0.03 mmol), *N*-methylmorpholine (10 μ l, 0.091 mmol) and *tert*-butyl glycyl-D-valinate hydrochloride, (10.4 mg, 0.039 mmol) were dissolved in DCM (2 ml). After 10 minutes, TBTU (11.9 mg, 0.037 mmol) was added and the mixture was stirred overnight. The intermediate *tert*-butylester was confirmed. *M/z*: 727.8 (M-H). The reaction mixture was extracted between water (10 ml, acidified to pH of 3 with KHSO₄ (2M)) and DCM (3x10 ml). The combined organic phases were washed with water (2x20 ml), dried over Na₂SO₄, filtered and concentrated. The oily residue was dissolved in DCM (2 ml) and TFA (1.3 ml) was added. The mixture was stirred overnight. The solvent was evaporated and co-evaporation with toluene was performed to assist the removal of TFA. The residue was dissolved in methanol (2 ml) and sodium borohydride (12.3 mg, 0.33 mmol) was added. After 15 minutes, ammonium acetate (17mg) was added and the solvent was removed under reduced pressure. The residue was purified on preparative HPLC on C8 column. A gradient from 20% to 50 % MeCN in 0.1M ammonium acetate buffer was used as eluent. After lyophilisation, the title compound was obtained as a white solid (15.1 mg, 77 %). H-NMR (400 MHz, DMSO-*d*₆): 0.78 (d, 3H), 0.80 (d, 3H), 0.95-1.03 (m, 1H), 2.83-3.00 (m, 2H), 3.75-3.80 (m, 2H), 3.93-4.00 (t, 1H), 4.30 (d, 0.5H), 4.36-4.38 (brs, 0.5H), 4.52 (s, 2H), 4.69-4.77 (m, 1H), 5.02 (d, 0.5H), 5.06 (d, 0.5H), 6.96-7.00 (m, 2H), 7.18-7.22 (m, 2H), 7.31-7.38 (m, 8H), 7.60-7.74 (m, 1H), 8.28 (t, 1H). *M/z*: 671.9 (M+H).

Example 62

***N*-{[4-((2*R*,3*R*)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-lysine**

5

N-{[4-((2*R*,3*R*)-1-(4-Chlorophenyl)-3-{[2-(4-chlorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (14.6 mg, 0.026 mmol) and *N*-methylmorpholine (20 μ l, 0.18 mmol) were dissolved in DCM (2 ml). *tert*-Butyl *N*⁶-(*tert*-butoxycarbonyl)-D-lysinate hydrochloride (11.1 mg, 0.033 mmol) was added and after 5 minutes TBTU (9.8 mg, 0.031 mmol) was added to the suspension. The mixture was stirred overnight. Additional *tert*-butyl *N*⁶-(*tert*-butoxycarbonyl)-D-lysinate (4.8 mg, 0.014 mmol), *N*-methylmorpholine (10 μ l, 91 μ mol) and TBTU (4.6 mg, 0.014 mmol) were added and the mixture was stirred for 2.5 h. The formation of the ester was confirmed. *M/z*: 855.4 (M-H). DCM (3 ml) and water (5 ml) were added and the solution was acidified to pH 3 with KHSO₄ (2M). The organic phase was washed with water (2x5 ml). The combined water phases were extracted with DCM (2x5 ml). The organic phase was dried over Na₂SO₄, filtered and concentrated under reduced pressure. The oily residue was dissolved in DCM (1.5 ml) and TFA (1 ml) was added. The mixture was stirred for 2.5 h. The mixture was concentrated and co-evaporation with toluene was performed to assist the removal of TFA. The residue was dissolved in methanol (2 ml) and sodium borohydride (10.4 mg, 0.027 mmol) was added. After 15 minutes, ammonium acetate buffer (0.1M, 1.5 ml) was added and the solvent was removed under reduced pressure. The residue was purified with preparative HPLC on a C8 column. A gradient from 20% to 50 % MeCN in 0.1M ammonium acetate buffer was used as eluent. After lyophilisation, the title compound was obtained as a white solid (10.8 mg, 59 %). H-NMR (400 MHz, DMSO-*d*₆): 1.18-1.36 (m, 2H), 1.41-1.70 (m, 4H), 2.71 (t, 2H), 2.84-2.97 (m, 2H), 3.72-3.75 (brd, 2H), 3.93 (m, 1H), 4.30 (d, 0.5H), 4.34 (d, 0.5H), 4.52 (s, 2H), 4.68-4.77 (m, 1H), 5.03 (d, 0.5H), 5.07 (d, 0.5H), 6.98 (d, 2H), 7.20 (d, 2H), 7.31-7.38 (m, 8H), 7.63-7.72 (m, 1H), 8.34 (t, 1H). *M/z*: 705.1.

30

Example 63

***(2R)*-2-[(*N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl)amino]-4-phenylbutanoic acid**

35

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (14.4 mg, 0.027 mmol) and *N*-methylmorpholine (15 μ l, 0.14 mmol) were dissolved in DMF (3 ml). After 5 minutes, TBTU (10.3 mg, 0.032 mmol) was added and the mixture was stirred at 30°C for 20 minutes. *(2R)*-2-Amino-4-phenylbutanoic acid (5.7 mg, 0.032 mmol) was added and the mixture was stirred at ambient temperature for 1.5 hours. The formation of the intermediate acid was confirmed. *M/z*: 702.0. MeOH (2.5 ml) and sodium borohydride were added and the mixture was stirred for 20 minutes. Ammonium acetate (31 mg) was added. The mixture was concentrated and purified with preparative HPLC on a C8 column. A gradient from 20% to 45% MeCN in 0.1M ammonium acetate buffer was used as eluent. After lyophilisation, the title compound was obtained as a white solid (9.4 mg, 50 %). H-NMR (400 MHz, DMSO-*d*₆): 1.74-1.87 (m, 1H), 1.89-1.99 (m, 1H),

50

2.49-2.53 (m, 2H), 2.82-2.98 (m, 2H), 3.77 (d, 2H), 3.94-4.01 (m, 1H), 4.27 (d, 0.5H), 4.31 (d, 0.5H), 4.53 (s, 2H), 4.68-4.77 (m, 1H), 5.02 (d, 0.5H), 5.05 (d, 0.5H), 6.99 (d, 2H), 7.05-7.16 (m, 7H), 7.19-7.26 (m, 4H), 7.30-7.38 (m, 4H), 7.78-7.88 (dd, 1H), 8.35 (t, 1H). M/z: 702.0 (M-H).

5

Example 64

10 (2R)-2-[(N-[[4-((2R,3R)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl]glycyl)amino]-4-(4-hydroxyphenyl)butanoic acid

15 N-[[4-((2R,3R)-1-(4-Fluorophenyl)-3-[[2-(4-fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl]glycine (15.2 mg, 0.028 mmol) and N-methylmorpholine (15 μ l, 0.14 mmol) were dissolved in DMF (2 ml). TBTU (10.5 mg, 0.033 mmol) was added and after 20 minutes (2R)-2-amino-4-(4-hydroxyphenyl)butanoic acid hydrobromide (9.2 mg, 0.033 mmol) was added. The reaction mixture was stirred for 1.5 h. The formation of the intermediate acid was confirmed. M/z: 718.3. MeOH (2 ml) and sodium borohydride (10.7 mg, 0.28 mmol) were added and the mixture was stirred for 20 minutes. Ammonium acetate 20 (34 mg) was added and the methanol was removed under reduced pressure. The residue was purified with preparative HPLC on a C8 column. A gradient from 20% to 45% MeCN in 0.1M ammonium acetate buffer was used as eluent. After lyophilisation, the title compound was obtained as a white solid (9.6 mg, 47%). H-NMR (400 MHz, DMSO-*d*₆): 1.75-1.86 (m, 1H), 1.89-1.98 (m, 1H), 2.41 (t, 2H), 2.81-2.98 (m, 2H), 3.78 (d, 2H), 3.97-4.05 (m, 1H), 4.27 25 (d, 0.5H), 4.31 (d, 0.5H), 4.53 (s, 2H), 4.67-4.76 (m, 1H), 5.02 (d, 0.5H), 5.05 (d, 0.5H), 6.62 (d, 2H), 6.91 (d, 2H), 6.98 (d, 2H), 7.05-7.16 (m, 4H), 7.20-7.25 (m, 2H), 7.30-7.39 (m, 4H), 7.87-7.97 (m, 1H), 8.30 (t, 1H), 8.91-9.30 (br, 1H). M/z: 718.0 (M-H).

30 Example 65

35 N-[[4-((2R,3R)-1-(4-fluorophenyl)-3-[[2-hydroxy-2-(4-methoxyphenyl)ethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl]glycyl-D-alanine

40 N-[[4-((2R,3R)-1-(4-Fluorophenyl)-3-[[2-(4-methoxyphenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl]glycine (0.04 g, 0.072 mmol) and N-methylmorpholine (0.022 g, 0.217 mmol) were dissolved in CH₂Cl₂ (4 ml). *tert*-Butyl D-alaninate hydrochloride (0.016 g, 0.087 mmol) and TBTU (0.030 g, 0.094 mmol) were added. After 2 h, the reaction mixture was purified on silica gel and eluted with EtOAc/ CH₂Cl₂ (25/75). Pure fractions were collected and concentrated. The residue was dissolved in CH₂Cl₂ (3 ml) and TFA (2 ml). The hydrolysis was completed after 2h. The reaction mixture was concentrated and MeOH (3 ml) and NaBH₄ (0.011 g, 0.290 mmol) were added. The mixture was stirred for 5 minutes. 45 The reaction was quenched by the addition of 0.1M NH₄OAc buffer (2 ml) and the solvent evaporated. The residue was purified by preparative HPLC using an eluent of 0-50% CH₃CN in 0.1M NH₄OAc buffer. Freeze-drying of the pure fractions afforded the title compound (0.030 g, 66%) as a colourless solid. M/z: 624.2, (M -1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 1.16 (d, 3H), 2.83-2.93 (m, 2H), 3.68-3.74 (m, 5H), 3.88-3.95 (m, 1H), 4.23-4.26 (m, 1H),

4.51 (s, 2H), 4.60-4.70 (m, 1H), 5.00-5.03 (m, 1H), 6.81-7.37 (m, 12H), 7.74-7.79 (m, 1H), 8.29-8.34 (m, 1H).

5 Example 66

1-[(N-[[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl)amino]cyclopropanecarboxylic acid

- 10 N-Methyl morpholine (0.037 g, 0.370 mmol) and TBTU (0.039 g, 0.120 mmol) were added to a solution of *N*-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.025 g, 0.046 mmol) in DMF (2 ml) at 30 °C. After 1h, 1-aminocyclopropane carboxylic acid (0.019 g, 0.185 mmol) was added. After 1h,
- 15 the reaction was quenched by the addition of water (1 ml). After 10 minutes, MeOH (2 ml) and NaBH₄ (0.035 g, 0.925 mmol) were added. Full conversion to the corresponding alcohol was obtained after 5 minutes. The reaction was quenched by the addition of 0.1M NH₄OAc buffer (2 ml). The product was purified by preparative HPLC (eluent 0-50% CH₃CN in 0.1M NH₄OAc buffer). Freeze-drying of the pure fractions afforded the title compound (0.045 g,
- 20 78%) as a colourless solid. ¹H NMR [(CD₃)₂SO], 400 MHz] δ 0.78-0.88 (m, 2H), 1.08-1.22 (m, 2H), 2.84-2.94 (m, 2H), 3.63-3.72 (m, 2H), 4.24-4.29 (m, 1H), 4.48-4.52 (m, 2H), 4.68-4.75 (m, 1H), 5.03-5.06 (m, 1H), 6.96-7.37 (m, 12H), 7.78-8.36 (m, 2H).

25 Example 67

***N*-[[4-((2R,3R)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-methyl-D-valine**

- 30 TBTU (0.020 g, 0.063 mmol) was added to a solution of *N*-{[4-((2R,3R)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.030 g, 0.052 mmol) and *N*-methylmorpholine (0.016 g, 0.157 mmol) in CH₂Cl₂ (5 ml) at 30 °C. After 30 minutes, D-tert-leucine (0.008 g, 0.063 mmol) was added and the mixture was stirred
- 35 for 30 minutes. The reaction mixture was concentrated. Toluene (2 ml) was added and evaporated. MeOH (3 ml) and sodium borohydride (0.020 g, 0.523 mmol) were added. Full conversion to the corresponding alcohol was obtained after 5 minutes. The reaction was quenched by the addition of 0.1M NH₄OAc (1 ml) buffer and the mixture was concentrated. The residue was purified by preparative HPLC using an eluent of 0-50% CH₃CN in 0.1M
- 40 NH₄OAc buffer. Freeze-drying of the pure fractions afforded the title compound (0.021 g, 58%) as a colourless solid. ¹H NMR [(CD₃)₂SO], 400 MHz] δ 0.85 (s, 9H), 2.82-2.98 (m, 2H), 3.75-3.81 (m, 2H), 3.91-3.96 (m, 1H), 4.29-4.37 (m, 1H), 4.52 (s, 2H), 4.70-4.78 (m, 1H), 5.01-5.06 (m, 1H), 6.97-6.99 (m, 2H), 7.19-7.21 (m, 2H), 7.32-7.36 (m, 8H), 7.52-7.63 (m, 1H), 8.27-8.32 (m, 1H).

Example 68

***N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-tryptophan**

- 5 TBTU (0.016 g, 0.051 mmol) was added to a solution of *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.025 g, 0.046 mmol) and *N*-methymorpholine (0.014 g, 0.139 mmol) in DMF (2 ml) at 30 °C. After 1h, DMSO (1 ml) and D-tryptophan (0.019 g, 0.092 mmol) were added. After 10 minutes, the reaction was quenched by the addition of water (1ml). The mixture was stirred for 10 minutes
- 10 and MeOH (1 ml) and NaBH₄ (0.035 g, 0.925 mmol) were added. After 5 minutes, full conversion to the corresponding alcohol was obtained. The reaction was quenched by the addition of 0.1M NH₄OAc buffer (1 ml). The reaction mixture was concentrated and the residue was purified by preparative HPLC using an eluent of 0-50% CH₃CN in 0.1M NH₄OAc buffer. This gave the title compound (0.028 g, 83%) as a colourless solid. M/z: 727.0
- 15 (M-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 2.79-3.18 (m, 4H), 3.61-3.80 (m, 2H), 4.26-4.34 (m, 2H), 4.43-4.54 (m, 2H), 4.68-4.78 (m, 1H), 4.97-5.04 (m, 1H), 6.84-7.55 (m, 17H), 7.65-7.82 (m, 1H), 8.22-8.25 (m, 1H), 10.73 (s, 1H).

20 **Example 69**

***N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-histidine**

- 25 TBTU (0.016 g, 0.051 mmol) was added to a solution of *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.025 g, 0.046 mmol) and *N*-methymorpholine (0.014 g, 0.139 mmol) in DMF (2 ml) at 30 °C. After 1 h, D-histidine (0.014 g, 0.092 mmol) and tetrabutylammoniumbromide (0.003 g, 0.009
- 30 mmol) were added. The reaction mixture was stirred overnight (30% conversion) and the reaction was quenched by the addition of water (2 ml). Purification by preparative HPLC using an eluent of 0-50% CH₃CN in 0.1M NH₄OAc buffer afforded the intermediate ketone, which was reduced by the addition of MeOH (3 ml) and NaBH₄ (0.005 g, 0.139 mmol). Full conversion to the corresponding alcohol was obtained after 5 minutes. The reaction was
- 35 quenched by the addition of 0.1M NH₄OAc buffer (2 ml) and the mixture was concentrated. Purification by preparative HPLC using an eluent of 0-40% CH₃CN in 0.1M NH₄OAc buffer afforded the title compound (0.001 g, 4.5%) as a colourless solid. M/z: 680.0. ¹H NMR [(CD₃)₂SO], 400 MHz] δ 2.82-2.93 (m, 4H), 3.71-3.80 (m, 2H), 4.11-4.30 (m, 2H), 4.52 (s, 2H), 4.68-4.73 (m, 1H), 5.04-5.07 (m, 1H), 6.68-7.50 (m, 14H), 7.90-7.96 (m, 1H), 8.27-8.33
- 40 (m, 1H).

Example 70

- 45 ***N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-(2-naphthyl)-D-alanine**

- TBTU (0.019 g, 0.060 mmol) was added to a solution of *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.025
- 50

g, 0.046 mmol) and N-methylmorpholine (0.014 g, 0.139 mmol) in CH₂Cl₂ (5 ml) at 30 °C. After 1h, DMF (3 ml), DMSO (1 ml) and D-(2-naphthyl)alanine (0.011 g, 0.051 mmol) were added. The reaction was quenched by the addition of water (1 ml) after 1 h. NaBH₄ (0.035 g, 0.925 mmol) was added. After 5 minutes, full conversion to the corresponding alcohol was obtained. The reaction was quenched by the addition of 0.1M NH₄OAc buffer (1 ml) and the mixture was concentrated. Purification by preparative HPLC using an eluent of 0-55% CH₃CN in 0.1M NH₄OAc buffer afforded the title compound (0.017 g, 48%) as a colourless solid. M/z: 738.0 (M-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 2.75-3.27 (m, 4H), 3.55-3.83 (m, 2H), 4.25-4.55 (m, 4H), 4.68-4.79 (m, 1H), 4.92-5.02 (m, 1H), 6.72-7.80 (m, 20H), 8.26-8.30 (m, 1H).

Example 71

15 *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-methyl-D-valine

TBTU (0.021 g, 0.067 mmol) was added to a solution of *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.030 g, 0.056 mmol) and N-methyl morpholine (0.017 g, 0.166 mmol) in CH₂Cl₂ (5 ml) at 30 °C. After 1.5h, D-tert-leucine (0.011 g, 0.083 mmol) was added. Full conversion to the corresponding amide was obtained after 30 minutes. The reaction was quenched by the addition of water (1 ml). After 10 minutes, MeOH (3 ml) and NaBH₄ (0.042 g, 1.11 mmol) were added. After 5 minutes, the reaction was quenched by the addition of 0.1M NH₄OAc buffer (1 ml). The reaction mixture was concentrated and purified by preparative HPLC using an eluent of 0-50% CH₃CN in 0.1M NH₄OAc buffer. The title compound (0.025 g, 69%) was obtained as a colourless solid. M/z: 654.0 (M-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 0.86 (s, 9H), 2.82-2.98 (m, 2H), 3.76-3.81 (m, 2H), 3.92-3.96 (m, 1H), 4.26-4.33 (m, 1H), 4.52 (s, 2H), 4.68-4.76 (m, 1H), 5.02-5.07 (m, 1H), 6.97-7.37 (m, 12H), 7.58-7.63 (m, 1H), 8.29-8.34 (m, 1H).

Example 72

35 *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-(3*R*,4*S*,5*R*)-3,4,5,6-tetrahydroxy-D-norleucine

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.025 g, 0.046 mmol) was dissolved in DMSO (2 ml) at 30 °C. N-Methylmorpholine (0.014 g, 0.139 mmol) and TBTU (0.018 g, 0.056 mmol) were added. After 1h, D-glucosaminic acid (0.018 g, 0.092 mmol) and tetrabutylammoniumbromide (0.001 g, 0.005 mmol) were added. The mixture was stirred for 15 minutes. Additional TBTU (18 mg, 0.056 mmol) was added. After 30 minutes, approximately 30% amide formation had occurred. The reaction was quenched by the addition of 0.1M NH₄OAc buffer (2 ml). The intermediate ketone was purified by preparative HPLC using an eluent of 0-50% CH₃CN in 0.1M NH₄OAc buffer and freeze-dried. MeOH (3 ml) and NaBH₄ (0.005 g, 0.139 mmol) were added. After 5 minutes, the reaction was quenched by the addition of 0.1M NH₄OAc buffer (1 ml). Concentration of the mixture and purification by preparative HPLC using an eluent of 0-50% CH₃CN in 0.1M NH₄OAc buffer gave the title compound (0.005 g, 16%) as a colourless

solid. M/z: 718.0 (M-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 2.82-2.94 (m, 2H), 3.34-3.56 (m, 4H), 3.76-3.80 (m, 2H), 3.87-3.90 (m, 1H), 4.07-4.11 (m, 1H), 4.27-4.32 (m, 1H), 4.52 (s, 2H), 4.68-4.76 (m, 1H), 5.02-5.05 (m, 1H), 6.97-7.39 (m, 12H), 7.63-7.70 (m, 1H), 8.28-8.35 (m, 1H).

5

Example 73

10 *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-β-phenyl-D-phenylalanine

TBTU (0.018 g, 0.056 mmol) was added to a solution of *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.025 g, 0.046 mmol) and *N*-methylmorpholine (0.023 g, 0.231 mmol) in CH₂Cl₂ (5 ml) at 30 °C. After 1.5h, β-phenyl-D-phenylalanine trifluoro acetic acid salt (0.033 g, 0.092 mmol) was added. The mixture was stirred for 5 minutes. Water (1 ml) was added and the mixture was concentrated. MeOH (3 ml) and NaBH₄ (0.017 g, 0.462 mmol) were added. After 5 minutes, the reaction was quenched by the addition of 0.1M NH₄OAc buffer (1 ml) followed by concentration of the mixture. Purification by preparative HPLC using an eluent of 0-45% CH₃CN in 0.1M NH₄OAc buffer and lyophilisation gave the title compound (0.021 g, 59%) as a colourless solid. M/z: 764.1 (M-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 2.82-2.95 (m, 2H), 3.42-3.49 (m, 1H), 3.66-3.74 (m, 1H), 4.25-4.33 (m, 2H), 4.42 (d, 1H), 4.47 (d, 1H), 4.69-4.76 (m, 1H), 5.03-5.12 (m, 2H), 6.94-7.38 (m, 22H), 8.10-8.14 (m, 1H), 8.18-8.24 (m, 1H).

25

Example 74

30 (2*R*)-4-cyclohexyl-2-[(*N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl)amino]butanoic acid

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-[[2-(4-fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.025 g, 0.046 mmol) was dissolved in DMF (2 ml) at 30 °C. *N*-Methyl morpholine (0.034 g, 0.333 mmol) and TBTU (0.043 g, 0.133 mmol) were added. After 1h, sodium (2*R*)-2-amino-4-cyclohexylbutanoate (0.039 g, 0.189 mmol), DMSO (2 ml) and tetrabutylammoniumbromide (0.004 g, 0.011 mmol) were added. The mixture was stirred for 1 h and water (1 ml) was added. After 1h, MeOH (2 ml) and NaBH₄ (0.084 g, 2.220 mmol) were added. Full conversion to the corresponding alcohol was obtained after 5 minutes. The reaction was quenched by the addition of 0.1M NH₄OAc buffer (2 ml). The mixture was purified by preparative HPLC (eluent 0-50% CH₃CN in 0.1M NH₄OAc buffer). Freeze-drying of the pure fractions gave the title compound (0.034 g, 43%) as a colourless solid. M/z: 708.1 (M-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 0.75-1.62 (m, 15H), 2.82-2.99 (m, 2H), 3.68-3.78 (m, 2H), 3.90-3.96 (m, 1H), 4.23-4.35 (m, 1H), 4.50 (s, 2H), 4.69-4.75 (m, 1H), 5.01-5.07 (m, 1H), 6.95-7.39 (m, 12H), 7.68-7.80 (m, 1H), 8.20-8.34 (m, 1H).

Example 75

(2R)-cyclopentyl[(N-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl)amino]acetic acid

- 5 *N*-{[4-((2R,3R)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.035 g, 0.065 mmol) was dissolved in DMF (2 ml) at 30 °C. *N*-Methyl morpholine (0.026 g, 0.259 mmol) and TBTU (0.027 g, 0.084 mmol) were added. After 1h, (2R)-amino(cyclopentyl)acetic acid (0.014 g, 0.097 mmol) was added. The mixture was stirred for 1 h and water (1 ml) was added. After 10 minutes, MeOH (2 ml) and NaBH₄ (0.037 g, 0.971 mmol) were added. Full conversion to the corresponding alcohol was obtained after 5 minutes. The reaction was quenched by the addition of 0.1M NH₄OAc buffer (2 ml). The mixture was purified by preparative HPLC (eluent 0-50% CH₃CN in 0.1M NH₄OAc buffer). Freeze-drying of the pure fractions gave the title compound (0.018 g, 42%) as a colourless solid. *M/z*: 666.0 (M-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 1.19-1.62 (m, 8H), 2.09-2.19 (m, 1H), 2.83-2.95 (m, 2H), 3.78 (d, 2H), 4.06-4.10 (m, 1H), 4.25-4.30 (m, 1H), 4.51 (s, 2H), 4.68-4.75 (m, 1H), 5.03-5.06 (m, 1H), 6.97-7.37 (m, 12H), 7.95-8.00 (m, 1H), 8.22 (t, 1H).

20 **Example 76**

***N*-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-methyl-D-isovaline**

- 25 *N*-{[4-((2R,3R)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.035 g, 0.065 mmol) was dissolved in DMF (2 ml) at 30 °C. *N*-Methyl morpholine (0.026 g, 0.259 mmol) and TBTU (0.027 g, 0.084 mmol) were added. After 1h, 3-methyl-D-isovaline (0.013 g, 0.097 mmol) was added. The mixture was stirred for 2 h and water (1 ml) was added. After 10 minutes, MeOH (2 ml) and NaBH₄ (0.037 g, 0.971 mmol) were added. Full conversion to the corresponding alcohol was obtained after 5 minutes and 0.1M NH₄OAc buffer (2 ml) was added. The mixture was purified by preparative HPLC (eluent 0-50% CH₃CN in 0.1M NH₄OAc buffer). Freeze-drying of the pure fractions afforded the title compound (0.020g, 47%) as a colourless solid. *M/z*: 654.0 (M-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 0.79 (d, 3H), 0.87 (d, 3H), 1.29 (s, 3H), 2.00-2.07 (m, 1H), 2.84-2.94 (m, 2H), 3.73 (d, 2H), 4.25-4.28 (m, 1H), 4.51 (s, 2H), 4.69-4.75 (m, 1H), 5.03-5.06 (m, 1H), 6.97-7.37 (m, 12H), 7.82 (s, 1H), 8.24 (t, 1H).

Example 77

- 40 *N*-{[4-((2R,3R)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*S*-(*tert*-butyl)-D-cysteine

- 45 *N*-{[4-((2R,3R)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (20 mg, 0.037 mmol) and *N*-methylemorpholine (20 µl, 0.18 mmol) were dissolved in DMF (3 ml). TBTU (14.6mg, 0.046mmol) was added and the mixture was stirred at 30°C for 45 minutes. *S*-(*tert*-Butyl)-D-cysteine hydrochloride (9.7 mg, 0.045 mmol) was added and the reaction mixture was stirred for 1.5 h. The formation of the ketone of the title compound was confirmed. *M/z*: 700.0. Methanol (2 ml) and sodium

borohydride (14.3 mg, 0.38 mmol) were added and the mixture was stirred for 30 minutes. Ammonium acetate buffer (0.1M, 2 ml) was added and the mixture was concentrated. The residue was purified with preparative HPLC on a C8 column. A gradient from 20% to 45 % MeCN in 0.1M NH₄OAc buffer was used as eluent. The MeCN was removed from the

collected fraction under reduced pressure. The remaining water solution was acidified to pH 1 with HCl (1M) and extracted with DCM. The organic phase was concentrated under reduced pressure and the residue was dissolved in MeCN and water. After lyophilisation, the title compound was obtained as a white solid (16.5 mg, 65%). H-NMR (400 MHz, DMSO-d₆): 1.21 (s, 9H), 2.71-2.78 (m, 1H), 2.82-2.86 (m, 3H), 2.74-2.80 (m, 2H), 2.18-2.26 (m, 1H), 4.27 (d, 0.5H), 4.30 (d, 0.5H), 4.51 (s, 2H), 2.67-2.76 (m, 1H), 5.03 (d, 0.5H), 5.05 (d, 0.5), 6.98 (d, 2H), 7.05-7.17 (m, 4H), 7.20-7.25 (m, 2H), 7.30-7.39 (m, 4H), 7.94-8.06 (b, 1H), 8.26 (t, 1H). M/z: 700.0 (M-H) and 702.1 (M+H).

15

Preparation of starting materials for the above Examples

Methods

20

Method 1

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-alanine

[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetic acid (50 mg, 0.1mmol), *tert*-butyl D-alaninate (23 mg, 0.12mmol) and 4-methylmorpholine (31 mg, 0.31 mmol) were dissolved in DCM (1.5 ml) and stirred at room temperature for 5 minutes. TBTU (40 mg, 0.12 mmol) was added and the reaction mixture was stirred for 2 h. TFA (0.7 ml) was added and the solution was stirred for 90 minutes. TFA and DCM were removed under reduced pressure and the residue was dissolved in DCM and washed with water. The organic layer was dried over sodium sulphate and the solvent was removed under reduced pressure giving 55 mg (95 %) of the title product. M/z: 553.0 (M-1).

40 Method 2

[4-((2*R*,3*R*)-3-[[5,5-dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl]thio]-4-oxo-1-phenylazetidin-2-yl)phenoxy]acetic acid

45

To a solution of ethyl [4-((2*R*,3*R*)-3-[[5,5-dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl]thio]-4-oxo-1-phenylazetidin-2-yl)phenoxy]acetate (0.86 g, 1.53 mmol) in MeOH (25 ml) was

added water (2.5 ml) and triethylamine (1.55 g). The reaction was stirred for 48 h at 50 °C. The solvent was evaporated and the residue was purified by preparative HPLC using a gradient of 20-70% CH₃CN in 0.1M NH₄OAc buffer. Freeze-drying of the pure fractions gave the title compound as a colourless solid. ¹H NMR [(CD₃)₂SO], 400 MHz] δ 0.50 (s, 3H), 1.15 (s, 3H), 2.93 (d, 1H), 2.96 (d, 1H), 3.21-3.37 (m, 4H), 4.13 (d, 1H), 4.28 (s, 2H), 4.97 (d, 1H), 6.83-7.41 (m, 14H).

Method 3

10 *N*-[4-[(2*R*,3*R*)-4-oxo-3-[(2-oxo-2-phenylethyl)thio]-1-phenylazetidin-2-yl]phenoxy]acetyl]glycine

15 Glycine tert-butyl ester (0.015 g, 0.112 mmol) and *N*-Methylmorpholine (0.028 g, 0.281 mmol) were added to a solution of [4-[(2*R*,3*R*)-3-[(5,5-dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl]thio]-4-oxo-1-phenylazetidin-2-yl]phenoxy]acetic acid (0.050 g, 0.094 mmol) in CH₂Cl₂ (5 ml). After 10 minutes, TBTU (0.039 g, 0.122 mmol) was added. The reaction was
20 stirred overnight. The resulting tert-butyl ester was purified on silica gel and eluted with EtOAc/CH₂Cl₂ (25/75). The pure fractions were collected and concentrated. CH₂Cl₂ (4 ml) and TFA (1 ml) were added. The solvent was evaporated after 2 h and the residue was purified by preparative HPLC using an eluent of 20-70% CH₃CN in 0.1M NH₄OAc buffer. Freeze-drying of the pure fractions gave the title product as a colourless solid. M/z: 503.5 (M-1). ¹H NMR (CD₃CN), 400 MHz] δ 3.84 (d, 2H), 4.18 (d, 1H), 4.24 (s, 2H), 4.50 (s, 2H), 5.04 (d, 1H), 6.99-7.12 (m, 3H), 7.25-7.66 (m, 9H), 7.94-7.96 (m, 2H).

Method 4

30 (2*R*)-([4-[(2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl]phenoxy]acetyl]amino)(phenyl)acetic acid

35 [4-[(2*R*,3*R*)-1-(4-Fluorophenyl)-3-[[2-(4-fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl]phenoxy]acetic acid (0.40 g, 0.827 mmol) was dissolved in CH₂Cl₂ (40 ml) and *tert*-butyl (2*R*)-amino(phenyl)acetate (0.206 g, 0.993 mmol) and *N*-methylmorpholine (0.251 g, 2.48 mmol) were added. After 10 minutes, TBTU (0.345 g, 1.076 mmol) was added. The reaction was stirred overnight. The resulting tert-butyl ester was concentrated and purified on silica gel
40 (eluted with EtOAc/CH₂Cl₂ 25/75). The pure fractions were collected and concentrated. CH₂Cl₂ (25 ml) and TFA (3 ml) were added. The mixture was stirred for 5 days and the solvent was removed under reduced pressure. The residue was purified by preparative HPLC using a gradient of 20-70% CH₃CN in 0.1M NH₄OAc buffer. Freeze-drying of the pure fractions gave the title compound as a colourless solid. M/z: 615.50 (M-1). ¹H NMR
45 [(CD₃)₂SO], 400 MHz] δ 4.35 (d, 1H), 4.36 (d, 1H), 4.40 (d, 1H), 4.53 (d, 1H), 4.58 (d, 1H), 4.94 (d, 1H), 5.19 (d, 1H), 6.97-7.40 (m, 15H), 8.02-8.06 (m, 2H), 8.26-8.32 (m, 1H).

Method 5

***N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-*D*-valine**

[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetic acid, (50.0 mg, 0.10 mmol) was dissolved in DCM (2 ml). *tert*-Butyl *D*-valinate hydrochloride (28.4 mg, 0.14 mmol) and *N*-methylmorpholine (3.0 μ l, 0.31 mmol) were added. After 5 minutes, TBTU (43.7 mg, 0.14 mmol) was added and the mixture was stirred overnight. The intermediate *tert*-butylester of the title compound was confirmed. *M/z*: 637.1 (M-H). The solvent was removed under reduced pressure. The yellow residue was dissolved in formic acid (1.5 ml) and heated at 50°C for 5 h. The solvent was evaporated and the residue was purified by preparative HPLC on a C8 column. A gradient from 20 to 50% MeCN in 0.1 M ammonium acetate buffer was used as eluent. After lyophilisation, the title compound was obtained as a white solid (30.5 mg, 51 %). ¹H-NMR (400 MHz, DMS-*d*₆): 0.74 (t, 6H), 1.98-2.07 (m, 1H), 3.84 (brs, 1H), 4.32 (d, 1H), 4.35 (s, 1H), 4.36 (s, 1H), 4.50 (brs, 2H), 5.16 (d, 1H), 6.96 (d, 2H), 7.10-7.17 (m, 2H), 7.19-7.24 (m, 2H), 7.31-7.38 (m, 4H), 7.66 (brs, 1H), 7.99-8.04 (m, 2H). *M/z*: 583.0 (M+H) and 581.0 (M-H).

Method 6

***N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*D*-valine**

A mixture of 3-(*R*)-4-(*R*)-1-(4-Fluorophenyl)-3-[(4-fluorobenzoyl)methylthio]-4-{4-[*N*-(carboxymethyl) carbamoylmethoxy]phenyl}azetidin-2-one (0.0229g, 0.042 mmol), (*R*)-valin *tert*-butylester hydrochloride (0.0121 g, 0.058 mmol) and *N*-methylmorpholine (0.012 ml, 0.111 mmol) in DCM (2ml) was stirred at room temperature. TBTU (0.018 g, 0.056 mmol) was added and the mixture was stirred overnight. Trifluoroacetic acid (0.65 ml) was added and after a couple of hours the hydrolysis was complete according to LC-MS. The solvent was removed under reduced pressure and the residue was purified by preparative HPLC on a Kromasil C8- column using a gradient of 5-100% MeCN in 0.15% trifluoroacetic acid buffer as eluent. The solvent was removed under reduced pressure and 0.022 g (81 %) of the title product was obtained. *M/z* 640.06.

Method 7

***N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*L*-threonine**

A mixture of 3-(*R*)-4-(*R*)-1-(4-Fluorophenyl)-3-[(4-fluorobenzoyl)methylthio]-4-{4-[*N*-(carboxymethyl) carbamoylmethoxy]phenyl}azetidin-2-one (0.0188g, 0.035 mmol), *tert*-butyl *O*-(*tert*-butyl)-*L*-threoninate (0.0151 g, 0.065 mmol) and *N*-methylmorpholine (0.012 ml, 0.111 mmol) in DCM (2ml) was stirred at room temperature. TBTU (0.018 g, 0.056 mmol) was added and the mixture was stirred overnight. Trifluoroacetic acid (0.65 ml) was added and after a couple of hours the hydrolysis was complete according to LC-MS. The solvent was removed under reduced pressure and the residue was purified by preparative HPLC on a Kromasil C8- column using a gradient of 5-100% MeCN in 0.15% trifluoroacetic acid buffer as eluent. The solvent was removed under reduced pressure and 0.014 g (63 %) of the title product was obtained. *M/z* 641.92.

Method 8

5 *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-L-asparagine

10 A mixture of 3-(*R*)-4-(*R*)-1-(4-Fluorophenyl)-3-[(4-fluorobenzoyl)methylthio]-4-{4-[*N*-(carboxymethyl) carbamoylmethoxy]phenyl}azetidin-2-one (0.0213 g, 0.039 mmol), *tert*-butyl L-asparaginate hydrochloride (0.0141 g, 0.063 mmol) and *N*-methylmorpholine (0.012 ml, 0.111 mmol) in DCM (2ml) was stirred at room temperature. TBTU (0.018 g, 0.056 mmol) was added and the mixture was stirred overnight. Trifluoroacetic acid (0.65 ml) was added and after a couple of hours the hydrolysis was complete according to LC-MS. The
15 solvent was removed under reduced pressure and the residue was purified by preparative HPLC on a Kromasil C8- column using a gradient of 5-100% MeCN in 0.15% trifluoroacetic acid buffer as eluent. The solvent was removed under reduced pressure and 0.020 g (77 %) of the title product was obtained. *M/z* 655.11.

20 Method 9

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-L-methionine

25 A mixture of 3-(*R*)-4-(*R*)-1-(4-Fluorophenyl)-3-[(4-fluorobenzoyl)methylthio]-4-{4-[*N*-(carboxymethyl) carbamoylmethoxy]phenyl}azetidin-2-one (0.0197 g, 0.036 mmol), *tert*-butyl L-methioninate hydrochloride (0.0144 g, 0.060 mmol) and *N*-methylmorpholine (0.012 ml, 0.111 mmol) in DCM (2ml) was stirred at room temperature. TBTU (0.018 g, 0.056 mmol) was added and the mixture was stirred overnight. Trifluoroacetic acid (0.65 ml) was
30 added and after a couple of hours the hydrolysis was complete according to LC-MS. The solvent was removed under reduced pressure and the residue was purified by preparative HPLC on a Kromasil C8- column using a gradient of 5-100% MeCN in 0.15% trifluoroacetic acid buffer as eluent. The solvent was removed under reduced pressure and 0.015 g (61 %) of the title product was obtained. *M/z* 672.10.

Method 10

tert-butyl *N*-[(benzyloxy)carbonyl]glycyl-D-valinate

40 A mixture of *N*-[(benzyloxy)carbonyl]glycine (, 2.4g, 11.5 mmol), *tert*-butyl D-valinate hydrochloride (2.4 g, 11.4 mmol) and *N*-methylmorpholine (2.53 ml, 22.9 mmol) in DCM (20ml) was stirred at room temperature. TBTU (4.79 g, 14.9 mmol) was added and the mixture was stirred for three days. The solvent was removed under reduced pressure. Water
45 was added and the mixture was extracted two times with toluen. The organic layer was washed with brine, dried (Na₂SO₄), filtered and concentrated. The crude product was purified by flash chromatography using DCM:EtOAc:aceton 4:1:1 as eluent to give 3.92 g (94%) of the title compound. NMR (500 MHz, CD₃COOD) 0.88-0.99 (m, 6H), 1.48 (s, 9H), 2.08-2.19 (m 1H), 3.85 (ABq, 2H), 4.24 (d, 1H), 5.12 (ABq, 2H), 7.28-7.41 (m, 5H).

50

Method 11

tert-butyl glycyl-D-valinate hydrochloride

5
10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
1000

tert-Butyl N-[(benzyloxy)carbonyl]glycyl-D-valinate (3.89g, 10.7 mmol) and Pd on charcoal (95%, 0.3g) were mixed in EtOH (95%, 80 ml) and stirred under H₂-atmosphere for 2 h. The mixture was filtered through Celite 521 and the solvent was evaporated under reduced pressure. MeCN (25 ml) and pyridine hydrochloride (1.25g, 10.8 mmol) were added. The solvent was evaporated under reduced pressure to give 2.3g (81%) of the title product. NMR (500 MHz, CD₃COOD) 0.96-1.01 (m, 6H), 1.49 (s, 9H), 2.13-2.23 (m 1H), 3.76 (AB, 2H), 4.28-4.33 (m, 1H).

Method 12

N-[[4-((2R,3R)-1-(4-fluorophenyl)-3-[[2-(4-methoxyphenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl]glycyl-D-valine

20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
1000

A mixture of [4-((2R,3R)-1-(4-fluorophenyl)-3-[[2-(4-methoxyphenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetic acid (0.0153g, 0.031 mmol), tert-butyl glycyl-D-valinate hydrochloride (0.0099 g, 0.037 mmol) and N-methylmorpholine (0.010 ml, 0.091 mmol) in DCM (2ml) was stirred at room temperature. TBTU (0.016 g, 0.050 mmol) was added and the mixture was stirred for 3.5 h. Trifluoroacetic acid (0.5 ml) was added and after 3.5 h the solvent was removed under reduced pressure. The residue was purified by preparative HPLC on a Kromasil C8- column using a gradient of 5-100% MeCN in 0.15% trifluoroacetic acid buffer as eluent. The solvent was removed under reduced pressure and 0.015 g (74 %) of the title product was obtained. M/z 652.20.

Method 13

N-[[4-((2R,3R)-1-(4-fluorophenyl)-3-[[2-(4-methoxyphenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl]glycine

35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
1000

A mixture of [4-((2R,3R)-1-(4-fluorophenyl)-3-[[2-(4-methoxyphenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetic acid (0.150g, 0.30 mmol), tert-butyl glycinate hydrochloride (0.0635 g, 0.38 mmol) and N-methylmorpholine (0.10 ml, 0.91 mmol) in DCM (2ml) was stirred at room temperature. TBTU (0.128 g, 0.40 mmol) was added and the mixture was stirred overnight. Trifluoroacetic acid (4.0 ml) was added and after 2 h the solvent was removed under reduced pressure. The residue was purified by preparative HPLC on a Kromasil C8- column using 35% MeCN in 0.1M ammonium acetate buffer as eluent. The solvent was removed under reduced pressure and 0.159 g (95 %) of the title product was obtained. M/z 553.02.

Method 14

***N*-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-2-butylnorleucine**

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (0.020 g, 0.037 mmol) and NMM (0.040 ml, 0.363 mmol) were dissolved in DCM (5 ml) at 30°C. TBTU (a total of 0.016 g, 0.050 mmol) were added in portions and the mixture was stirred for 1 h. 2-Butylnorleucine (0.007 g, 0.037 mmol) was added and the mixture was stirred at 30°C for 18 h. The reaction mixture was concentrated under reduced pressure and the residue was purified by preparative HPLC, using a gradient of 20-50% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, 0.009 g (34 % yield) of the title product was obtained as a white solid. *M/z*: 710.1. ¹H NMR (DMSO, 400 MHz): δ 0.73-0.82 (m, 6H), 0.88-1.22 (m, 8H), 1.56-1.69 (m, 2H), 1.96-2.07 (m, 2H), 3.71 (d, 2H), 4.32 (d, 1H), 4.36 (ABq, 2H), 4.52 (s, 2H), 5.16 (d, 1H), 6.95-7.01 (m, 2H), 7.11-7.26 (m, 4H), 7.30-7.40 (m, 4H), 7.61 (s, 1H), 7.98-8.06 (m, 2H), 8.25-8.42 (m, 1H).

Method 15

***N*-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-L-alanine**

A solution of [4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetic acid (0.020 g, 0.041 mmol), L-Alanine tert-butyl ester hydrochloride (0.009 g, 0.050 mmol) and N-Methylmorpholine (0.018 ml, 0.163 mmol) in DCM (4 ml) was stirred for 5 min. TBTU (0.017 g, 0.053 mmol) was added. The formation of the ester was confirmed after 3 h. *M/z*: 611.1. TFA (3 ml) was added. After 2h, the mixture was diluted with toluene (2 ml) and the solvent was removed under reduced pressure. The residue was purified by preparative HPLC, using a gradient of 20-50% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, 0.023 g (>98%) of the title product was obtained as a white solid. *M/z*: 555.1. ¹H NMR (DMSO, 400 MHz): δ 1.17 (d, 3H), 3.73-3.82 (m, 1H), 4.33 (d, 1H), 4.35 (ABq, 2H), 4.43 (s, 2H), 5.15 (d, 1H), 6.92-7.98 (m, 2H), 7.10-7.24 (m, 4H), 7.29-7.39 (m, 4H), 7.84 (d, 1H), 7.97-8.04 (m, 2H).

Method 16

***N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-L-phenylalanine**

Methyl *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-L-phenylalaninate (20 mg, 0.029 mmol) was dissolved in 1.6 ml MeOH and 0.2 ml H₂O. Et₃N (0.2 ml, 1.44 mmol) was added and the mixture was stirred overnight. The mixture was heated to 80°C for 7 h. The mixture was purified by preparative HPLC using a C8 column (25x300mm). A gradient from 20% to 40% MeCN in 0.1 M ammonium acetate was used as mobile phase. The product fraction was concentrated and lyophilized to yield 7 mg (36%). *M/z*: 688.

Method 17

methyl *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-L-phenylalaninate

5

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (66 mg, 0.12 mmol) was dissolved in 2 ml DCM. *N*-methylmorpholine (40 μ l, 0.36 mmol), L-(*S*)-phenylalanine methyl ester hydrochloride (33 mg, 0.15 mmol) and finally TBTU (45 mg, 0.14 mmol) were added. The mixture was stirred overnight. The crude mixture was purified by flash chromatography on 5g SiO₂. EtOAc:Hex (1:1), DCM and finally DCM:Acetone (4:1) were used as eluents. The collected fraction was concentrated to yield 67mg (78%) of the title compound. *M/z*: 702 (*M*+1).

10

Method 18

15

***tert*-butyl *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-4-methylleucinate**

20

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine (20 mg, 0.037 mmol) was dissolved in 3 ml DCM. *N*-Methylmorpholine (9 μ l, 0.082 mmol) and TBTU (14 mg, 0.044 mmol) were added. After 5 minutes, *tert*-butyl 4-methylleucinate (9 mg, 0.045 mmol) was added and the mixture was stirred for 2 h. Additional *tert*-butyl 4-methylleucinate (ca 3 mg, 0.015 mmol) was added. After 15 min, water (2 ml) was added and the mixture was acidified to a pH of 2 using 2M KHSO₄. The aqueous phase was extracted with 2 ml DCM and the combined organic phases were washed with 3 ml water, dried over Na₂SO₄ and filtered. Removal of the solvent under reduced pressure gave the title compound. *M/z*: 724.

25

30

Method 19

***N*-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycylglycine**

35

A solution of of [4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetic acid (0.200 g, 0.414 mmol), glycylglycine methyl ester hydrochloride (0.090 g, 0.493 mmol) and *N*-methylmorpholine (0.150 ml) in DCM (5 ml) was stirred for 10 min. TBTU (0.170 g) was added and the mixture was stirred for 20 h. The formation of the ester was confirmed. *M/z*: 612.0. The solvent was removed under reduced pressure. The residue was dissolved in a mixture of MeOH (5 ml), water (1 ml) and Et₃N (0.5 ml). The solution was stirred at 50°C for 18 h. DBN (0.050 ml, 0.405 mmol) was added and the mixture was stirred for 2 h at 50°C. Ammonium acetate buffer (0.1 M, 3 ml) was added and the mixture was concentrated. The residue was purified by preparative HPLC, using a gradient of 20-50% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, the title product (0.094 g, 38 % yield) was obtained as a white solid. *M/z*: 598.2. ¹H NMR (DMSO, 400 MHz): 3.50 (d, 2H), 3.75 (d, 2H), 4.32 (d, 1H), 4.35 (ABq, 2H), 4.46-4.53 (m,

40

45

2H), 5.15 (d, 1H), 6.94-7.00 (m, 2H), 7.10-7.25 (m, 4H), 7.29-7.39 (m, 4H), 7.68-7.81 (m, 1H), 7.98-8.04 (m, 2H), 8.30-8.36 (m, 1H).

5 Method 20

Ethyl {[(5,5-dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl]thio}acetate

10

Ethyl [(2-oxo-2-phenylethyl)thio]acetate (10.8 g, 45.3 mmol) was dissolved in toluene (250 ml). 2,2-Dimethyl-1,3-propanediol (37.6 g, 0.36 mol) and p-toluene sulfonic acid (cat., 500 mg) were added. The mixture was stirred at reflux in a Dean-Stark apparatus for two hours and at room temperature overnight. The mixture was concentrated under reduced pressure.

15 The crude oil was purified by flash-chromatography (hexane:EtOAc-7:1) to give 11.2 g (70 %) of the title compound as a colourless oil. ¹H-NMR (CDCl₃, 200 MHz): δ 0.6 (s, 3H), 1.2-1.3 (t, 3H), 1.4 (s, 3H), 3.0 (s, 2H), 3.2 (s, 2H), 3.5 (s, 4H), 4.1-4.2 (q, 2H), 7.3-7.6 (m, 5H).

Method 21

20

{[(5,5-Dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl]thio}acetic acid

25 Ethyl {[(5,5-dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl]thio}acetate (11.2 g, 34.3 mmol) was dissolved in THF (150 ml) and cooled to 0°C. LiOH (2.88 g, 68.7 mmol) in water (40 ml) was added and the mixture was stirred for 19 h. The solvents were evaporated. The crude product was extracted between water and diethyl ether. The aqueous layer was acidified to a pH of 6 using 2M HCl and extracted twice with CH₂Cl₂. The combined CH₂Cl₂ layers were dried

30 (Na₂SO₄) and concentrated under reduced pressure to give 9.8 g (96 %) of the title compound as a white solid. ¹H-NMR (CDCl₃, 200 MHz): δ 0.6 (s, 3H), 1.4 (s, 3H), 3.0 (s, 2H), 3.3 (s, 2H), 3.5 (s, 4H), 7.3-7.6 (m, 5H).

35 Method 22

(4S)-3-({[(5,5-Dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl]thio}acetyl)-4-phenyl-1,3-oxazolidin-2-one

40

{[(5,5-Dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl]thio}acetic acid (9.9 g, 33.0 mmol) was dissolved in dry CH₂Cl₂ (250 ml) and cooled to 0°C. N,N'-Dicyclohexylcarbodiimide (DCC, 7.63 g, 37.0 mmol) and 4-(dimethylamino)pyridine (DMAP, 8.57 g, 70.0 mmol) were added and the mixture was stirred at 0°C for 20 minutes. (S)-(+)-4-Phenyl-2-oxazolidinone (5.38 g,

45 33.0 mmol) was added and the mixture was stirred at room temperature for 70 h. The mixture was filtered, concentrated under reduced pressure and purified by flash-chromatography (hexane:EtOAc-7:3). This afforded 10.2 g (70 %) of the title compound as a white solid. ¹H-NMR (CDCl₃, 200 MHz): 0.6 (s, 3H), 1.3 (s, 3H), 2.8 (s, 2H), 3.4 (s, 4H), 3.8 (s, 2H), 4.1-4.15 (dd, 1H), 4.6-4.8 (t, 1H), 5.35-5.45 (dd, 1H), 7.25-7.45 (m, 9H).

50

Method 23**5 Ethyl 4-((1*S*,2*R*)-1-anilino-2-(((5,5-dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl)thio)-3-oxo-3-[(4*S*)-2-oxo-4-phenyl-1,3-oxazolidin-3-yl]propyl)phenoxy)acetate**

10 Tetraisopropyl orthotitanate (0.5 ml, 1.7 mmol) was added to a solution of TiCl_4 (1M in CH_2Cl_2 , 5.1 ml, 5.1 mmol) in CH_2Cl_2 (50 ml) at 0°C under inert atmosphere. The mixture was stirred for ten minutes. (4*S*)-3-(((5,5-Dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl)thio)acetyl)-4-phenyl-1,3-oxazolidin-2-one (3.0 g, 6.8 mmol) in dry CH_2Cl_2 (60 ml) was added dropwise over 20 minutes and the mixture was stirred for ten minutes. Ethyl {4-[(phenylimino)methyl]phenoxy}acetate (3.8 g, 13.6 mmol) in dry CH_2Cl_2 (60 ml) was
15 added dropwise over 30 minutes and the mixture was cooled to -40°C and stirred for 20 minutes. Ethyl diisopropyl amine (2.3 ml, 13.6 mmol) was added dropwise over ten minutes and the mixture was stirred at -40°C for six hours. The mixture was cooled to -78°C and isopropanol (90 ml) was added. The mixture was slowly warmed to room temperature overnight. H_2O (100 ml) was added and the mixture was stirred for 35 minutes at room
20 temperature. NH_4Cl (10%) was added and the mixture was extracted twice with diethyl ether. The combined organic layers were washed with water, dried (MgSO_4) and concentrated under reduced pressure. Purification by flash-chromatography (hexane:EtOAc 5:1 then 7:3 then 6:4) gave 2.13 g (43 %) of the title compound as a yellow solid. $^1\text{H-NMR}$ (CDCl_3 , 200 MHz): δ 0.6 (s, 3H), 1.2-1.4 (m, 6H), 2.8-3.0 (m, 2H), 3.3-3.5 (m, 4H), 4.5-4.7 (m, 3H), 5.3-5.5 (m,
25 1H), 5.7-5.8 (d, 1H), 6.4-6.5 (d, 2H), 6.6-6.9 (m, 4H), 6.9-7.0 (d, 2H), 7.0-7.5 (m, 7H). M/z : 747.3 ($\text{M}^+ + \text{Na}$)

Method 24

30

Ethyl 4-((2*R*,3*R*)-3-(((5,5-dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl)thio)-4-oxo-1-phenylazetididin-2-yl)phenoxy]acetate

35

Ethyl 4-((1*S*,2*R*)-1-anilino-2-(((5,5-dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl)thio)-3-oxo-3-[(4*S*)-2-oxo-4-phenyl-1,3-oxazolidin-3-yl]propyl)phenoxy)acetate (2.1 g, 2.9 mmol) was dissolved in dry toluene (200 ml) and heated to 90°C under inert atmosphere. N,O -Bis(trimethylsilyl)acetamide (BSA, 2.1 ml, 8.7 mmol) was added and the mixture was stirred
40 at 90°C for one hour. At 45°C tetrabutylammonium fluoride (TBAF, cat., 0.1 g) was added and the mixture was stirred at 45°C for 18 h. The mixture was concentrated under reduced pressure. The residue was purified by flash-chromatography (hexane:EtOAc 5:1). This afforded 0.98 g (60 %) of the title compound as a yellow oil. $^1\text{H-NMR}$ (CDCl_3 , 200 MHz): 0.6 (s, 3H), 1.2-1.4 (m, 6H), 3.0-3.2 (t, broad, 2H), 3.3-3.5 (m, 4H), 3.95 (d, 1H), 4.2-4.4 (q,
45 2H), 4.6 (s, 2H), 4.8 (d, 1H), 6.9-7.1 (m, 3H), 7.2-7.6 (m, 11H). MS (CI) M/z : 584.2 ($\text{M}^+ + \text{Na}$).

Method 25

50

***tert*-Butyl (4-((1*S*,2*R*)-2-(((5,5-dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl)thio)-1-((4-fluorophenyl)amino)-3-oxo-3-((4*S*)-2-oxo-4-phenyl-1,3-oxazolidin-3-yl)propyl)phenoxy)acetate**

- 5 Tetraisopropyl orthotitanate (0.51 ml, 1.8 mmol) was added to a solution of TiCl₄ (1M in CH₂Cl₂, 5.1 ml, 5.1 mmol) in CH₂Cl₂ (50 ml) at 0°C under inert atmosphere. The mixture was stirred for ten minutes. (4*S*)-3-(((5,5-Dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl)thio)acetyl)-4-phenyl-1,3-oxazolidin-2-one (3.0 g, 6.8 mmol) in dry CH₂Cl₂ (50 ml) was added dropwise over 20 minutes. After ten minutes, *tert*-butyl (4-((4-fluorophenyl)imino)methyl)phenoxy)acetate (4.5 g, 13.6 mmol) in dry CH₂Cl₂ (50 ml) was added dropwise over 30 minutes. The mixture cooled to -30°C and stirred for 20 minutes. Ethyl diisopropyl amine (2.3 ml, 13.4 mmol) in 20 ml dry CH₂Cl₂ was added dropwise over ten minutes and the mixture was stirred at -30°C for six hours. The mixture was cooled to -78°C. Isopropanol (60 ml) was added and the temperature was risen overnight. H₂O (100 ml)
- 15 was added and the mixture was stirred for 35 minutes at room temperature. The mixture was extracted twice with diethyl ether. The combined organic layers were washed with water, dried (MgSO₄) and concentrated under reduced pressure. Purification by flash-chromatography (hexane:EtOAc 5:1) afforded 2.95 g (56 %) of the title compound as a yellow solid. M/z: 793.3 (M⁺ + Na).

20

Method 26

- 25 ***tert*-Butyl {4-((2*R*,3*R*)-3-(((5,5-dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl)thio)-1-(4-fluorophenyl)-4-oxoazetidin-2-yl)phenoxy}acetate**

- 30 *tert*-Butyl (4-((1*S*,2*R*)-2-(((5,5-dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl)thio)-1-((4-fluorophenyl)amino)-3-oxo-3-((4*S*)-2-oxo-4-phenyl-1,3-oxazolidin-3-yl)propyl)phenoxy)acetate (2.6 g, 3.4 mmol) was dissolved in dry toluene (250 ml) and heated to 90°C under inert atmosphere. N,O-Bis(trimethylsilyl)acetamide (BSA, 2.5 ml, 10.3 mmol) was added and the mixture was stirred at 90°C for one hour. The mixture was cooled to 45°C and tetrabutylammonium fluoride (TBAF, cat., 0.5 g) was added. The mixture was stirred at 45°C for two hours. The mixture was concentrated under reduced pressure and purified by flash-chromatography (hexane:EtOAc 7:1). This afforded 0.65 g (31 %) of the
- 35 title compound as a yellow solid. ¹H-NMR (CDCl₃, 200 MHz): δ 0.6 (s, 3H), 1.2 (s, 3H), 1.5 (s, 9H), 2.9-3.1 (q, broad, 2H), 3.4 (q, 4H), 4.0 (s, 1H), 4.5 (s, 2H), 4.8 (s, 1H), 6.9-7.0 (m, 4H), 7.2-7.3 (m, 5H), 4.3-4.4 (m, 4H).

40 Method 27

(4-((2*R*,3*R*)-1-(4-Fluorophenyl)-4-oxo-3-((2-oxo-2-phenylethyl)thio)azetidin-2-yl)phenoxy)acetic acid

- 45 *tert*-butyl {4-((2*R*,3*R*)-3-(((5,5-dimethyl-2-phenyl-1,3-dioxan-2-yl)methyl)thio)-1-(4-fluorophenyl)-4-oxoazetidin-2-yl)phenoxy}acetate (1.34 g, 2.21 mmol) was dissolved in formic acid (20 ml) and stirred for 90 minutes. The mixture was concentrated under reduced pressure (temperature < 30°C). The crude oil was purified by flash-chromatography (hexane:acetone:formic acid 60:40:0.1) to afford 0.7 g (68 %) of the title compound as a pale
- 50 yellow solid. ¹H-NMR (CDCl₃, 200 MHz): δ 4.15 (d, 1H), 4.22 (d, 2H), 4.71 (s, 2H), 4.91 (d,

1H), 6.92-7.00 (m, 4H), 7.24-7.30 (m, 4H), 7.46-7.63 (m, 3H), 7.94-7.99 (d, 2H). MS (CI)
M/z: 464.2 (M-1).

5 Method 28

Ethyl ({[2-(4-methoxyphenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl}thio)acetate

10 Ethyl {[2-(4-methoxyphenyl)-2-oxoethyl]thio}acetate (9.57 g, 35.7 mmol) was dissolved in benzene (250 ml). 2,2-Dimethyl-1,3-propanediol (29.7 g, 0.29 mol) and p-toluene sulfonic acid (500 mg) were added. The mixture was stirred at reflux in a Dean-Stark apparatus for three hours and at room temperature overnight. The mixture was concentrated under reduced pressure. CH₂Cl₂ was added and the solution was washed twice with H₂O and with brine. The organic phase was dried (MgSO₄) and filtered. Concentration under reduced pressure afforded
15 a crude oil which was purified by flash-chromatography (hexane:EtOAc 7:1) to afford 5.95 g (47 %) of the title compound as a colorless oil. ¹H-NMR (CDCl₃, 200 MHz): δ 0.6 (s, 3H), 1.2-1.4 (m, 6H), 3.0 (s, 2H), 3.2 (s, 2H), 3.4 (s, 4H), 3.8 (s, 3H), 4.2 (q, 2H), 6.9 (d, 2H), 7.4 (d, 2H).

20

Method 29

{[2-(4-Methoxyphenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl}thio)acetic acid

25 Ethyl ({[2-(4-methoxyphenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl}thio)acetate (5.95 g, 16.8 mmol) was dissolved in THF (75 ml) and cooled to 0°C. LiOH (2.11 g, 50.4 mmol) in water (40 ml) was added and the mixture was stirred for 90 minutes. Water was added and the mixture was extracted twice with diethyl ether. The aqueous layer was acidified using 2M HCl until pH=4 and extracted twice with CH₂Cl₂. The combined CH₂Cl₂ layers were dried
30 (Na₂SO₄) and concentrated under reduced pressure to afford 5.4 g (>98%) of the title compound as a white solid. ¹H-NMR (CDCl₃, 200 MHz): δ 0.6 (s, 3H), 1.4 (s, 3H), 3.0 (s, 2H), 3.4 (s, 2H), 3.5 (s, 4H), 3.9 (s, 3H), 6.9 (d, 2H), 7.4 (d, 2H).

35

Method 30

(4S)-3-[({[2-(4-Methoxyphenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl}thio)acetyl]-4-phenyl-1,3-oxazolidin-2-one

40 ({[2-(4-Methoxyphenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl}thio)acetic acid (5.4 g, 16.5 mmol) was dissolved in dry CH₂Cl₂ (80 ml) and cooled to 0°C. N,N'-dicyclohexylcarbodiimide (DCC, 3.76 g, 18.2 mmol) in 20 ml of CH₂Cl₂ and 4-(dimethylamino)pyridine (DMAP, 4.04 g, 33.1 mmol) were added and the mixture was stirred
45 at 0°C for 30 minutes. (S)-(+)-4-Phenyl-2-oxazolidinone (2.69 g, 16.5 mmol) was added and the mixture was stirred at room temperature for 17 h. The mixture was filtered, concentrated under reduced pressure and purified by flash-chromatography (hexane:EtOAc 4:1 then 2:1). This afforded 5.69 g (73 %) of the title compound as a white solid. ¹H-NMR (CDCl₃, 200 MHz): δ 0.6 (s, 3H), 1.3 (s, 3H), 2.8 (s, 2H), 3.4 (s, 4H), 3.8 (s, 2H), 3.8 (s, 3H), 4.3 (dd, 1H),
50 4.7 (t, 1H), 5.4 (dd, 1H), 6.9 (d, 2H), 7.3 (m, 7H).

Method 31

- 5 *tert*-Butyl (4-((1*S*,2*R*)-1-[(4-fluorophenyl)amino]-2-([2-(4-methoxyphenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl)thio)-3-oxo-3-[(4*S*)-2-oxo-4-phenyl-1,3-oxazolidin-3-yl]propyl)phenoxy)acetate

- 10 Tetraisopropyl orthotitanate (0.31 ml, 1.06 mmol) was added to a solution of TiCl₄ (1M in CH₂Cl₂, 3.18 ml, 3.18 mmol) in CH₂Cl₂ (50 ml) at 0°C under inert atmosphere. The mixture was stirred for ten minutes. (4*S*)-3-([2-(4-Methoxyphenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl)thio)acetyl]-4-phenyl-1,3-oxazolidin-2-one (2.00 g, 4.24 mmol) in dry CH₂Cl₂ (50 ml) was added dropwise over 20 minutes. The mixture was stirred for ten minutes. *tert*-Butyl (4-((4-fluorophenyl)imino)methyl)phenoxy)acetate (2.79 g, 8.48 mmol) in dry CH₂Cl₂ (50 ml) was added dropwise over 30 minutes. The mixture was cooled to -30°C and stirred for 20 minutes. Ethyl diisopropyl amine (1.45 ml, 8.48 mmol) in 10 ml dry CH₂Cl₂ was added dropwise over ten minutes and the mixture was stirred at -30°C for five hours. The mixture was cooled to -78°C. Isopropanol (60 ml) was added and the temperature was allowed to rise overnight. H₂O (100 ml) was added and the mixture was stirred for 20 minutes and extracted twice with diethyl ether. The combined organic layers were washed with water, dried (MgSO₄) and concentrated under reduced pressure. Purification by flash-chromatography (hexane:EtOAc 3:1) afforded 2.00 g (59 %) of the title compound as a yellow solid. ¹H-NMR (CDCl₃, 200 MHz): δ 0.6 (s, 3H), 1.3 (s, 3H), 1.5 (s, 9H), 2.6-2.9 (m, 2H), 3.3-3.5 (m, 4H), 3.8 (s, 3H), 4.1-4.3 (m, 2H), 4.5 (s, 2H), 4.6-4.8 (m, 2H), 5.0-5.4 (s, broad, 1H), 5.4 (m, 1H), 5.7 (d, 1H), 6.4 (m, 2H), 6.6-6.8 (m, 4H), 6.8-7.0 (m, 4H), 7.1-7.4 (m, 7H).
- 25

Method 32

- 30 *tert*-Butyl {4-[(2*R*,3*R*)-1-(4-fluorophenyl)-3-([2-(4-methoxyphenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl)thio)-4-oxoazetidin-2-yl]phenoxy}acetate

- tert*-Butyl (4-((1*S*,2*R*)-1-[(4-fluorophenyl)amino]-2-([2-(4-methoxyphenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl)thio)-3-oxo-3-[(4*S*)-2-oxo-4-phenyl-1,3-oxazolidin-3-yl]propyl)phenoxy)acetate (2.0 g, 2.5 mmol) was dissolved in dry toluene (200 ml) and heated to 90°C under inert atmosphere. N,O-Bis(trimethylsilyl)acetamide (BSA, 1.8 ml, 7.5 mmol) was added and the mixture was stirred at 90°C for one hour. The mixture was cooled to 45°C and tetrabutylammonium fluoride (TBAF, 150 mg) was added and the mixture was stirred at 45°C for two hours. The mixture was concentrated under reduced pressure and filtered through a short column of silica (hexane:EtOAc 4:1). Purification of the crude oil by flash-chromatography (hexane:EtOAc 5:1) gave 0.65 g (41 %) of the title compound as a yellow solid. ¹H-NMR (CDCl₃, 200 MHz): δ 0.6 (s, 3H), 1.3 (s, 3H), 1.5 (s, 9H), 3.0-3.2 (m, 2H), 3.3-3.5 (m, 4H), 3.8 (s, 3H), 4.0 (d, 1H), 4.5 (s, 2H), 4.8 (d, 1H), 6.8-7.0 (m, 6H), 7.2-7.4 (m, 6H).
- 40
- 45

Method 33

- 50 [4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-([2-(4-methoxyphenyl)-2-oxoethyl]thio)-4-oxoazetidin-2-yl]phenoxy]acetic acid

tert-Butyl {4-[(2*R*,3*R*)-1-(4-fluorophenyl)-3-([2-(4-methoxyphenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl)thio]-4-oxoazetidin-2-yl]phenoxy}acetate (0.65 g, 1.02 mmol) was dissolved in formic acid (10 ml) and stirred for 90 minutes. The mixture was concentrated under reduced pressure (temperature < 30°C) and the crude oil was purified by flash-chromatography (hexane:acetone:formic acid 60:40:0.1) to afford 0.45 g (88 %) of the title compound as a pale yellow solid. ¹H-NMR (CDCl₃, 200 MHz): δ 3.9 (s, 3H), 4.1 (d, 1H), 4.1 (s, 2H), 4.7 (s, 2H), 4.9 (d, 1H), 6.9-7.1 (m, 6H), 7.2-7.4 (m, 4H), 7.9-8.0 (d, 2H). MS (CI) M/z: 494.1 (M⁺ - 1), 495.1 (M⁺)

Method 34

Ethyl ({[5,5-dimethyl-2-(4-methylphenyl)-1,3-dioxan-2-yl]methyl}thio)acetate

Ethyl {[2-(4-methylphenyl)-2-oxoethyl]thio}acetate (12.2 g, 48.4 mmol) was dissolved in benzene (350 ml). 2,2-Dimethyl-1,3-propanediol (40.3 g, 0.387 mol) and *p*-toluene sulfonic acid (1g) were added. The mixture was stirred at reflux in a Dean-Stark apparatus for two hours, cooled and concentrated under reduced pressure. CH₂Cl₂ was added and the organic phase was washed twice with brine and dried (MgSO₄). Filtration and concentration under reduced pressure afforded a crude oil, which was purified by flash-chromatography (hexane:EtOAc 8:1) to afford 10.0 g (61 %) of the title compound as a clear oil. ¹H-NMR (CDCl₃, 200 MHz): δ 0.6 (s, 3H), 1.2-1.4 (m, 6H), 2.4 (s, 3H), 3.0 (s, 2H), 3.2 (s, 2H), 3.4-3.5 (m, 4H), 4.2 (q, 2H), 7.2-7.4 (m, 4H).

Method 35

([5,5-Dimethyl-2-(4-methylphenyl)-1,3-dioxan-2-yl]methyl)thio)acetic acid

Ethyl ({[5,5-dimethyl-2-(4-methylphenyl)-1,3-dioxan-2-yl]methyl}thio)acetate (10.1 g, 29.8 mmol) was dissolved in THF (150 ml) and cooled to 0°C. LiOH (3.76 g, 89.5 mmol) in water (50 ml) was added and the mixture was stirred for two hours at room temperature. The mixture was concentrated under reduced pressure. Water was added and the mixture was extracted with diethyl ether. The aqueous layer was acidified using 2M HCl to pH 3 and extracted twice with CH₂Cl₂. The combined CH₂Cl₂ layers were dried (Na₂SO₄) and concentrated under reduced pressure to afford 8.7 g (94 %) of the title compound as a white solid. ¹H-NMR (CDCl₃, 200 MHz): δ 0.6 (s, 3H), 1.2 (s, 3H), 2.4 (s, 3H), 3.0 (s, 2H), 3.4 (s, 2H), 3.4 (s, 4H), 7.2-7.4 (m, 4H).

Method 36

(4*S*)-3-[(5,5-Dimethyl-2-(4-methylphenyl)-1,3-dioxan-2-yl]methyl)thio)acetyl]-4-phenyl-1,3-oxazolidin-2-one

([5,5-Dimethyl-2-(4-methylphenyl)-1,3-dioxan-2-yl]methyl)thio)acetic acid (8.7 g, 28.0 mmol) was dissolved in dry CH₂Cl₂ (120 ml) and cooled to 0°C. N,N'-Dicyclohexylcarbodiimide (DCC, 6.35 g, 30.8 mmol) in 30 ml of CH₂Cl₂ and 4-

10 *tert*-Butyl (4-((1*S*,2*R*)-2-((5,5-dimethyl-2-(4-methylphenyl)-1,3-dioxan-2-yl)methyl)thio)-1-[(4-fluorophenyl)amino]-3-oxo-3-[(4*S*)-2-oxo-4-phenyl-1,3-oxazolidin-3-yl]propyl)phenoxy)acetate

Tetraisopropyl orthotitanate (0.37 ml, 1.23 mmol) was added to a solution of TiCl_4 (1M in CH_2Cl_2 , 3.3 ml, 3.3 mmol) in CH_2Cl_2 (50 ml) at 0°C under inert atmosphere. The mixture was stirred for ten minutes. (4*S*)-3-[($\{[5,5\text{-Dimethyl-2-(4-methylphenyl)-1,3-dioxan-2-yl]methyl\}$ thio)acetyl]-4-phenyl-1,3-oxazolidin-2-one (2.0 g, 4.4 mmol) in dry CH_2Cl_2 (50 ml) was added dropwise over 20 minutes and the mixture was stirred for ten minutes. *tert*-Butyl (4-[[4-(4-fluorophenyl)imino]methyl]phenoxy)acetate (2.9 g, 8.8 mmol) in dry CH_2Cl_2 (50 ml) was added dropwise over 30 minutes. The mixture was cooled to -30°C and stirred for 20 minutes. Ethyl diisopropyl amine (1.5 ml, 8.8 mmol) in 10 ml dry CH_2Cl_2 was added dropwise over ten minutes. The mixture was stirred at -30°C for four hours. The mixture was cooled to -78°C and isopropanol (60 ml) was added. The temperature was allowed to reach room temperature over two hours. H_2O (100 ml) was added and the mixture was stirred for 20 minutes and extracted twice with diethyl ether. The combined organic layers were washed with water, dried (MgSO_4) and concentrated under reduced pressure. Purification by flash-chromatography (hexane:EtOAc 5:1 then 4:1) afforded 2.55 g (74 %) of the title compound as a white solid. $^1\text{H-NMR}$ (CDCl_3 , 200 MHz): δ 0.6 (s, 3H), 1.3 (s, 3H), 1.5 (s, 9H), 2.4 (s, 3H), 2.6-2.9 (m, 2H), 3.3-3.5 (m, 4H), 4.1-4.3 (m, 1H), 4.5 (s, 2H), 4.6-4.8 (m, 2H), 5.0-5.3 (s, broad, 1H), 5.2 (m, 1H), 5.7 (d, 1H), 6.4 (m, 2H), 6.6-6.8 (m, 4H), 6.9 (m, 2H), 7.1-7.4 (m, 9H).

35

***tert*-Butyl {4-[(2*R*,3*R*)-3-({[5,5-dimethyl-2-(4-methylphenyl)-1,3-dioxan-2-yl]methyl}thio)-1-(4-fluorophenyl)-4-oxoazetidin-2-yl]phenoxy}acetate**

40 *tert*-Butyl (4-((1*S*,2*R*)-2-((5,5-dimethyl-2-(4-methylphenyl)-1,3-dioxan-2-yl)methyl)thio)-1-
[(4-fluorophenyl)amino]-3-oxo-3-[(4*S*)-2-oxo-4-phenyl-1,3-oxazolidin-3-
yl]propyl]phenoxy)acetate (2.55 g, 3.25 mmol) was dissolved in dry toluene (250 ml) and
heated to 90°C under inert atmosphere. N,O-Bis(trimethylsilyl)acetamide (BSA, 2.38 ml, 9.75
mmol) was added and the mixture was stirred at 90°C for one hour. The mixture was cooled
45 to 45°C and tetrabutylammonium fluoride (TBAF, 0.25 g) was added and the mixture was
stirred at 45°C for one hour. The mixture was concentrated under reduced pressure and
purified by flash-chromatography (hexane:EtOAc 6:1). This afforded 1.06 g (52 %) of the
title compound as a white solid. ¹H-NMR (CDCl₃, 200 MHz): δ 0.6 (s, 3H), 1.3 (s, 3H), 1.5

(s, 9H), 2.4 (s, 3H), 3.0-3.2 (m, 2H), 3.3-3.5 (m, 4H), 3.9 (d, 1H), 4.5 (s, 2H), 4.8 (d, 1H), 6.8-7.0 (m, 6H), 7.1-7.4 (m, 6H).

5 Method 39

[4-((2R,3R)-1-(4-Fluorophenyl)-3-[[2-(4-methylphenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetic acid

- 10 *tert*-Butyl {4-[(2R,3R)-3-([5,5-dimethyl-2-(4-methylphenyl)-1,3-dioxan-2-yl]methyl)thio]-1-(4-fluorophenyl)-4-oxoazetidin-2-yl]phenoxy}acetat (1.04 g, 1.67 mmol) was dissolved in formic acid (20 ml) and stirred for 60 minutes. The mixture was concentrated under reduced pressure (temperature < 30°C) and the crude oil was purified by flash-chromatography (hexane:acetone:formic acid 60:40:0.1) to afford 0.72 g (90 %) of the title compound as a pale yellow solid. ¹H-NMR (CDCl₃, 200 MHz): δ 2.4 (s, 3H), 4.1 (d, 1H), 4.2 (s, 2H), 4.7 (s, 2H), 4.9 (d, 1H), 6.9 (m, 6H), 7.2-7.4 (m, 4H), 7.8 (d, 2H). MS (CI) M/z: 478.1 (M⁺ - 1).

Method 40

- 20 *tert*-Butyl (4-[(*E*)-[(4-chlorophenyl)imino]methyl]phenoxy)acetate

A suspension of NaH (60 % in mineral oil, 3.69 g, 92.2 mmol) in dry DMF (70 ml) was cooled to 0°C. A solution of 4-hydroxybenzaldehyde (10.0 g, 82.0 mmol) in dry DMF (35 ml) was added dropwise. The mixture was stirred at 0°C for 40 minutes. *tert*-Butyl bromoacetate (12.1 ml, 82.5 mmol) was added and the mixture was stirred at room temperature for 17 h. The mixture was concentrated under reduced pressure. Diethyl ether was added and the mixture was washed with 10% NH₄Cl, water and brine. The organic phase was dried (MgSO₄), concentrated under reduced pressure and purified by flash-chromatography (10% - 20% EtOAc in hexane). This gave *tert*-butyl (4-formylphenoxy)acetate (17.4 g, 73.4 mmol, 90 % yield) as an colourless oil. This intermediate was dissolved in dry toluene (120 ml) and 4-chloroaniline (9.37 g, 73.4 mmol) was added. The mixture was refluxed in a Dean-Stark apparatus for 20 h, cooled and concentrated under reduced pressure. Hexane was added and the formed precipitate was filtered, washed twice with cold hexane and dried. This afforded 20.0 g (79%) of the title compound as a yellow solid. ¹H-NMR (CDCl₃, 200 MHz): δ 1.5 (s, 9H), 4.6 (s, 2H), 7.0 (d, 2H), 7.2 (d, 2H), 7.4 (d, 2H), 7.8 (d, 2H), 8.4 (s, 1H). MS (CI) M/z: 368.0 (M⁺+Na, 100), 369.0 (20), 370.0 (30), 371.0 (10).

Method 41

Ethyl ([2-(4-chlorophenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl)thio}acetate

- 45 Ethyl {[2-(4-chlorophenyl)-2-oxoethyl]thio}acetate (8.15 g, 29.9 mmol) was dissolved in toluene (165 ml). 2,2-Dimethyl-1,3-propanediol (24.8 g, 238 mmol) and *p*-toluene sulfonic acid (300 mg) were added. The mixture was stirred at reflux in a Dean-Stark apparatus for 2.5 h and concentrated under reduced pressure. The crude product was dissolved in CH₂Cl₂. The mixture was washed with H₂O (3x), brine, dried (Na₂SO₄) and concentrated under reduced pressure. The residue was purified by flash-chromatography (Hexane:EtOAc 9:1) to give 3.36

g (31 %) of the title compound as an yellow oil. MS (CI) M/z: 381.0 ($M^+ + Na$, 100), 382.0 (15), 383.0 (30), 384 (5)

5 Method 42

(([2-(4-Chlorophenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl}thio)acetic acid

Ethyl ([2-(4-chlorophenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl}thio)acetate (3.36 g, 9.36 mmol) was dissolved in THF (45 ml) and cooled to 0°C. LiOH (0.79 g, 18.8 mmol) in water (12 ml) was added and the mixture was stirred for 18 h at room temperature. The solvents were evaporated. The crude product was extracted between water and diethyl ether. The aqueous layer was acidified using 2M HCl to pH 6 and extracted twice with CH₂Cl₂. The combined CH₂Cl₂ layers were dried (Na₂SO₄) and concentrated under reduced pressure to afford 2.98 g (96 %) of the title compound as an yellow oil. ¹H-NMR (CDCl₃, 300 MHz): δ 0.62 (s, 3H), 1.34 (s, 3H), 2.95 (s, 2H), 3.34 (s, 2H), 3.45 (s, 4H), 7.4 (m, 4H)

Method 43

(4S)-3-[[[2-(4-Chlorophenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl}thio)acetyl]-4-phenyl-1,3-oxazolidin-2-one

(([2-(4-Chlorophenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl}thio)acetic acid (2.00 g, 6.05 mmol) was dissolved in dry CH₂Cl₂ (45 ml) and cooled to 0°C. N,N'-Dicyclohexylcarbodiimide (DCC, 1.43 g, 6.93 mmol) and 4-(dimethylamino)pyridine (DMAP, 1.70 g, 10.4 mmol) were added and the mixture was stirred at 0°C for 20 minutes. (S)-(+)-4-Phenyl-2-oxazolidinone (1.40 g, 11.5 mmol) was added and the mixture was stirred at room temperature for 70 h. The mixture was filtered, concentrated under reduced pressure and purified by flash-chromatography (20% - 30% EtOAc in hexane). This afforded 1.27 g (44 %) of the title compound as a white solid. ¹H-NMR (CDCl₃, 300 MHz): δ 0.62 (s, 3H), 1.34 (s, 3H), 2.78 (s, 2H), 3.42 (s, 4H), 3.85 (s, 2H), 4.27-4.33 (dd, 1H), 4.69-4.78 (t, 1H), 5.34-5.42 (dd, 1H), 7.29-7.38 (m, 9H).

Method 44

tert-Butyl (4-((1S,2R)-1-[(4-chlorophenyl)amino]-2-([2-(4-chlorophenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl}thio)-3-oxo-3-[(4S)-2-oxo-4-phenyl-1,3-oxazolidin-3-yl]propyl}phenoxy)acetate

Tetraisopropyl orthotitanate (0.23 ml, 0.77 mmol) was added to a solution of TiCl₄ (1M in CH₂Cl₂, 2.25 ml, 2.25 mmol) in CH₂Cl₂ (22 ml) at 0°C under inert atmosphere. The mixture was stirred for ten minutes. (4S)-3-[[[2-(4-Chlorophenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl}thio)acetyl]-4-phenyl-1,3-oxazolidin-2-one (1.44 g, 3.03 mmol) in dry CH₂Cl₂ (25 ml) was added dropwise over 45 minutes and the mixture was stirred for ten minutes. *tert*-Butyl (4-((*E*)-[(4-chlorophenyl)imino]methyl}phenoxy)acetate (2.09 g, 6.04 mmol) in dry CH₂Cl₂ (25 ml) was added dropwise over 80 minutes. The mixture was cooled to -40°C and stirred for 20 minutes. Ethyl diisopropyl amine (1.03 ml, 6.02 mmol) was added dropwise over 40 minutes and the mixture was stirred at -40°C for ten minutes. The mixture was cooled

to -78°C and isopropanol (80 ml) was added. The temperature was allowed to reach room temperature over three hours. NH₄Cl (10%, 80 ml) was added and the mixture was stirred for 35 minutes. Brine (200 ml) was added and the mixture was extracted twice with 400 ml diethyl ether. The combined organic layers were dried (MgSO₄) and concentrated under reduced pressure. Purification by flash-chromatography (10% - 20% EtOAc in hexane) afforded 1.69 g (65 %) of the title compound as a white solid. ¹H-NMR (CDCl₃, 200 MHz): δ 0.65 (s, 3H), 1.3 (s, 3H), 1.5 (s, 9H), 2.6-2.9 (m, 2H), 3.4 (m, 4H), 4.2 (m, 1H), 4.5 (s, 2H), 4.6-4.8 (m, 2H), 5.3 (m, 1H), 5.7 (d, 1H), 6.4 (m, 2H), 6.7-7.4 (m, 15H).

10

Method 45

tert-Butyl {4-[(2*R*,3*R*)-1-(4-chlorophenyl)-3-([2-(4-chlorophenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl)thio]-4-oxoazetidin-2-yl]phenoxy}acetate

15

tert-Butyl (4-{(1*S*,2*R*)-1-[(4-chlorophenyl)amino]-2-([2-(4-chlorophenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl)thio}-3-oxo-3-[(4*S*)-2-oxo-4-phenyl-1,3-oxazolidin-3-yl]propyl)phenoxy}acetate (1.69 g, 2.06 mmol) was dissolved in dry toluene (140 ml) and heated to 90°C under inert atmosphere. N,O-Bis(trimethylsilyl)acetamide (BSA, 1.48 ml, 6.05 mmol) was added and the mixture was stirred at 90°C for one hour. The mixture was cooled to 45°C and tetrabutylammonium fluoride (TBAF, 0.1 g) was added and the mixture was stirred at 45°C for one hour and at room temperature overnight. The mixture was concentrated under reduced pressure and purified by flash-chromatography (10% - 20% EtOAc in hexane). This afforded 0.84 g (61 %) of the title compound as a white solid. ¹H-NMR (CDCl₃, 200 MHz): δ 0.6 (s, 3H), 1.2 (s, 3H), 1.5 (s, 9H), 2.9-3.1 (t, broad, 2H), 3.4 (s, 4H), 4.0 (s, 1H), 4.5 (s, 2H), 4.8 (s, 1H), 6.9 (d, 2H), 7.2-7.4 (m, 10H).

30 Method 46

[4-((2*R*,3*R*)-1-(4-Chlorophenyl)-3-([2-(4-chlorophenyl)-2-oxoethyl]thio)-4-oxoazetidin-2-yl]phenoxy}acetic acid

tert-Butyl {4-[(2*R*,3*R*)-1-(4-chlorophenyl)-3-([2-(4-chlorophenyl)-5,5-dimethyl-1,3-dioxan-2-yl]methyl)thio]-4-oxoazetidin-2-yl]phenoxy}acetate (1.69 g, 2.57 mmol) was dissolved in formic acid (25 ml) and stirred for two hours. The mixture was concentrated under reduced pressure (temperature < 30°C) and the crude oil was purified by flash-chromatography (hexane:acetone:formic acid 60:40:0.1) to afford 1.08 g (81 %) of the title compound as a pale yellow solid. ¹H-NMR (CDCl₃, 200 MHz): δ 4.0-4.2 (m, 3H), 4.7 (s, 2H), 4.9 (d, 1H), 6.9 (d, 2H), 7.2-7.4 (m, 6H), 7.5 (d, 2H), 7.9 (d, 2H). MS (CI) M/z: 514.2 (M), 515.2 (30), 516.1 (70), 517.2 (20).

Method 47

45

Sodium (2*R*)-2-amino-4-cyclohexylbutanoate

Ethyl (2*R*)-2-amino-4-cyclohexylbutanoate hydrochloride (4.00 g, 18.75 mmol) was dissolved in MeOH/Water (10/5 ml). Sodium hydroxide (1.50 g, 37.50 mmol) was added. The reaction mixture was stirred for 1 h and concentrated to give the title compound as a colourless solid

(containing 100 mol% of sodium chloride). ^1H NMR $[(\text{CD}_3)_2\text{SO}]$, 400 MHz] δ 0.83-0.97 (m, 2H), 1.10-1.33 (m, 6H), 1.48-1.78 (m, 7H), 3.15 (dd, 1H).

5 Method 48

N-{[4-((2*R*,3*R*)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycine

- 10 [4-((2*R*,3*R*)-1-(4-Chlorophenyl)-3-{[2-(4-chlorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetic acid, 302.1 mg, 0.585 mmol) was dissolved in DCM (6 ml). *N*-Methylmorpholine (190 μl , 1.728 mmol) and *tert*-butyl glycinate hydrochloride (133.4 mg, 0.80 mmol) were added. After 10 minutes, TBTU (224.3 mg, 0.67 mmol) was added and the reaction mixture was stirred for 60 h. The intermediate *tert*-butylester of the title compound
- 15 was confirmed. M/z : 626.88 (M-H). DCM (10 ml) and water (15 ml) were added and the mixture was acidified to pH of 3 using KHSO_4 (2M). The organic phase was washed with water (2x15 ml). The combined aqueous phases were extracted with DCM (10 ml), dried over Na_2SO_4 , filtered and concentrated. A solution of the residue (500 mg) in DCM (10 ml) and TFA (4 ml) was stirred overnight. The solvent was removed under reduced pressure. Toluene
- 20 was added and evaporated to assist the removal of TFA. The residue was purified with preparative HPLC on a C8 column. A gradient from 20 to 50 % MeCN in 0.1M ammonium acetate buffer was used as eluent. After lyophilisation, the title compound was obtained as a white solid (166.9 mg, 50 %). ^1H -NMR (400 MHz, $\text{DMS}-d_6$): 3.51 (d, 2H), 4.33 (d, 1H), 4.34 (s, 1H), 4.36 (s, 1H), 4.47 (s, 2H), 5.17 (d, 1H), 6.96 (d, 2H), 7.16-7.21 (m, 2H), 7.35 (d, 4H),
- 25 7.54-7.59 (m, 2H), 7.83-7.90 (brs, 1H), 7.91-7.95 (m, 2H). M/z : 571.04 (M-H) and 572.88 (M+H).

Method 49

- 30 (2*R*)-cyclohexyl[(*N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-oxoethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl)amino]acetic acid

TBTU (0.0092 g, 0.029 mmol) was added to a mixture of 3-(*R*)-4-(*R*)-1-(4-Fluorophenyl)-3-[(4-fluorobenzoyl)methylthio]-4-{4-[*N*-(carboxymethyl) carbamoylmethoxy]phenyl}azetidin-2-one (0.016g, 0.030 mmol) and *N*-methylmorpholin (0.101 ml, 0.98 mmol) in DMF (2ml).

35 The mixture was stirred overnight under N_2 -atmosphere. Additional TBTU (0.0092 g, 0.029 mmol) was added and the mixture was stirred at 35°C for 2 h. (2*R*)-amino(cyclohexyl)acetic acid hydrochloride (0.0068 g, 0.035

mmol) was added. The mixture was stirred at 35°C for 2 h and at room temperature overnight.

- 40 The solvent was removed under reduced pressure and the residue was purified by preparative HPLC on a Kromasil C8- column using a gradient of 5-100% MeCN in 0.15% trifluoroacetic acid buffer as eluent. The solvent was removed under reduced pressure and 0.009 g (47 %) of the title product was obtained.

M/z 680.01

45

Examples of intermediates of formula (VI)

Method 50

***N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-alanine**

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-alanine was dissolved in methanol (1.5 ml). Sodium borohydride was added and the mixture was stirred for 30 minutes. An ammonium acetate/H₂O solution (2 ml) was added and the methanol was evaporated. The product was purified by preparative HPLC (CH₃CN/ 0.1 % ammoniumacetate buffer 20:80-100:0). The fractions containing product were lyophilized and 27 mg (48%) of the title product was obtained. *M/z*: 555.0 (*M*-1).

Method 51

***N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}-L-tryptophan**

[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-[[2-(4-fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetic acid (0.050 g, 0.103 mmol) was dissolved in CH₂Cl₂ (5 ml). Tryptophane tert-butyl ester hydrochloride (0.037g, 0.12 mmol) and *N*-methylmorpholine (31 mg, 0.31 mmol) were added. After 10 minutes, TBTU (43 mg, 0.13 mmol) was added and the mixture was stirred for 4 h. The crude ester was purified on silica gel and eluted with EtOAc/CH₂Cl₂, 25/75. The fractions containing pure ester were concentrated. CH₂Cl₂ (5 ml) and TFA (1 ml) were added and the reaction was stirred for 4 h. The mixture was concentrated and the remaining trace of TFA was azeotropically removed by co-evaporation with toluene (2 X 5 ml). The residue was dissolved in 5 ml of MeOH and sodium borohydride (0.016 g, 0.414 mmol) was added. The reaction was quenched by the addition of 0.1M NH₄OAc buffer (1 ml) after 5 minutes. The mixture was concentrated and purified by preparative HPLC (gradient 20-50% CH₃CN in 0.1M ammonium acetate buffer). Freeze-drying of the pure fractions gave the title compound as a colourless solid (0.040 g, 58%). *M/z*: 670.3 (*M*-1). ¹H NMR [(CD₃)₂SO], 400 MHz] δ 2.85-2.95 (m, 2H), 3.07-3.12 (m, 1H), 3.22-3.27 (m, 1H), 4.24-4.27 (m, 1H), 4.34-4.38 (m, 1H), 4.41 (s, 2H), 4.70-4.76 (m, 1H), 5.01-5.04 (m, 1H), 6.80-7.35 (m, 16H), 7.50-7.53 (m, 1H), 7.85-7.92 (m, 1H), 10.76 (s, 1H).

Method 52

***N*²-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}-L-glutamine**

[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetic acid (50 mg, 0.103 mmol), tert-butyl L-glutamate hydrochloride (30 mg, 0.124 mmol) and *N*-methylmorpholine (40 mg, 0.396 mmol) were dissolved in methylene chloride (1 ml). TBTU (40 mg, 0.125 mmol) was added and the mixture was stirred for 90 min. The solvent was evaporated and the residue was dissolved in formic acid (1 ml). The mixture was heated to 45-50 °C for 4 h. The reaction mixture was evaporated under reduced pressure. Toluene (5 ml) was added and evaporated. The residue was dissolved in methanol (1

ml). NaBH₄ (30 mg, 0.793 mmol) was added and the mixture was stirred for 15 min. Acetic acid (50 mg, 0.83 mmol) was added and the reaction mixture was evaporated under reduced pressure. The residue was purified by preparative HPLC using acetonitrile/ammonium acetate buffer (35:65) as eluent. After freeze-drying 47 mg (74%) of the title compound was

5 obtained. ¹H-NMR, 300 MHz, DMSO): 1.72-2.16 (m, 4H), 2.81-2.95 (m, 2H), 4.08-4.20 (m, 1H), 4.26-4.31 (m, 1H), 4.50 (s, 2H), 4.65-4.78 (m, 1H), 5.03-5.08 (m, 1H), 6.68 (s, 1H), 6.89-7.44 (m, 14H), 8.29 (d, 1H).

Method 53

10

N-{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}-D-serine

A solution of [4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-oxoethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetic acid (0.050 g, 0.103 mmol), O-(tert-butyl)-D-serine tert-butyl ester hydrochloride (0.032 g, 0.147 mmol) and N-methylmorpholine (0.035 ml, 0.318 mmol) in DCM (4 ml) was stirred for 5 min. TBTU (0.044 g, 0.137 mmol) was added. The formation of the ester was confirmed after 3 h. M/z: 683.1. TFA (2 ml) was added and the mixture was stirred for 22 h. The solvent was removed under reduced pressure. The residue was dissolved in MeOH (4 ml) and NaBH₄ (totally 0.130 g, 3.44 mmol) was added in small portions. The reaction was quenched by the addition of 0.1M ammonium acetate buffer (3 ml). The methanol was removed under reduced pressure. The remaining solution was purified by preparative HPLC using a gradient of 20-60% MeCN in 0.1M ammonium acetate buffer as eluent. After freeze-drying, 0.021 g (36 % yield) of the title product was obtained as a white solid. M/z: 573.1. ¹H NMR (DMSO, 400 MHz): δ 2.84-2.96 (m, 2H), 3.47 (dd, 1H), 3.69 (dd, 1H), 3.97-4.06 (m, 1H), 4.27-4.32 (m, 1H), 4.52 (ABq, 2H), 4.68-4.77 (m, 1H), 5.04-5.09 (m, 1H), 5.65 (bs, 1H), 6.99 (d, 2H), 7.07-7.41 (m, 10H), 7.89 (d, 1H).

30

Absorption

35 Absorption of the compounds of formula (I) was tested in a Caco-2 cells model (Gastroenterology 1989, 96, 736):

Compound (I)	Caco value (10 ⁻⁶ cm/sec)
<i>N</i> -{[4-((2 <i>R</i> ,3 <i>R</i>)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-valine	0.06
<i>N</i> -{[4-((2 <i>R</i> ,3 <i>R</i>)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-tyrosine	0.07
<i>N</i> -{[4-((2 <i>R</i> ,3 <i>R</i>)-1-(4-fluorophenyl)-3-[[2-(4-fluorophenyl)-2-hydroxyethyl]thio]-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-	0.2

dated 10/8/11/2004
101340

©BLS&L

04-07-2004
04-07-2004


- 110 -

lysine	
1-(4-Fluorophenyl)-3-(R)-[2-(4-fluorophenyl)-2-hydroxyethylthio]-4-(R)-{4-[N-{N-[2-(phenyl)-1-(R)-(carboxy)ethyl]carbamoylmethyl}carbamoylmethoxy]phenyl}azetidin-2-one	0.09

101340

Claims

1. A compound of formula (I):



5

(I)

wherein:

R^1 is hydrogen, C_{1-6} alkyl, C_{3-6} cycloalkyl or aryl; wherein said C_{1-6} alkyl may be optionally substituted by one or more hydroxy, amino, guanidino, carbamoyl, carboxy, C_{1-6} alkoxy, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, C_1 - C_6 alkylcarbonylamino

- 10 C_{1-6} alkylS(O) $_a$ wherein a is 0-2, C_{3-6} cycloalkyl or aryl; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C_{1-6} alkyl or C_{1-6} alkoxy;

R^2 is hydrogen, a branched or unbranched C_{1-6} alkyl, C_{3-6} cycloalkyl or aryl; wherein said C_{1-6} alkyl may be optionally substituted by one or more hydroxy, amino, guanidino,

- 15 carbamoyl, carboxy, C_{1-6} alkoxy, $(C_1-C_4)_3Si$, N -(C_{1-6} alkyl)amino, N,N -(C_{1-6} alkyl) $_2$ amino, C_{1-6} alkylS(O) $_a$ wherein a is 0-2, C_{3-6} cycloalkyl or aryl; and wherein any aryl group may be optionally substituted by one or two substituents selected from halo, hydroxy, C_{1-6} alkyl or C_{1-6} alkoxy;

R^3 is hydrogen, alkyl, halo or C_{1-6} alkoxy;

- 20 R^4 is hydrogen, halo or C_{1-6} alkoxy;

R^5 is hydrogen, branched or unbranched C_{1-6} alkyl, arylalkyl, or aryl C_{1-6} alkyl;

R^6 is hydrogen, C_{1-6} alkyl, or aryl C_{1-6} alkyl;

wherein R^5 and R^2 may form a ring with 2-7 carbon atoms and wherein R^6 and R^2 may form a ring with 3-6 carbon atoms;

- 25 or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof; with the proviso that said compound is not 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-2-hydroxyethylsulphanlyl]-4-[4-(N -(N -(R)-1-(carboxy)-2-(hydroxy)ethyl]carbamoylmethyl]

carbamoylmethoxy]phenyl]azetidin-2-one; or 3-(R)-4-(R)-1-(phenyl)-3-[2-(4-fluorophenyl)-2-hydroxyethylsulphanyl]-4-{4-[N-((R)- α -{N-[(S)-1-(carboxy)-2-(hydroxy)ethyl]carbamoyl}benzyl)carbamoylmethoxy]phenyl}azetidin-2-one.

5 2. A compound according to claim 1, wherein:

R^1 is hydrogen or phenyl.

3. A compound according to claim 1 or 2, wherein:

R^2 is hydrogen, a branched or unbranched C_{1-6} alkyl, C_{3-6} cycloalkyl or aryl; wherein said

10 C_{1-6} alkyl may be optionally substituted by one or more hydroxy, amino, acylamino, C_{1-6} alkylS(O)_a wherein a is 0-2, C_{3-6} cycloalkyl or aryl; and wherein any aryl group may be optionally substituted by hydroxy.

4. A compound according to any of the preceding claims, wherein:

15 R^3 is hydrogen, C_1-C_6 alkyl, halo or methoxy.

5. A compound according to any of the preceding claims, wherein:

R^3 is hydrogen, methyl, chlorine, fluorine or methoxy.

20 6. A compound according to any of the preceding claims, wherein:

R^4 is hydrogen or halo.

7. A compound according to any of the preceding claims, wherein:

R^4 is chlorine or fluorine.

25

8. A compound according to any of the preceding claims, wherein:

R^6 is hydrogen, aryl C_{1-6} or R^6 and R^2 form a ring with 3-6 carbon atoms.

9. A compound according to claim 1, wherein:

30 R^1 is hydrogen;

R^2 is a branched or unbranched C_{1-4} alkyl, optionally substituted by a C_{3-6} cycloalkyl;

R^3 and R^4 are halo;

R^5 is hydrogen or C_{1-6} alkyl; and

R^6 is hydrogen.

10. One or more compounds chosen from:

5 N -{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl- N^6 -acetyl-D-lysine;

10 1-(4-Fluorophenyl)-3-(*R*)-[2-(4-fluorophenyl)-2-hydroxyethylthio]-4-(*R*)-{4-[*N*-{*N*-[2-(phenyl)-1-(*R*)-(carboxy)ethyl]carbamoylmethyl}carbamoylmethoxy]phenyl}azetidin-2-one;

N -{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-valine;

15 N -{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-tyrosine;

N -{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-proline;

20 N -{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-lysine;

N -{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-hydroxy-2-(4-methoxyphenyl)ethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-valine;

25 N -{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-2-butylnorleucine;

30 N -{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-*S*-methyl-L-cysteine;

N -{[4-((2*R*,3*R*)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-cyclohexyl-D-alanine;

35 N -{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-cyclohexyl-D-alanine;

40 N -{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-4-methylleucine;

N -{[4-((2*R*,3*R*)-1-(4-Fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}-L-alanyl-D-valine;

45 N -{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-hydroxy-2-(4-methylphenyl)ethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-valine;

- 114 -

N-{[4-((2*R*,3*R*)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-D-valine;


5 *N*-{[4-((2*R*,3*R*)-1-(4-chlorophenyl)-3-{[2-(4-chlorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-methyl-D-valine;

N-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-(2-naphthyl)-D-alanine;

10 *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-3-methyl-D-valine;

15 *N*-{[4-((2*R*,3*R*)-1-(4-fluorophenyl)-3-{[2-(4-fluorophenyl)-2-hydroxyethyl]thio}-4-oxoazetidin-2-yl)phenoxy]acetyl}glycyl-(3*R*,4*S*,5*R*)-3,4,5,6-tetrahydroxy-D-norleucine;

101340

ABSTRACTTITLE : CHEMICAL COMPOUNDS

5 Compounds of formula (I):

(I)

(wherein variable groups are as defined within) pharmaceutically acceptable salts, solvates,
10 solvates of such salts and prodrugs thereof and their use as cholesterol absorption inhibitors
for the treatment of hyperlipidaemia are described. Processes for their manufacture and
pharmaceutical compositions containing them are also described.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record.**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.